

Jin Bai Kim

On singular matrices

*Czechoslovak Mathematical Journal*, Vol. 18 (1968), No. 2, 274–277

Persistent URL: <http://dml.cz/dmlcz/100831>

## Terms of use:

© Institute of Mathematics AS CR, 1968

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON SINGULAR MATRICES

JIN BAI KIM, Morgatown

(Received November 11, 1966)

1. PENROSE [7] discussed a generalized inverse for matrices, and he established the following theorem.

**Theorem A.** *For any matrix  $A$ , the four equations  $AXA = A$ ,  $XAX = X$ ,  $(AX)^* = AX$ , and  $(XA)^* = XA$  have a unique solution  $X$ , where  $A^*$  denotes the conjugate transpose of  $A$ .*

This unique solution  $X$  is called the generalized inverse of  $A$ . If we remove the third and fourth equations in Theorem A above, a solution  $X$  (of the equations  $XAX = X$  and  $AXA = A$ ) is not in general unique.

Then the natural question is:

**Problem.** *What is the cardinal number of the set of all solutions  $X$  of the equations  $AXA = A$  and  $XAX = X$  for a matrix  $A$  in the set  $M_n(F)$  of all  $n$  by  $n$  matrices over a field  $F$ ?*

The purpose of this note is to prove (Theorem 1) that if  $A \in M_n(F)$  then the cardinal number of the set of all solutions  $X$  of the equations  $AXA = A$  and  $XAX = X$  is equal to  $|F|^{2(\text{rank of } A)(n - (\text{rank of } A))}$ .

This result leads to a new definition in the class of regular semigroups (see Definition 2) and gives new examples of regular semigroups with zero (see Theorem 2).

2. Let  $F$  be a field.  $M_n(F)$  denotes the set of all  $n$  by  $n$  matrices over the field  $F$  with binary operation, the usual matrix multiplication. By Theorem A,  $M_n(F)$  is a regular semigroup. We define  $V(A) = \{X \in M_n(F) : AXA = A \text{ and } XAX = X\}$  which will be called an inverse set of  $A$  in  $M_n(F)$ .  $\varrho(A)$  denotes the rank of a matrix  $A$  in  $M_n(F)$ , and  $|T|$  denotes the cardinal number of a set  $T$ .

**Lemma 1.** *Let  $A \in M_n(F)$  and let  $X \in V(A)$ . Then  $\varrho(A) = \varrho(X)$ .*

**Proof.** From  $AXA = A$  and  $XAX = X$ ,  $\varrho(A) = \varrho(AXA) \leq \varrho(X) = \varrho(XAX) \leq \varrho(A)$  by Theorem 1.4 of [6, p. 83]; hence  $\varrho(A) = \varrho(X)$ .

**Lemma 2.** *The cardinal number of an inverse set  $V(A)$  of a matrix  $A$  in  $M_n(F)$  is invariant under elementary row or column operation on  $A$ , that is,  $|V(A)| = |V(EA)| = |V(AH)|$ , where  $E$  and  $H$  are elementary matrices (see Definition of elementary matrices on page 91 in [6]).*

*Proof.* Let  $A \in M_n(F)$  and let  $E$  be an elementary matrix in  $M_n(F)$ . Let  $X \in V(A)$  and let  $E^{-1}$  be the inverse matrix of the non-singular matrix  $E$ . Then  $EA = E(AXA) = EA(XE^{-1})EA$  and  $XE^{-1} = (XAX)E^{-1} = XE^{-1}(EA)XE^{-1}$ ; hence  $V(A)E^{-1} \subseteq V(EA)$  and  $|V(A)| \leq |V(EA)|$ . Similarly, we obtain  $V(EA)E \subseteq V(A)$  and  $|V(EA)| \leq |V(A)|$ . Thus  $|V(A)| = |V(EA)|$ . Analogously, we have  $|V(A)| = |V(AH)|$ , where  $H$  is an elementary matrix. This proves Lemma 2.

We need the following well known theorem.

**Theorem B.** *Every  $m$  by  $n$  matrix  $A$  is equivalent to a matrix  $C = (c_{ij})$  where  $c_{ii} = 1, i = 1, 2, \dots, \varrho(A)$ , and  $c_{ij} = 0$ , otherwise. The matrix  $C$  is called the canonical form of  $A$  (see Theorem 3.4 on page 106 in [6]).*

For  $1 \leq k \leq n$  let  $C_k = (d_{ij})$  where  $d_{ii} = 1$  for  $i = 1, 2, \dots, k$  and  $d_{ij} = 0$ , otherwise.

According to Lemma 2 and Theorem B, to solve the problem we need only consider  $C_k, k = 1, 2, \dots, n$ .

The main lemma follows.

**Lemma 3.** *Let  $k$  and  $n$  be positive integers with  $k \leq n$ . Let  $F_q$  be a Galois field with  $q$  elements. If  $C_k \in M_n(F_q)$ , then  $|V(C_k)| = q^{2k(n-k)} = q^{2(e(C_k))(n-e(C_k))}$ .*

*Proof.* Let  $k < n$ . Let  $X$  be an element of the inverse set  $V(C_k)$ . Then  $C_kXC_k = C_k$  and  $XC_kX = X$ . By direct calculation, it is not hard to see that  $X = (x_{ij})$  takes the form:

$$x_{ij} = \begin{cases} 1 & \text{if } i = j \text{ and } i = 1, 2, \dots, k; \\ 0 & \text{if } i \neq j \text{ and } \{i, j\} \subset \{1, 2, \dots, k\}; \\ x_{ij} & \text{if } i = 1, 2, \dots, k \text{ and } j = k + 1, k + 2, \dots, n; \\ x_{ij} & \text{if } i = k + 1, k + 2, \dots, n \text{ and } j = 1, 2, \dots, k; \\ \sum_{t=1}^k x_{it}x_{tj} & \text{if } \{i, j\} \subset \{k + 1, k + 2, \dots, n\}, \end{cases}$$

where  $x_{ij}$  above are arbitrary in  $F_q$ . Thus we are able to choose  $2k(n - k)$  entries of  $X$  arbitrary so that the cardinal number of the set  $V(C_k)$  is equal to  $q^{2k(n-k)}$ . If  $k = n$ , then  $V(C_n) = \{C_n\}$ , and  $|V(C_n)| = 1$ . This proves Lemma 3.

**Theorem 1.** *If  $A \in M_n(F)$ , then the cardinal number of the inverse set  $V(A)$  is equal to  $|F|^{2e(A)(n-e(A))}$ .*

*Proof* follows from Lemmas 2, 3 and Theorem B.

### 3. APPLICATIONS AND A QUESTION

**Definition 1.** A semigroup  $S$  with  $0$  is said to be homogeneous  $n$  regular if  $|V(a)| = n$  for every  $a \in S \setminus 0$  [4].

Let  $n$  and  $k$  be two positive integers with  $k \leq n$ . We define  $S_{n,k}(F) = \{X \in M_n(F) : \varrho(X) \leq k\}$ , and let  $S_{n,n-1}(F) = S_n(F)$ .

We have corollaries and Theorem 2.

**Corollary 1.**  $S_2(F_q)$  is a homogeneous  $q^2$  regular semigroup with  $0$ , where  $F_q$  is a finite field with  $q$  elements.

$S_3(F_q)$  is a homogeneous  $q^4$  regular semigroup with  $0$ .

$S_{n,1}(F_q)$  is a homogeneous  $q^{2(n-1)}$  regular semigroup with  $0$ .

**Corollary 2.** If  $F$  is a field of characteristic  $0$  then  $S_n(F)$  is a homogeneous  $\infty$  regular semigroup with  $0$ .

We have a new definition in the class of regular semigroups with  $0$ .

**Definition 2.** Let  $S$  be a regular semigroup with  $0$ .  $S$  is called a  $[s, t]$  regular semigroup with  $0$  if  $s \leq |V(a)| \leq t$  for every  $a \in S \setminus 0$ , where  $s$  and  $t$  are positive integers with  $s < t$ .

**Theorem 2.** Let  $F_q$  be a Galois field with  $q$  elements. Then  $S_n(F_q)$  is a  $[q^{2(n-1)}, q^{2\lfloor n/2 \rfloor(n - \lfloor n/2 \rfloor)}]$  regular semigroup with  $0$ , where

$$\lfloor n/2 \rfloor = \begin{cases} n/2 & \text{if } n \text{ is even,} \\ (n-1)/2 & \text{if } n \text{ is odd.} \end{cases}$$

In  $S_3(F_q)$ , there are two non-zero idempotents  $e = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$  and  $f = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

with  $ef = fe = e$  and  $e \neq f$ . Hence  $f$  is not a primitive idempotent of the homogeneous  $q^4$  regular semigroup  $S_3(F_q)$ . This example shows that the condition "every idempotent of  $S$  is primitive" is not necessary for a regular semigroup  $S$  with  $0$  to be homogeneous  $n$  regular (see Theorems 1, 3, 7 and 8 in [4]).

Hence we raise the following question:

**Question.** What are necessary and sufficient conditions for a regular semigroup  $S$  with  $0$  to be homogeneous  $n$  regular?

The author wishes to express his gratitude to Professor P. H. DOYLE of Michigan State University for his suggestions.

### References

- [1] *A. H. Clifford and G. B. Preston*: The algebraic theory of semigroups, Math. Surveys No. 7, Amer. Math. Soc., Providence, Providence, R. I., 1961.
- [2] *R. E. Cline*: Representations for the generalized inverse of a partitioned matrix, J. Soc. Indust. Appl. Math. vol. 12, no. 3 (1964), 588—600.
- [3] *Henry P. Decell, Jr.*: An alternative form of the generalized inverse of an arbitrary complex matrix, Siam Review, vol. 7 (1965), 356—357.
- [4] *Jin Bai Kim*: Completely 0-simple and homogeneous  $n$  regular semigroups, Bull. Amer. Math. Soc., vol. 71 (1965) 867—871.
- [5] *Jin Bai Kim*: Completely 0-simple and homogeneous  $n$  regular semigroups, Proceedings of the Amer. Math. Soc. (to appear), April issue of 1966.
- [6] *Marvin Marcus and Henry Minc*: Introduction to Linear algebra, The Macmillan Company, New York, 1965.
- [7] *R. Penrose*: A generalized inverse for matrices, Proc. Cambridge Phil. Soc., 51 (1955), 406—413.

*Author's address*: Morgantown, West Virginia, U.S.A. (West Virginia Universit.)