

Summaries of articles published in this issue

Czechoslovak Mathematical Journal, Vol. 19 (1969), No. 3, (569)–(570),(572)

Persistent URL: <http://dml.cz/dmlcz/100923>

Terms of use:

© Institute of Mathematics AS CR, 1969

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SUMMARIES OF ARTICLES PUBLISHED IN THIS ISSUE

(Publication of these summaries is permitted)

JOHN C. HIGGINS, Provo: *A faithful canonical representation for finitely generated N -semigroups*. Czech. Math. J. 19 (94), (1969), 375–379. (Original paper.)

A N -semigroup has been represented by Tamura as the Cartesian product of an abelian group and the additive positive integers with a special operation. This paper shows how, for finitely generated N -semigroups, one may obtain canonical representations in terms of distinguished elements of the semigroup. It also gives criteria for determining when two such representations represent isomorphic N -semigroups.

NORMAN NOBLE, Worcester: *Countably compact and pseudocompact product*. Czech. Math. J. 19 (94), (1969), 390–397. (Original paper.)

Following Frolík, we use \mathfrak{C} (resp. \mathfrak{P}) to denote the class of spaces X such that $X \times Y$ is countably compact (resp. pseudocompact) whenever Y is. In the first section we show that countably compact k -spaces (indeed, countably compact spaces in which each non- P -point is a k -point) are in \mathfrak{C} . Corresponding results for pseudocompact spaces are given in the second section and in the third section we prove that the class \mathfrak{P} is closed under arbitrary products.

ŠTEFAN SCHWABIK, Praha: *Stetige Abhängigkeit von einem Parameter und invariante Mannigfaltigkeiten für verallgemeinerte Differentialgleichungen*. Czech. Math. J. 19 (94), (1969), 398–427. (Originalartikel.)

Es werden verallgemeinerte Differentialgleichungen behandelt, welche unstetige Funktionen als Lösungen zulassen und mit deren Hilfe man Prozesse mit in gewissen Zeitpunkten wirkenden Impulsen beschreiben kann.

MIROSLAV FIEDLER and VLASTIMIL PTÁK, Praha: *Cyclic products and an inequality for determinants*. Czech. Math. J. 19 (94), (1969), 428–451. (Original paper.)

A new inequality for square matrices is introduced as follows: $A \stackrel{w}{\geq} B$ if every cyclic product of the matrix A is \geq than the corresponding cyclic product of B . This inequality is used to prove a generalization of Kotelanski's determinantal inequality and to discuss completely the case of equality.

Jiří VANŽURA, Praha: *Invariants of submanifolds*. Czech. Math. J. 19 (94), (1969), 452–468. (Original paper.)

The Author studies scalar geometric invariants of submanifolds of a given manifold, provided with the geometric structure by giving a sheaf of germs of vector fields on it. These invariants are generalization of those known from the classical geometries.

MARIE KOPÁČKOVÁ-SUCHÁ, Praha: *On the weakly nonlinear wave equation involving a small parameter at the highest derivative.* Czech. Math. J. 19 (94), (1969), 469–491. (Original paper.)

The paper deals with the equation $L_\varepsilon u \equiv \varepsilon u_{tt} - u_{xx} + 2au_t + cu = g(t, x) + \varepsilon f(t, x, u, u_x, u_t)$ where ε is a sufficiently small positive number, $t \geq 0$, $x \in E_1$. The existence and uniqueness of the solution of the Cauchy problem and of the periodic solution for this equation is proved. The behaviour of these solutions for $\varepsilon \rightarrow 0$ is studied.

ALOIS ŠVEC, Praha: *Submanifolds of Klein spaces.* Czech. Math. J. 19 (94), (1969), 492–499. (Original paper.)

It is shown that for the solution of the equivalence problem for submanifolds in Klein spaces only a finite process is needed.

IVO VRKOČ, Praha: *The class of functions fulfilling the inequality $\|f(x+z) - f(x) - f(y+z) + f(y)\| \leq \|x-y\| \omega(\|z\|)$.* Czech. Math. J. 19 (94), (1969), 500–514. (Original paper.)

In the article transformations $f: D \rightarrow B$, $D \subset A$ are investigated where A is a linear space with a semi-norm and B is a Banach space. Conditions are given for f to have the unique extension f^* on a set $L(D)$ and for the extension f^* to have the Gateaux differential at every point of D .

JIRÍ VANŽURA, Praha: *Remark on involutive subspaces and regular bases.* Czech. Math. J. 19 (94), (1969), 515–517. (Original paper.)

Given an involutive subspace $g^k \subset W \otimes S^k V^*$ and a subspace $V_1 \subset V$ of dimension r there is a regular basis v_1, \dots, v_n of a suitable prolongation g^{k_0} of g^k such that $v_1, \dots, v_r \in V_1$.

LADISLAV BICAN, Praha: *Some properties of completely decomposable torsion free abelian groups.* Czech. Math. J. 19 (94), (1969), 518–533. (Original paper.)

In this paper there are studied the conditions for the isomorphism $G \cong H$ where H is a completely decomposable torsion free abelian group and G a torsion free extension of H (subgroup of H , respectively). These conditions are formulated in terms of imbedding of H in G (G in H , respectively). As consequences some new criteria for complete decomposability of a torsion free abelian group are obtained.

JINDŘICH VEVERKA, Brno: *On couples of line congruences with the projective connection having Rozenfeld image with the character smaller than 5.* Czech. Math. J. 19 (94), (1969), 534–546. (Original paper.)

The Author studies the couples of non-parabolic line congruences in a three-dimensional space and in the space with the projective connection having Rozenfeld image in a five-dimensional projective space with a character smaller than five. Necessary and sufficient conditions for the character of Rozenfeld image to be smaller than five are found.

VLASTIMIL PTÁK, Praha: *Openness of linear mappings in LF-space*. Czech. Math. J. 19 (94), (1969), 547–552. (Original paper.)

Let E and F be two LF -spaces and T a sequentially open mapping of E into F . If we denote by R the range of T then R is sequentially closed in F . In the present paper we define a notion of orthogonality for subspaces R of spaces of type LF which describes the mutual position of R and the spaces of a defining sequence. Roughly speaking, R is said to be orthogonal in F provided R does not get too close, with increasing j , to the spaces of a defining sequence F_j (this is shown to be independent of the choice of a defining sequence). If T is sequentially open and if the range R of T is orthogonal in F then T is open.

VLASTIMIL PTÁK, Praha: *Simultaneous extensions of two functionals*. Czech. Math. J. 19 (94), (1969), 553–566. (Original paper.)

Given two subspaces R and Y of a Banach space F and two continuous linear functionals $r^* \in R'$ and $y^* \in Y'$ which coincide on $R \cap Y$, does there exist a continuous linear functional $x' \in F'$ which extends both r^* and y^* ? Clearly this is not always true. On the other hand, if the mutual position of R and Y is favourable, simultaneous extensions exist. There is also an intermediate situation where extensions do not exist, but the problem may be solved approximately: given $\varepsilon > 0$ there exists a $z' \in F'$ which is an extension of r^* and such that its restriction to Y differs from y^* less than ε in the norm of Y' . Necessary and sufficient conditions for each of these cases are given. In a forthcoming note they will be applied to the study of open mappings in spaces of type LF .

LADISLAV MIŠÍK, Bratislava: *Über f -durchschnittliche Eigenschaften*. Czech. Math. J. 19(94), (1969), 380–389. (Originalartikel.)

Hier ist die Frage für einige Mengeneigenschaften studiert, ob sie f -durchschnittlich oder nicht f -durchschnittlich sind.