

Bedřich Pondělíček

A certain equivalence on a semigroup

Czechoslovak Mathematical Journal, Vol. 21 (1971), No. 1, 109–117

Persistent URL: <http://dml.cz/dmlcz/101006>

Terms of use:

© Institute of Mathematics AS CR, 1971

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A CERTAIN EQUIVALENCE ON A SEMIGROUP

BEDŘICH PONDĚLÍČEK, Poděbrady

(Received November 5, 1969)

Let S be a periodic semigroup. We shall introduce the equivalence \bar{K} : for $a, b \in S$, $a\bar{K}b$ if and only if there exists an idempotent e and positive integers m, n such that $a^m = e = b^n$. In [1] J. T. Sedlock studies necessary and sufficient conditions on a periodic semigroup S in order that \bar{K} coincide with any one of the Green relations [2]. In this paper we consider arbitrary semigroups having similar properties.

I

In this section, S will be a fixed non-empty set. The mapping $\mathbf{U} : \exp S \rightarrow \exp S$ is said to be \mathcal{C} -closure operation if the mapping \mathbf{U} satisfies the following conditions:

- (1) $\mathbf{U}(\emptyset) = \emptyset$;
- (2) $A \subset B \subset S \Rightarrow \mathbf{U}(A) \subset \mathbf{U}(B)$;
- (3) $A \subset \mathbf{U}(A)$ for each $A \subset S$;
- (4) $\mathbf{U}(\mathbf{U}(A)) = \mathbf{U}(A)$ for each $A \subset S$.

For $x \in S$ we write simply $\mathbf{U}(x)$ instead of $\mathbf{U}(\{x\})$. The set of all \mathcal{C} -closure operations for a set S will be denoted by $\mathcal{C}(S)$.

A \mathcal{C} -closure operation \mathbf{U} is said to be \mathcal{Q} -closure operation if

$$(5) \quad \mathbf{U}\left(\bigcup_{i \in I} A_i\right) = \bigcup_{i \in I} \mathbf{U}(A_i) \quad \text{for } A_i \subset S \ (i \in I \neq \emptyset)$$

holds. Let $\mathcal{Q}(S)$ be the set of all \mathcal{Q} -closure operations for a set S . Evidently $\mathcal{Q}(S) \subset \mathcal{C}(S)$.

Let $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$, then we define

$$(6) \quad \mathbf{U} \leq \mathbf{V} \Leftrightarrow \mathbf{U}(A) \subset \mathbf{V}(A) \quad \text{for each } A \subset S .$$

The ordered set $\mathcal{C}(S)$ is a lattice. If $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$, then

$$(7) \quad (\mathbf{U} \wedge \mathbf{V})(A) = \mathbf{U}(A) \cap \mathbf{V}(A) \quad \text{for each } A \subset S.$$

If $\mathbf{U}, \mathbf{V} \in \mathcal{Q}(S)$, then

$$(8) \quad \mathbf{U} \vee \mathbf{V} \in \mathcal{Q}(S);$$

$$(9) \quad \mathbf{U} \leq \mathbf{V} \Leftrightarrow \mathbf{U}(x) \subset \mathbf{V}(x) \quad \text{for each } x \in S.$$

A subset A of S will be called **U-closed** if $\mathbf{U}(A) = A$. The set of all **U-closed** subsets of S will be denoted by $\mathcal{F}(\mathbf{U})$. If $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$, then

$$(10) \quad \mathcal{F}(\mathbf{U} \vee \mathbf{V}) = \mathcal{F}(\mathbf{U}) \cap \mathcal{F}(\mathbf{V});$$

$$(11) \quad \mathbf{U} \leq \mathbf{V} \Leftrightarrow \mathcal{F}(\mathbf{V}) \subset \mathcal{F}(\mathbf{U}).$$

Let $\mathbf{U} \in \mathcal{C}(S)$. We define $\mathbf{U}^* \in \mathcal{Q}(S)$. If $A \subset S$, then $x \in \mathbf{U}^*(A)$ if and only if $\mathbf{U}(x) \cap A \neq \emptyset$. For $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$ we have

$$(12) \quad \mathbf{U} \leq \mathbf{V} \Rightarrow \mathbf{U}^* \leq \mathbf{V}^*;$$

$$(13) \quad x \in \mathbf{U}(y) \Leftrightarrow y \in \mathbf{U}^*(x) \quad \text{for each } x, y \in S;$$

$$(14) \quad \mathbf{U}(x) = \mathbf{U}^{**}(x) \quad \text{for each } x \in S;$$

$$(15) \quad \mathbf{U} = \mathbf{U}^{**} \Leftrightarrow \mathbf{U} \in \mathcal{Q}(S).$$

(See [3].)

Definition 1. Let $\mathbf{U} \in \mathcal{C}(S)$. We shall introduce the equivalence $\bar{\mathbf{U}}$ on S by: for $x, y \in S$, $x\bar{\mathbf{U}}y$ if and only if $\mathbf{U}(x) = \mathbf{U}(y)$. For any element x of S , let \mathbf{U}_x denote the $\bar{\mathbf{U}}$ -class of S containing x .

Lemma 1. Let $\mathbf{U} \in \mathcal{C}(S)$. If $x, y \in S$, then $x\bar{\mathbf{U}}y$ if and only if $x \in \mathbf{U}(y)$ and $y \in \mathbf{U}(x)$.

Proof. If $x\bar{\mathbf{U}}y$, then by (3) $x \in \mathbf{U}(x) = \mathbf{U}(y)$ and $y \in \mathbf{U}(y) = \mathbf{U}(x)$. If $x \in \mathbf{U}(y)$ and $y \in \mathbf{U}(x)$, then by (2), (4) we have $\mathbf{U}(x) \subset \mathbf{U}(\mathbf{U}(y)) = \mathbf{U}(y)$. Similarly we obtain $\mathbf{U}(y) \subset \mathbf{U}(x)$. Thus $\mathbf{U}(x) = \mathbf{U}(y)$ and $x\bar{\mathbf{U}}y$.

Theorem 1. Let $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$. Then the following conditions are equivalent:

1. $\bar{\mathbf{U}} \subset \bar{\mathbf{V}}$;
2. for every $x \in S$, $\mathbf{U}_x \subset \mathbf{V}(x)$;
3. for every $A \in \mathcal{F}(\mathbf{V})$, $A = \bigcup_{x \in A} \mathbf{U}_x$.

Proof. 1 \Rightarrow 2. Let $x \in S$, then $\mathbf{U}_x \subset \mathbf{V}_x$. If $y \in \mathbf{U}_x$, then $y \in \mathbf{V}_x$. By Definition 1 and (3) we have $y \in \mathbf{V}(y) = \mathbf{V}(x)$. Thus $\mathbf{U}_x \subset \mathbf{V}(x)$.

2 \Rightarrow 3. If $x \in A \in \mathcal{F}(\mathbf{V})$, then $\mathbf{V}(x) \subset \mathbf{V}(A) = A$. Hence $\mathbf{U}_x \subset A$. This implies $A = \bigcup_{x \in A} \mathbf{U}_x$.

3 \Rightarrow 1. Let $x \overline{\mathbf{U}}y$. Evidently $\mathbf{V}(x) \in \mathcal{F}(\mathbf{V})$ and thus $y \in \mathbf{U}_y = \mathbf{U}_x \subset \mathbf{V}(x)$. Similarly we obtain $x \in \mathbf{U}_y \subset \mathbf{V}(y)$. From Lemma 1 it follows that $x \overline{\mathbf{V}}y$.

Corollary. *If $\mathbf{U} \in \mathcal{C}(S)$, then for every $A \in \mathcal{F}(\mathbf{U})$, $A = \bigcup_{x \in A} \mathbf{U}_x$.*

Theorem 2. *Let $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$. If $\mathbf{U} \leq \mathbf{V}$, then $\overline{\mathbf{U}} \subset \overline{\mathbf{V}}$.*

Proof. If $x \overline{\mathbf{U}}y$, then by Lemma 1 and (6) we have $x \in \mathbf{U}(y) \subset \mathbf{V}(y)$ and $y \in \mathbf{U}(x) \subset \mathbf{V}(x)$. It follows from Lemma 1 that $x \overline{\mathbf{V}}y$.

Theorem 3. *Let $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$, then $\overline{\mathbf{U} \wedge \mathbf{V}} = \overline{\mathbf{U}} \cap \overline{\mathbf{V}}$.*

Proof. It follows from Theorem 2 that $\overline{\mathbf{U} \wedge \mathbf{V}} \subset \overline{\mathbf{U}}$, $\overline{\mathbf{U} \wedge \mathbf{V}} \subset \overline{\mathbf{V}}$. This implies $\overline{\mathbf{U} \wedge \mathbf{V}} \subset \overline{\mathbf{U}} \cap \overline{\mathbf{V}}$. If $x(\overline{\mathbf{U}} \cap \overline{\mathbf{V}})y$, then $x \overline{\mathbf{U}}y$ and $x \overline{\mathbf{V}}y$. We have thus $\mathbf{U}(x) = \mathbf{U}(y)$ and $\mathbf{V}(x) = \mathbf{V}(y)$ so that $\mathbf{U}(x) \cap \mathbf{V}(x) = \mathbf{U}(y) \cap \mathbf{V}(y)$. By (7) we have $x \overline{\mathbf{U} \wedge \mathbf{V}}y$. Hence $\overline{\mathbf{U}} \cap \overline{\mathbf{V}} \subset \overline{\mathbf{U} \wedge \mathbf{V}}$ which implies $\overline{\mathbf{U} \wedge \mathbf{V}} = \overline{\mathbf{U}} \cap \overline{\mathbf{V}}$.

Theorem 4. *Let $\mathbf{U} \in \mathcal{Q}(S)$. Then the following conditions are equivalent:*

1. $\mathbf{U} = \mathbf{U}^*$;
2. for every $x \in S$, $\mathbf{U}(x) = \mathbf{U}_x$;
3. for every $x \in S$, $\mathbf{U}_x \in \mathcal{F}(\mathbf{U})$.

Proof. 1 \Rightarrow 2. It follows from Theorem 1 that $\mathbf{U}_x \subset \mathbf{U}(x)$ for every $x \in S$. Let $y \in \mathbf{U}(x)$. According to (13) we have $x \in \mathbf{U}^*(y) = \mathbf{U}(y)$. Since $\mathbf{U}(x) = \mathbf{U}(y)$, we have $y \in \mathbf{U}_x$, hence $\mathbf{U}(x) \subset \mathbf{U}_x$. This implies $\mathbf{U}(x) = \mathbf{U}_x$.

2 \Rightarrow 3. Evident.

3 \Rightarrow 1. It follows from (15) that $\mathbf{U} = \mathbf{U}^{**}$. Let $x \in S$. If $y \in \mathbf{U}^*(x)$, then by (13) $x \in \mathbf{U}(y)$. Since $y \in \mathbf{U}_y$, we have $\mathbf{U}(y) \subset \mathbf{U}(\mathbf{U}_y) = \mathbf{U}_y$ so that $x \in \mathbf{U}_y$. This implies $y \in \mathbf{U}(y) = \mathbf{U}(x)$ and $\mathbf{U}^*(x) \subset \mathbf{U}(x)$ for every $x \in S$. It follows from (9) that $\mathbf{U}^* \leq \mathbf{U}$. By (12) we have $\mathbf{U} = \mathbf{U}^{**} \leq \mathbf{U}^*$. Hence $\mathbf{U} = \mathbf{U}^*$.

Theorem 5. *Let $\mathbf{U}, \mathbf{V} \in \mathcal{C}(S)$. If $\mathbf{U} = \mathbf{U}^*$, then $\mathbf{U} \leq \mathbf{V}$ if and only if $\overline{\mathbf{U}} \subset \overline{\mathbf{V}}$.*

Proof. If $\mathbf{U} \leq \mathbf{V}$, then by Theorem 2 we have $\overline{\mathbf{U}} \subset \overline{\mathbf{V}}$. Suppose now that $\overline{\mathbf{U}} \subset \overline{\mathbf{V}}$. Evidently $\mathbf{U} = \mathbf{U}^* \in \mathcal{Q}(S)$. Let $A \subset S$. If $y \in \mathbf{U}(A)$, then by (5) we have $y \in \mathbf{U}(x)$ for some $x \in A$. According to Theorem 4, Theorem 1 and (2), we have $y \in \mathbf{U}_x \subset \mathbf{V}(x) \subset \mathbf{V}(A)$. This implies $\mathbf{U}(A) \subset \mathbf{V}(A)$. It follows from (6) that $\mathbf{U} \leq \mathbf{V}$.

II

Let now S be an arbitrary semigroup. Let $A \subset S$, $A \neq \emptyset$. Put $\mathbf{L}(A) = S^1A = SA \cup A$ and $\mathbf{R}(A) = AS^1 = AS \cup A$. Finally $\mathbf{L}(\emptyset) = \emptyset = \mathbf{R}(\emptyset)$. It is clear that $\mathbf{L}, \mathbf{R} \in \mathcal{Q}(S)$ and $\mathcal{F}(\mathbf{L})$ is the set of all left ideals of S (including \emptyset), $\mathcal{F}(\mathbf{R})$ is the set of all right ideals of S (including \emptyset). Put $\mathbf{M} = \mathbf{L} \vee \mathbf{R}$, $\mathbf{H} = \mathbf{L} \wedge \mathbf{R}$. Evidently $\mathbf{M} \in \mathcal{Q}(S)$ and $\mathbf{H} \in \mathcal{C}(S)$. It follows from (10) and (7) that $\mathcal{F}(\mathbf{M})$ is the set of all two-sided ideals of S (including \emptyset) and $\mathcal{F}(\mathbf{H})$ is the set of all quasi-ideals of S (including \emptyset).

Put $\mathbf{P}(\emptyset) = \emptyset$. If $A \subset S$, $A \neq \emptyset$, then by $\mathbf{P}(A)$ we denote the subsemigroup generated by all elements of A . Evidently $\mathbf{P} \in \mathcal{C}(S)$ and $\mathcal{F}(\mathbf{P})$ is the set of all subsemigroups of S (including \emptyset). Clearly $\mathbf{P} \leq \mathbf{H}$.

Lemma 2. *Let $A \subset S$. Then $A \in \mathcal{F}(\mathbf{P}^*)$ if and only if the implication*

$$(16) \quad x^n \in A \Rightarrow x \in A$$

holds for every $x \in S$ and for every positive integer n .

Proof. 1. Let $A \in \mathcal{F}(\mathbf{P}^*)$. If $x^n \in A$ for some $x \in S$ and for some positive integer n , then by (2) and (4) we have $\mathbf{P}^*(x^n) \subset A$. Since $x^n \in \mathbf{P}(x)$, it follows from (13) that $x \in \mathbf{P}^*(x^n) \subset A$.

2. Let (16) hold for every $x \in S$ and for every positive integer n . Evidently $\mathbf{P}^* \in \mathcal{Q}(S)$. If $A \neq \emptyset$, then by (5) we have $\mathbf{P}^*(A) = \bigcup_{x \in A} \mathbf{P}^*(x)$. If $y \in \mathbf{P}^*(A)$, then $y \in \mathbf{P}^*(x)$ for some $x \in A$. According to (13) $x \in \mathbf{P}(y)$ and thus $x = y^n$ for some positive integer n . Since $y^n \in A$, it follows from (16) that $y \in A$. Hence $\mathbf{P}^*(A) \subset A$ so that, by (3), $A = \mathbf{P}^*(A) \in \mathcal{F}(\mathbf{P}^*)$.

Lemma 3. *Let $A \subset S$. Then $A \in \mathcal{F}(\mathbf{P}^{**})$ if and only if the implication*

$$x \in A \Rightarrow x^n \in A$$

holds for every positive integer n .

Proof is analogous to the proof of Lemma 2.

Definition 2. Put $\mathbf{K} = \mathbf{P}^* \vee \mathbf{P}^{**}$.

Lemma 4. $\mathbf{K} = \mathbf{K}^*$.

Proof. According to (8) and (15), we have $\mathbf{K} = \mathbf{K}^{**}$. From $\mathbf{P}^* \leq \mathbf{K}$ and (12) we obtain $\mathbf{P}^{**} \leq \mathbf{K}^*$. It follows from $\mathbf{P}^{**} \leq \mathbf{K}$, (12) and (15) that $\mathbf{P}^* = \mathbf{P}^{***} \leq \mathbf{K}^*$. Thus $\mathbf{K} = \mathbf{P}^* \vee \mathbf{P}^{**} \leq \mathbf{K}^*$ and by (12) we have $\mathbf{K}^* \leq \mathbf{K}^{**} = \mathbf{K}$. This implies $\mathbf{K} = \mathbf{K}^*$.

Lemma 5. *If $x, y \in S$, then $x\bar{\mathbf{K}}y$ if and only if there exist positive integers n, m such that $x^n = y^m$.*

Proof. 1. Let $x^n = y^m$ for some positive integers n, m . By (14) and (6) we have $y^m \in \mathbf{P}(y) = \mathbf{P}^{**}(y) \subset \mathbf{K}(y)$. This and Lemma 2 implies that $x \in \mathbf{P}^*(x^n) = \mathbf{P}^*(y^m) \subset \mathbf{K}(y^m) \subset \mathbf{K}(y)$. Similarly we obtain $y \in \mathbf{K}(x)$ and thus by Lemma 1 we have $x\bar{\mathbf{K}}y$.

2. If $x\bar{\mathbf{K}}y$, then by Lemma 1 $x \in \mathbf{K}(y)$. Let $A = \{u/u^n = y^m \text{ for some positive integers } n, m\}$. It follows from Lemma 2, Lemma 3 and (10) that $A \in \mathcal{F}(\mathbf{P}^*) \cap \mathcal{F}(\mathbf{P}^{**}) = \mathcal{F}(\mathbf{P}^* \vee \mathbf{P}^{**}) = \mathcal{F}(\mathbf{K})$. Since $y \in A$, hence $x \in \mathbf{K}(y) \subset A$. We have thus $x^n = y^m$ for some positive integers n, m .

A semigroup S is called *right regular* (*left regular*) if $x \in x^2S$ ($x \in Sx^2$) for every $x \in S$.

Theorem 6. *The following conditions on a semigroup S are equivalent:*

1. S is right regular;
2. $\mathbf{P}^* \leq \mathbf{R}$;
3. $\mathbf{K} \leq \mathbf{R}$;
4. $\bar{\mathbf{K}} \subset \bar{\mathbf{R}}$.

Proof. $1 \Rightarrow 2$. Let S be a right regular semigroup. Let A be a right ideal of S . If $x^n \in A$ ($x \in S, n \geq 2$), then there exists $a \in S$ such that $x = x^2a$ and $x^{n-1} = x^na \in Aa \subset A$. Similarly we obtain $x^{n-i} \in A$ for any positive integer $i < n$. From here it follows that $x \in A$. By Lemma 2 we have $A \in \mathcal{F}(\mathbf{P}^*)$. It follows from (11) that $\mathbf{P}^* \leq \mathbf{R}$.

$2 \Rightarrow 3$. Suppose $\mathbf{P}^* \leq \mathbf{R}$. Evidently $\mathbf{P} \leq \mathbf{R}$. It follows from (12) and (15) that $\mathbf{P}^{**} \leq \mathbf{R}^{**} = \mathbf{R}$. Thus $\mathbf{K} = \mathbf{P}^* \vee \mathbf{P}^{**} \leq \mathbf{R}$.

$3 \Rightarrow 4$. This follows from Theorem 2.

$4 \Rightarrow 1$. If $\bar{\mathbf{K}} \subset \bar{\mathbf{R}}$, then by Lemma 4 and Theorem 5 we have $\mathbf{P}^* \leq \mathbf{K} \leq \mathbf{R}$. According to (11) $x^2 \in \mathbf{R}(x^2) \in \mathcal{F}(\mathbf{R}) \subset \mathcal{F}(\mathbf{P}^*)$. It follows from Lemma 2 that $x \in \mathbf{R}(x^2) = x^2S^1$. We shall show that $x \in x^2S$. Indeed, if $x = x^2$, then $x = x^3 \in x^2S$. Hence, S is right regular.

The following left-right dual of Theorem 6 is also true.

Theorem 7. *The following conditions on a semigroup S are equivalent:*

1. S is left regular;
2. $\mathbf{P}^* \leq \mathbf{L}$;
3. $\mathbf{K} \leq \mathbf{L}$;
4. $\bar{\mathbf{K}} \subset \bar{\mathbf{L}}$.

Theorem 8. *The following conditions on a semigroup S are equivalent:*

1. S is a union of groups;
2. S is left regular and right regular;
3. $\mathbf{P}^* \leq \mathbf{H}$;
4. $\mathbf{K} \leq \mathbf{H}$;
5. $\bar{\mathbf{K}} \subset \bar{\mathbf{H}}$.

Proof. $1 \Rightarrow 2$. Evident.

$2 \Rightarrow 3 \Rightarrow 4$. This follows from Theorem 6 and Theorem 7.

$4 \Rightarrow 5$. This follows from Theorem 2.

$5 \Rightarrow 1$. Suppose $\bar{K} \subset \bar{H}$. According to Theorem 3, Theorem 6 and Theorem 7, S is right regular and left regular. From here and (7) we obtain $x \in x^2S \cap Sx^2 \subset \mathbf{R}(x^2) \cap \mathbf{L}(x^2) = \mathbf{H}(x^2)$. On the other hand, we have $x^2 \in xS \cap Sx \subset \mathbf{R}(x) \cap \mathbf{L}(x) = \mathbf{H}(x)$. It follows from Lemma 1 and Theorem 3 that $x^2 \in \mathbf{H}_x = \mathbf{R}_x \cap \mathbf{L}_x$. According to [2] S is a union of groups.

A semigroup S is called *intraregular* if $x \in Sx^2S$ for every $x \in S$.

Theorem 9. *The following conditions on a semigroup S are equivalent:*

1. S is intraregular;
2. $\mathbf{P}^* \leq \mathbf{M}$;
3. $\mathbf{K} \leq \mathbf{M}$;
4. $\bar{\mathbf{K}} \subset \bar{\mathbf{M}}$.

(See [4].)

Proof. $1 \Rightarrow 2$. Let S be an intraregular semigroup. Let A be a two-sided ideal of S . If $x^n \in A$ ($x \in S$, $n \geq 2$), then there exist $a, b \in S$ such that $x^{n-1} = ax^{2(n-1)}b \in Sx^nS \subset SAS \subset A$. Similarly we obtain $x^{n-i} \in A$ for any positive integer $i < n$. This implies $x \in A$ and it follows from Lemma 2 that $A \in \mathcal{F}(\mathbf{P}^*)$ so that, by (11), $\mathbf{P}^* \leq \mathbf{M}$.

$2 \Rightarrow 3 \Rightarrow 4$. The proof is analogous to the proof of Theorem 6.

$4 \Rightarrow 1$. If $\bar{\mathbf{K}} \subset \bar{\mathbf{M}}$, then by Lemma 4 and Theorem 5 we have $\mathbf{P}^* \leq \mathbf{K} \leq \mathbf{M}$. It follows from (11) that $x^2 \in \mathbf{M}(x^2) \in \mathcal{F}(\mathbf{M}) \subset \mathcal{F}(\mathbf{P}^*)$. According to Lemma 2, $x \in \mathbf{M}(x^2) = S^1x^2S^1$. We shall prove that $x \in Sx^2S$. If $x \in Sx^2$, then $x = ax^2$ for some $a \in S$, thus $x = a(ax^2)x \in Sx^2S$. Similarly, $x \in x^2S$ implies $x \in Sx^2S$. If $x = x^2$, then $x = x^4 \in Sx^2S$. Hence, S is intraregular.

Remark 1. If S is a periodic semigroup, then from Corollary 2.3 [1], Theorem 3.8 [1] we have:

The conditions of Theorems 6, 7, 8 and 9 and the following condition on a periodic semigroup S are equivalent

$$\bar{\mathbf{K}} = \bar{\mathbf{H}}.$$

A semigroup S is called *left (right) weakly commutative* if for every $a, b \in S$ there exist $x \in S$ and a positive integer k such that $(ab)^k = bx$ ($(ab)^k = xa$).

Lemma 6. *If $\mathbf{L} \leq \mathbf{R}$, then a semigroup S is left weakly commutative.*

Proof. Let $a, b \in S$. By (6) we have $ab \in S^1b = \mathbf{L}(b) \subset \mathbf{R}(b) = bS^1$. If $ab = bx$ for some $x \in S$, then $(ab)^1 = bx$. If $ab = b$, then $(ab)^2 = b(ab)$. Hence, S is left weakly commutative.

Lemma 7. *If $\mathbf{R} \leq \mathbf{L}$, then the semigroup S is right weakly commutative.*

Lemma 8. *If S is a right regular and left weakly commutative semigroup, then $\mathbf{L} \leq \mathbf{R}$.*

Proof. Let $a \in S$. If $x \in Sa$, then $x = ua$ for some $u \in S$. Thus the hypothesis that S is left weakly commutative implies that there exists $v \in S$ and a positive integer k such that $x^k = (ua)^k = av \in aS \in \mathcal{F}(\mathbf{R})$. According to Theorem 6, (11) and Lemma 2, we have $x \in aS$. Hence $Sa \subset aS$. This shows that $\mathbf{L}(a) \subset \mathbf{R}(a)$ for every $a \in S$. Therefore, by (9), we have $\mathbf{L} \leq \mathbf{R}$.

Lemma 9. *If S is a left regular and right weakly commutative semigroup, then $\mathbf{R} \leq \mathbf{L}$.*

Theorem 10. *The following conditions on a semigroup S are equivalent:*

1. S is a semilattice of right groups;
2. S is a union of groups and $\mathbf{L} \leq \mathbf{R}$;
3. S is a union of groups and it is left weakly commutative;
4. $\mathbf{P}^* \leq \mathbf{L} \leq \mathbf{R}$;
5. $\mathbf{K} \leq \mathbf{L} \leq \mathbf{R}$;
6. $\overline{\mathbf{K}} \subset \overline{\mathbf{L}} \subset \overline{\mathbf{R}}$.

Proof. $1 \Rightarrow 2$. It follows from Theorem 2 [5] that S is a union of groups. Let $a \in S$. If $x \in Sa$, then $x = ua$ for some $u \in S$. Let e and f be an identity for a and for x , respectively. Similarly, let a^{-1} and x^{-1} be an inverse for a and for x , respectively. Since $x = ua = uae = xe$, hence $f = x^{-1}x = x^{-1}xe = fe$. By Theorem 2 [5] we have $f = efe$. Thus $ef = f$. Then $x = fx = efx = ex = (aa^{-1})x = a(a^{-1}x)$. This implies $x \in aS$. Consequently $Sa \subset aS$ and we have thus $\mathbf{L}(a) \subset \mathbf{R}(a)$. By (9) we obtain $\mathbf{L} \leq \mathbf{R}$.

$2 \Rightarrow 3$. This follows from Lemma 6.

$3 \Rightarrow 4$. This follows from Theorem 8 and Lemma 8.

$4 \Rightarrow 5$. This follows from Theorem 7.

$5 \Rightarrow 6$. This follows from Theorem 2.

$6 \Rightarrow 1$. It follows from Theorem 3 that $\overline{\mathbf{L}} = \overline{\mathbf{H}}$ and $\overline{\mathbf{K}} \subset \overline{\mathbf{H}}$. By Theorem 8, S is a union of groups. Let e and f be idempotents of S . Put $y = fe$. Let g and y^{-1} be an identity and an inverse for y , respectively. Since $y = yg = feg \in Seg$ and $eg = ey^{-1}y \in Sy$, hence $\mathbf{L}(y) = \mathbf{L}(eg)$. Now the hypothesis that $\overline{\mathbf{L}} \subset \overline{\mathbf{R}}$ implies $\mathbf{R}(y) = \mathbf{R}(eg)$. From this it follows that $y = eg$ or $y = egu$ for some $u \in S$. Then $y \in eS$ and therefore $efe = ey = y = fe$. It follows from Theorem 2 [5] that S is a semilattice of right groups.

Remark 2. The following example shows that the implication

$$\bar{L} \subset \bar{R} \Rightarrow L \leq R$$

on a semigroup S does not hold in general.

Let $S = \{(i, n - i) \mid \text{for all positive integers } n \text{ and for } i = 0, 1\}$. Define in S a multiplication by

$$xy = (i, n + m)$$

where $x = (i, n) \in S$ and $y = (j, m) \in S$. Then S is a semigroup (see [6]). It is clear that $\bar{L} \subset \bar{R}$. On the other hand, if $a = (1, 0)$, then $R(a) = aS \not\subseteq S = L(a)$ and thus $L \not\leq R$.

Remark 3. If S is a periodic semigroup, then from Theorem 3 and from Remark 1 we have:

The conditions of Theorem 10 and the following condition on a periodic semigroup S are equivalent:

$$\bar{K} = \bar{L}.$$

The dual statement reads as follows:

Theorem 11. *The following conditions on a semigroup S are equivalent:*

1. S is a semilattice of left groups;
2. S is a union of groups and $R \leq L$;
3. S is a union of groups and it is right weakly commutative;
4. $P^* \leq R \leq L$;
5. $K \leq R \leq L$;
6. $\bar{K} \subset \bar{R} \subset \bar{L}$.

Remark 4. *The conditions of Theorem 11 and the following condition on a periodic semigroup S are equivalent:*

$$\bar{K} = \bar{R}.$$

A semigroup S is called *weakly commutative* if for every $a, b \in S$ there exist $x, y \in S$ and a positive integer k such that

$$(ab)^k = xa = by.$$

Lemma 10. *A semigroup S is weakly commutative if and only if it is left weakly commutative and right weakly commutative.*

Proof. If S is a weakly commutative semigroup, then it is clear that S is left and right weakly commutative.

Suppose that S is left weakly commutative and right weakly commutative. Then there exist $x, y \in S$ and positive integers k, l such that

$$(ab)^k = xa, \quad (ab)^l = by.$$

This implies that $(ab)^{k+l} = ua = bv$ where $u = (ab)^l x, v = y(ab)^k$.

Theorem 12. *The following conditions on a semigroup S are equivalent:*

1. S is a semilattice of groups;
2. S is a union of groups and $\mathbf{L} = \mathbf{R}$;
3. S is a union of groups and it is weakly commutative;
4. $\mathbf{P}^* \leq \mathbf{L} = \mathbf{R}$;
5. $\mathbf{K} \leq \mathbf{L} = \mathbf{R}$;
6. $\overline{\mathbf{K}} \subset \overline{\mathbf{L}} = \overline{\mathbf{R}}$.

Proof follows from Theorem 2 [5], Corollary 2 [5], Theorem 10, Theorem 11 and Lemma 10.

Remark 5. *The conditions of Theorem 12 and the following conditions on a periodic semigroup S are equivalent:*

1. $\overline{\mathbf{K}} = \overline{\mathbf{L}} = \overline{\mathbf{R}}$;
2. $\overline{\mathbf{K}} = \overline{\mathbf{M}}$.

Proof. Conditions of Theorem 12 \Leftrightarrow 1. This follows from Theorem 12, Remark 3 and Remark 4.

1 \Rightarrow 2. It follows from Theorem 12 that $\mathbf{L} = \mathbf{R}$. Then $\mathbf{L} = \mathbf{M}$ and thus, by Remark 3, $\overline{\mathbf{K}} = \overline{\mathbf{L}} = \overline{\mathbf{M}}$.

2 \Rightarrow 1. According to Remark 1, we have $\overline{\mathbf{M}} = \overline{\mathbf{K}} = \overline{\mathbf{H}}$. It follows from Theorem 2 that $\overline{\mathbf{K}} = \overline{\mathbf{L}} = \overline{\mathbf{R}}$.

References

- [1] Sedlock J. T.: Green's relations on a periodic semigroup, Czech. Math. J., 19 (1969), 318–323.
- [2] Green J. A.: On the structure of semigroups, Annals of Math., 54 (1951), 163–172.
- [3] Pondělíček B.: On a certain relation for closure operations on a semigroup, Czech. Math. J., 20 (1970), 220–231.
- [4] Szász G.: Halbgruppen, deren Elemente durch Primideale trennbar sind, Acta Math. Ac. Sc. Hung., 19 (1968), 187–189.
- [5] Petrich M.: Semigroups certain of whose subsemigroups have identities, Czech. Math. J., 16 (1966), 186–198.
- [6] Pondělíček B.: Right prime ideals and maximal right ideals in semigroup, Mat. Časopis Slovensk. Akad. Vied (to appear).

Author's address: Poděbrady-zámek, ČSSR (České vysoké učení technické).