

James R. Wall

Green's relations for stochastic matrices

*Czechoslovak Mathematical Journal*, Vol. 25 (1975), No. 2, 247–260

Persistent URL: <http://dml.cz/dmlcz/101315>

## Terms of use:

© Institute of Mathematics AS CR, 1975

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

GREEN'S RELATIONS FOR STOCHASTIC MATRICES

J. R. WALL, Auburn

(Received December 13, 1973)

A real nonnegative  $m \times n$  matrix  $A$  is stochastic if the sum of all the entries in each row is 1. If  $A^T$ , the transpose of  $A$ , is also stochastic, then  $A$  is doubly stochastic. The set  $S_n[D_n]$  of all  $n \times n$  stochastic [doubly stochastic] matrices forms a compact Hausdorff semigroup under matrix multiplication with the  $n \times n$  identity matrix  $I$  as identity element.

An element  $a$  of a semigroup  $S$  is regular if  $axa = a$  for some  $x \in S$ . For  $A$  and  $B$  regular elements of  $S_n[D_n]$ , we obtain necessary and sufficient conditions for each of the systems

$$(1) \quad AX = B, \quad BY = A, \quad X, Y \in S_n[D_n]$$

and dually,

$$(2) \quad XA = B, \quad YB = A, \quad X, Y \in S_n[D_n]$$

to be consistent.

**1. Preliminaries, idempotents.** Certain definitions and results from the theory of semigroups will be useful. The algebraic facts summarized below may be found in [2, Chapter 2] and those requiring compactness in [6].

Let  $S$  be a semigroup with identity, and let  $a, b \in S$ . The relation  $\mathcal{R}[\mathcal{L}, \mathcal{J}]$  is defined on  $S$  by  $a \mathcal{R} b [a \mathcal{L} b, a \mathcal{J} b]$  if  $a$  and  $b$  generate the same principal right [left, two-sided] ideal of  $S$ . The relation  $\mathcal{D}$  is defined as the join of  $\mathcal{R}$  and  $\mathcal{L}$ , and the relation  $\mathcal{H}$  as the intersection of  $\mathcal{R}$  and  $\mathcal{L}$ . These relations are called Green's relations on  $S$ , and each is an equivalence relation on  $S$ .

Equivalently,  $a \mathcal{R} b [a \mathcal{L} b]$  if and only if  $ax = b$  and  $by = a [xa = b, yb = a]$  for some  $x, y \in S$ . Thus the problem of solving matrix equations (1) in  $S_n$  may be stated in two other ways: first, as the problem of characterizing  $\mathcal{R}$  on  $S_n$ ; and second, of finding all matrices  $B$  in  $S_n$  which lie in the same  $\mathcal{R}$ -equivalence class with a given  $A$  in  $S_n$ . Similar remarks hold for  $\mathcal{L}$  and for  $D_n$ .

If a  $\mathcal{D}$ -class  $D$  of a semigroup  $S$  contains one regular element, then every element of  $D$  is regular. Each  $\mathcal{R}$ -class and each  $\mathcal{L}$ -class of  $S$  contained in  $D$  contains an

idempotent. The  $\mathcal{H}$ -classes of  $S$  which contain an idempotent are just the maximal subgroups of  $S$ .

Since  $S_n[D_n]$  is compact,  $\mathcal{D} = \mathcal{J}$  on  $S_n[D_n]$ . Every compact semigroup  $S$  contains a unique minimal two-sided ideal called the kernel of  $S$ . The kernel of  $S_n[D_n]$  is described in the following theorem, which also identifies the idempotents in  $S_n[D_n]$  of rank 1.

**Theorem 1.1.** (SCHWARZ [10]) (a) *An idempotent element of  $S_n[D_n]$  has rank 1 if and only if it has constant columns. Every such idempotent is a right zero element of  $S_n$ , and  $S_n$  contains no left zeros. The only rank 1 idempotent in  $D_n$  is the matrix having each entry equal to  $1/n$ , and it is the zero element of  $D_n$ .*

(b) *The kernel  $K$  of the semigroup  $S_n[D_n]$  is the set of all rank 1 idempotents in  $S_n[D_n]$ . Each maximal subgroup contained in  $K$  contains only one element. Thus each  $\mathcal{L}$ -class of  $K$  is a singleton, and  $K$  contains only one  $\mathcal{R}$ -class.*

The next two results identify all idempotents in  $S_n[D_n]$ . In the sequel the term "canonical idempotent" in  $S_n[D_n]$  will refer to an idempotent which is expressed exactly in the form and with the notation of these results.

**Theorem 1.2.** (DOOB [3]) *Let  $E$  be an idempotent in  $S_n$  with rank  $k$ . Then there exists an  $n \times n$  permutation matrix  $P$  such that*

$$PEP^T = \begin{pmatrix} E_1 & 0 & \dots & 0 & 0 \\ 0 & E_2 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & E_k & 0 \\ F_1 & F_2 & \dots & F_k & 0 \end{pmatrix}$$

where each  $E_i$  is a positive  $n_i \times n_i$  stochastic matrix of rank 1; each  $F_i$  is an  $n_{k+1} \times n_i$  matrix of the form  $F_i = D_i E_i'$  with each  $F_i$  an  $n_{k+1} \times n_{k+1}$  nonnegative diagonal matrix such that  $D_1 + \dots + D_k = I$ , and each  $E_i'$  consists of  $n_{k+1}$  rows each equal to a row of  $E_i$ ;  $n_1 + \dots + n_{k+1} = n$ ;  $n_1 \geq \dots \geq n_k \geq n_{k+1} \geq 0$ ; and  $n_k \geq 1$ .

It is understood here and in similar situations in what follows that the columns of zeros on the right of  $PEP^T$  or the blocks  $F_1, \dots, F_k$  may not appear.

**Corollary 1.3.** (Doob [3], Schwarz [11]) *Let  $E$  be an idempotent in  $D_n$  with rank  $k$ . Then there exists an  $n \times n$  permutation matrix  $P$  such that*

$$PEP^T = E_1 \oplus E_2 \oplus \dots \oplus E_k$$

where each  $E_i$  is the  $n_i \times n_i$  matrix having each entry equal to  $1/n_i$ ;  $n_1 + \dots + n_k = n$ ; and  $n_1 \geq n_2 \geq \dots \geq n_k \geq 1$ .

**2. Green's relation  $\mathcal{R}$  for  $S_n$ .** We begin by characterizing those elements of  $S_n$  which lie in the same  $\mathcal{R}$ -class with a canonical idempotent  $E$ . The form of such a matrix  $A$  depends upon the way  $E$  is partitioned into blocks of rows, that is, upon the numbers  $n_i$ , and upon the diagonal matrices  $D_i$ . In the course of the proof, a method for constructing solutions to the system (1) will be indicated for the case where  $B$  is idempotent.

**Theorem 2.1.** *Let  $E$  be a canonical idempotent in  $S_n$  with rank  $k$ , and let  $A \in S_n$ . Then the following statements are equivalent.*

- (a)  $A \mathcal{R} E$ ; that is, the system  $AX = E, EY = A, X, Y \in S_n$ , is consistent.
- (b) There exists an  $n \times n$  permutation matrix  $P$  such that

$$(3) \quad AP = \begin{pmatrix} A_1 & 0 & \dots & 0 & 0 \\ 0 & A_2 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & A_k & 0 \\ U_1 & U_2 & \dots & U_k & 0 \end{pmatrix}$$

where each  $A_i$  is an  $n_i \times t_i$  stochastic matrix of rank 1, and  $U_i = D_i A'_i$  with each row of  $A'_i$  equal to a row of  $A_i$ .

- (c) The rows of  $A$  can be partitioned into the form

$$A = \begin{pmatrix} M_1 \\ \dots \\ M_k \\ M_{k+1} \end{pmatrix}$$

where each  $M_i$  is an  $n_i \times n$  stochastic matrix of rank 1; the matrices  $M_1, \dots, M_k$  are mutually orthogonal; and  $M_{k+1} = D_1 M'_1 + \dots + D_k M'_k$  where each row of  $M'_i$  is a row of  $M_i$ .

**Proof.** Assume (a) holds. Partition  $A$  by

$$A = \begin{pmatrix} M_1 \\ \dots \\ M_k \\ M_{k+1} \end{pmatrix}$$

where each  $M_i$  is  $n_i \times n$ . Since  $EA = A, E_i M_i = M_i$  for  $i = 1, \dots, k$ , and  $F_1 M_1 + \dots + F_k M_k = M_{k+1}$ . Thus  $\text{rank}(M_i) \leq \text{rank}(E_i) = 1$ , so  $\text{rank}(M_i) = 1$  for  $i = 1, \dots, k$ . Since  $M_i$  is stochastic, the columns of  $M_i$  are constant for  $i = 1, \dots, k$ .

Partition  $X$  by

$$X = (X_1, \dots, X_k, X_{k+1})$$

where each  $X_i$  is  $n \times n_i$ . Then  $AE = E$  implies

$$M_i X_i = E_i \quad \text{for } i = 1, \dots, k;$$

$$M_i X_j = 0 \quad \text{for } i \neq j, \quad i, j = 1, \dots, k;$$

$$M_i X_{k+1} = 0 \quad \text{for } i = 1, \dots, k + 1;$$

and

$$M_{k+1} X_j = F_j \quad \text{for } j = 1, \dots, k.$$

For  $k > 1$ , suppose  $M_i > 0$  for some  $i = 1, \dots, k$ . Then, for any  $j \neq i, j = 1, \dots, k$ ,  $M_i X_j = 0$ , so  $X_j = 0$ . But then  $E_j = M_j X_j = 0$ , contrary to  $E_j > 0$ . Therefore  $M_i$  has at least one column of zeros for  $i = 1, \dots, k$ . Let  $t_i$  be the number of nonzero columns of  $M_i$ ,  $i = 1, \dots, k$ . If  $k = 1$ , it may be that  $t_1 = n$ ; in this case  $A = M_1 = A_1$ .

Now assume  $t_1 < n$ . Let  $P_1$  be an  $n \times n$  permutation matrix such that the first  $t_1$  columns of  $M_1 P_1$  are nonzero. Then, for  $j = 2, \dots, k + 1$ ,  $0 = M_1 X_j = M_1 P_1 P_1^T X_j$ , so the first  $t_1$  rows of  $P_1^T X_j$  are zero. Thus

$$AP_1 = \begin{pmatrix} A_1 & 0 \\ A_{21} & A_{22} \end{pmatrix}$$

where  $A_1$  is an  $n_1 \times t_1$  stochastic matrix of rank 1, and

$$P_1^T X = \begin{pmatrix} X_{11} & 0 \\ X_{21} & X_{22} \end{pmatrix}$$

where  $X_{11}$  is a  $t_1 \times n_1$  stochastic matrix with

$$P_1^T X_1 = \begin{pmatrix} X_{11} \\ X_{21} \end{pmatrix}.$$

If  $k = 1$ , then  $AP_1$  has the form

$$AP_1 = \begin{pmatrix} A_1 & 0 \\ U_1 & 0 \end{pmatrix}.$$

Assume now that  $k > 1$ , and, for  $j = 2, \dots, k$ , let  $M_j P_1 = (B_j, C_j)$  where  $B_j$  is  $n_j \times t_1$ . Then

$$0 = M_j X_1 = B_j X_{11} + C_j X_{21}$$

so  $B_j X_{11} = 0$ . Thus, if  $B_j$  has a positive column, then the corresponding row of  $X_{11}$  is a zero row, which is impossible since  $X_{11}$  is stochastic. Hence  $B_j = 0$  for  $j = 2, \dots, k$ .

By repeating the process  $k$  times, we obtain an  $n \times n$  permutation matrix  $P = P_1 P_2 \dots P_k$  such that

$$AP = \begin{pmatrix} A_1 & 0 & \dots & 0 & 0 \\ 0 & A_2 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & A_k & 0 \\ U_1 & U_2 & \dots & U_k & U_{k+1} \end{pmatrix}$$

where each  $A_i$  is an  $n_i \times t_i$  stochastic matrix of rank 1. At each stage the matrix  $P_i$  is chosen so that it fixes the first  $t_1 + \dots + t_{i-1}$  columns of  $AP_1 \dots P_{i-1}$ . Also  $P^T X$  has the form

$$P^T X = \begin{pmatrix} X_{11} & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & X_{kk} & 0 \\ W_1 & \dots & W_k & W_{k+1} \end{pmatrix}$$

where each  $X_{ii}$  is a  $t_i \times n_i$  stochastic matrix.

Now consider the block

$$U = (U_1, \dots, U_k, U_{k+1}).$$

Now the rank of  $A_1 \oplus \dots \oplus A_k$  is  $k = \text{rank}(A)$ , so each row of  $U$  is a linear combination of the first  $n - n_{k+1}$  rows of  $AP$ . Hence, since  $AP$  is stochastic,  $U_{k+1} = 0$ , and  $U_j = G_j A'_j$  for  $j = 1, \dots, k$ , where each  $A'_j$  consists of  $n_{k+1}$  rows each equal to a row of  $A_j$ , each  $G_j$  is an  $n_{k+1} \times n_{k+1}$  diagonal matrix, and  $G_1 + \dots + G_k = I$ . Thus, for  $j = 1, \dots, k$ ,

$$D_j E'_j = F_j = M_{k+1} P P^T X_j = U_j X_{jj} = G_j A'_j X_{jj} = G_j M'_j X_j = G_j E'_j.$$

But  $E'_j > 0$  and  $D_j - G_j$  is diagonal, so  $D_j = G_j$ . Hence  $U_j = D_j A'_j$  for  $j = 1, \dots, k$ . This completes the proof that (a) implies (b).

Assume now that (b) holds. Let  $Y$  be the  $n \times n$  stochastic matrix

$$Y = Y_1 \oplus \dots \oplus Y_k \oplus Z$$

where each  $Y_i$  is the  $t_i \times n_i$  stochastic matrix having each row equal to a row of  $E_i$ , and  $Z$  is any stochastic matrix of appropriate size. Then  $APY = E$ . But  $EAP = AP$ , so  $AP \mathcal{R} E$ . Now  $AP \mathcal{R} A$  and  $\mathcal{R}$  is transitive, so  $A \mathcal{R} E$ . Thus (a) holds.

That (b) and (c) are equivalent is clear, which completes the proof of the theorem.

The first of the following corollaries removes the restriction that the idempotent  $E$  be canonical, while the second gives necessary and sufficient conditions for two canonical idempotents to be  $\mathcal{R}$ -equivalent. For the proof, note that  $A \mathcal{R} B$  if and only if  $Q A Q^T \mathcal{R} Q B Q^T$  for  $Q$  any permutation matrix.

**Corollary 2.2.** *Let  $F$  be an idempotent in  $S_n$  and let  $A \in S_n$ . Then  $A \mathcal{R} F$  if and only if for some  $n \times n$  permutation matrices  $P$  and  $Q$ ,  $Q A Q^T P$  has the form (3) and  $Q F Q^T$  is a canonical idempotent.*

**Corollary 2.3.** *Two canonical idempotents in  $S_n$  are  $\mathcal{R}$ -equivalent if and only if their corresponding blocks have the same dimensions and the diagonal matrices which determine the last  $n_{k+1}$  rows of each are identical.*

Since there are uncountably many choices for the row in  $E_i$  whenever  $n_i \geq 2$ , Corollary 2.3 shows that any  $\mathcal{R}$ -class of  $S_n$  which contains one canonical idempotent  $E$  must contain uncountably many, provided only that at least one block  $E_i$  of  $E$  has order 2 or greater.

As a result of the next theorem, if a stochastic matrix  $A$  itself has the form (3), then this is sufficient for  $A$  to be regular. The matrix

$$A = \begin{pmatrix} \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \end{pmatrix}$$

is regular, since  $A$  is idempotent, but not of the form (3), so the condition is not necessary. Indeed,  $A$  cannot be put in the form (3) by any column permutation.

**Theorem 2.4.** *Let  $A \in S_n$ . If  $A$  has the form (3), then there exists a canonical idempotent  $E$  in  $S_n$  such that  $A \mathcal{R} E$ . Consequently,  $A$  is regular.*

*Proof.* Use the matrix  $Y$  introduced at the end of the proof of Theorem 2.1 with each  $Y_i$  having each entry equal to  $1/n_i$ .

The following corollary gives necessary and sufficient conditions for two regular stochastic matrices to lie in the same  $\mathcal{R}$ -class, and hence for the system (1) to be consistent. Regularity does not need to be assumed for the sufficiency.

**Corollary 2.3.** *Let  $A, B \in S_n$ . Then  $A$  and  $B$  are regular and  $A \mathcal{R} B$  if and only if there exist  $n \times n$  permutation matrices  $P_1, P_2$ , and  $Q$  such that  $QAQ^T P_1$  and  $QBQ^T P_2$  each have the form (3) with corresponding blocks having the same dimensions and with identical diagonal matrices determining the last  $n_{k+1}$  rows of each.*

*Proof.* Assume  $A$  and  $B$  are regular, and  $A \mathcal{R} B$ . Then there exists an  $n \times n$  permutation matrix  $Q$  such that  $QAQ^T$  and  $QBQ^T$  are both in the same  $\mathcal{R}$ -class as some canonical idempotent  $E$  of  $S_n$ . Thus there exist  $n \times n$  permutation matrices  $P_1$  and  $P_2$  such that  $QAQ^T P_1$  and  $QBQ^T P_2$  each have the form (3). The converse follows by using the canonical idempotent  $E$  of Theorem 2.4.

**3. Green's relation  $\mathcal{L}$  for  $S_n$ .** We again begin by characterizing those elements  $A$  in  $S_n$  which lie in the same  $\mathcal{L}$ -class with a canonical idempotent  $E$ . The result is similar to that for  $\mathcal{R}$ , but here the form of  $A$  depends upon the matrices  $E_i$  as well as the way in which  $E$  is partitioned into blocks of columns.

**Theorem 3.1.** *Let  $E$  be a canonical idempotent in  $S_n$  with rank  $k$ , and let  $A \in S_n$ . Then the following statements are equivalent.*

- (a)  $A \mathcal{L} E$ ; that is, the system  $XA = E, YE = A, X, Y \in S_n$  is consistent.

(b) There exists an  $n \times n$  permutation matrix  $P$  such that  $PA$  has the form

$$(4) \quad PA = \begin{pmatrix} A_1 & 0 & \dots & 0 & 0 \\ 0 & A_2 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & A_k & 0 \\ U_1 & U_2 & \dots & U_k & 0 \end{pmatrix}$$

where each  $A_i$  is a  $t_i \times n_i$  stochastic matrix of rank 1 having each row equal to a row of  $E_i$ ; and where  $U_i = G_i A'_i$ , where each  $A'_i$  consists of  $t = n - t_1 - \dots - t_k$  rows of  $A_i$ , each  $G_i$  is a  $t \times t$  diagonal matrix, and  $G_1 + \dots + G_k = I$ .

Proof. Assume (a) holds. Partition  $A$  by

$$A = (M_1, \dots, M_k, M_{k+1})$$

where each  $M_i$  is  $n \times n_i$ . Since  $AE = A$ ,  $M_{k+1} = 0$ , and  $M_i E_i = M_i$  for  $i = 1, \dots, k$ . Partition  $X$  by

$$X = \begin{pmatrix} X_1 \\ \dots \\ X_k \\ X_{k+1} \end{pmatrix}$$

where each  $X_i$  is  $n_i \times n$ . Then  $XA = E$  implies

$$\begin{aligned} X_i M_i &= E_i \quad \text{for } i = 1, \dots, k; \\ X_i M_j &= 0 \quad \text{for } i \neq j, \quad i, j = 1, \dots, k; \end{aligned}$$

and

$$X_{k+1} M_j = F_j \quad \text{for } j = 1, \dots, k.$$

Now  $M_i E_i = M_i$  and  $E_i$  positive imply that if  $M_i$  has a zero in any row, then that entire row must be a row of zeros. Thus a row of  $M_i$  must be either positive or a row of zeros. Now  $M_i \neq 0$ , so  $\text{rank}(M_i) = 1$  for  $i = 1, \dots, k$ .

If  $k = 1$ , then  $M_1 > 0$  since  $A = (M_1, 0)$  is stochastic. In this case  $M_1 E_1 = M_1$  implies each row of  $M_1$  is a row of  $E_1$ . Thus  $A$  itself has the form (4) where  $A_1 = M_1$  and the block  $U_1$  is missing.

Now assume  $k > 1$ . If  $M_i > 0$  for some  $i = 1, \dots, k$ , then, for any  $j \neq i$ ,  $X_j M_i = 0$  implies  $X_j = 0$ . Thus  $E_j = X_j M_j = 0$ , a contradiction. Hence each  $M_i$  has zero entries, thus rows of zeros. Let  $u_i$  be the number of positive rows in each  $M_i$ .

Let  $P_1$  be an  $n \times n$  permutation matrix such that the first  $u_1$  rows of  $P_1 M_1$  are nonzero. Then, for  $i = 2, \dots, k$ ,  $0 = X_i M_1 = X_i P_1^T P_1 M_1$ , so the first  $u_1$  columns of  $X_i P_1^T$  are zero columns. Thus

$$P_1 A = \begin{pmatrix} A_{11}^* & A_{12} & 0 \\ 0 & A_{22}^* & 0 \end{pmatrix}$$

where  $A_1^*$  is  $u_1 \times n_1$ ,  $A_{22}^*$  is  $u_1 \times (n - n_1 - n_{k+1})$ ; and

$$XP_1^T = \begin{pmatrix} X_{11} & X_{12} \\ 0 & X_{22} \\ X_{31} & X_{32} \end{pmatrix}$$

where  $X_{11}$  is  $n_1 \times u_1$ , and  $X_{31}$  is  $n_{k+1} \times u_1$ . For  $j = 2, \dots, k$ , let

$$P_1 M_j = \begin{pmatrix} B_j \\ C_j \end{pmatrix}$$

where  $B_j$  is  $u_1 \times n_j$ . Then

$$0 = X_1 M_j = X_{11} B_j + X_{12} C_j,$$

so  $X_{11} B_j = 0$ . If  $B_j$  has a positive row, then the corresponding column of  $X_{11}$  is a column of zeros. If each of the  $u_1$  rows of  $A_{12} = (B_2, \dots, B_k)$  has positive entries, then each column of  $X_{11}$  is a column of zeros. Thus  $E_1 = X_{11} A_1^* = 0$ , a contradiction. Hence  $A_{12}$  has at least one row of zeros. Let  $t_1$  be the number of rows of zeros in  $A_{12}$ . Let  $Q_1$  be an  $n \times n$  permutation matrix such that the first  $t_1$  rows of the matrix

$$Q_1 = \begin{pmatrix} A_{12} \\ A_{22}^* \end{pmatrix}$$

are rows of zeros, and the last  $u_1 - t_1$  rows are the rows of  $A_{12}$  which contain positive entries. Then

$$Q_1 P_1 A = \begin{pmatrix} A_1 & 0 & 0 \\ 0 & A_{22} & 0 \\ U_1^* & A_{32} & 0 \end{pmatrix}$$

where  $A_1$  is a  $t_1 \times n_1$  stochastic matrix and  $U_1^*$  is  $(u_1 - t_1) \times n_1$ . If  $t_1 = u_1$ , then  $U_1^* = 0$ . Now

$$Q_1 P_1 M_1 = \begin{pmatrix} A_1 \\ 0 \\ U_1^* \end{pmatrix},$$

so, since  $M_1 E_1 = M_1$ ,  $A_1 E_1 = A_1$ . Thus each row of  $A_1$  is a row of  $E_1$ . Furthermore,  $\text{rank}(Q_1 P_1 M_1) = \text{rank}(M_1) = 1$ , so  $U_1^* = G_1^* A_1'$  where  $G_1^*$  is a  $(u_1 - t_1) \times (u_1 - t_1)$  nonnegative diagonal matrix and  $A_1'$  consists of  $(u_1 - t_1)$  rows each equal to a row of  $A_1$ .

By repeating this process we obtain  $n \times n$  permutation matrices  $P_1, \dots, P_k$ ,  $Q_1, \dots, Q_k$  such that  $PA$  has the form (4) with  $P = Q_k P_k, \dots, Q_1 P_1$ . For  $i = 2, \dots, k$ , the number  $t_i$  is the number of rows of zeros in the submatrix at the  $i$ -th step which is analogous to  $A_{12}$ . If  $t_i = u_i$  for some  $i$ , then  $U_i = 0$  and so  $G_i = 0$ . Thus (b) holds.

Conversely, let  $Y$  be the  $n \times n$  stochastic matrix

$$Y = \begin{pmatrix} Y_1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & Y_k & 0 \\ Z_1 & \dots & Z_k & 0 \end{pmatrix}$$

where each  $Y_i$  is any  $n_i \times t_i$  stochastic matrix of rank 1; and where  $Z_i = D_i Y_i'$  with each  $Y_i'$  consisting of  $n_{k+1}$  rows each equal to a row of  $Y_i$ . Then, for  $i = 1, \dots, k$ ,  $Y_i E_i = A_i$  and

$$Z_i A_i = D_i Y_i' A_i = D_i A_i' = D_i E_i' = F_i$$

so  $YPA = E$ . But also  $PAE = PA$ , so  $PA \mathcal{L} E$ . Now  $A \mathcal{L} PA$  and the relation  $\mathcal{L}$  is transitive, so  $A \mathcal{L} E$ . This completes the proof.

The remaining results in this section are analogous to those at the end of the previous section. The proofs are similar to these previous results, and hence are omitted.

**Corollary 3.2.** *Let  $F$  be an idempotent in  $S_n$ , and let  $A \in S_n$ . Then  $A \mathcal{L} F$  if and only if there exist  $n \times n$  permutation matrices  $P$  and  $Q$  such that  $PQAQ^T$  has the form (4) and  $QFQ^T$  is a canonical idempotent.*

**Corollary 3.3.** *Two canonical idempotents in  $S_n$  are  $\mathcal{L}$ -equivalent if and only if their corresponding blocks  $E_i$  are identical.*

Since there are uncountably many choices for the diagonal matrices  $D_i$  in the canonical form whenever the blocks  $F_i$  appear and  $k \geq 2$ , Corollary 3.3 shows that an  $\mathcal{L}$ -class of  $S_n$  which contains one canonical idempotent must contain uncountably many canonical idempotents or else no other canonical idempotent except  $E$ .

The next theorem provides a sufficient condition for an element  $A$  of  $S_n$  to be regular. Again the matrix  $A$  given just before Theorem 2.4 is regular, but not of the form (4), so the condition is not necessary. In fact, no row permutation will bring  $A$  into the form (4).

**Theorem 3.4.** *Let  $A \in S_n$ . If  $A$  has the form (4), then there exists a canonical idempotent  $E$  in  $S_n$  such that  $A \mathcal{L} E$ . Consequently  $A$  is regular.*

As a corollary we obtain necessary and sufficient conditions for two regular elements of  $S_n$  to be in the same  $\mathcal{L}$ -class. Again, regularity is not required for the sufficiency.

**Corollary 3.5.** *Let  $A, B \in S_n$ . Then  $A$  and  $B$  are regular and  $A \mathcal{L} B$  if and only if there exist  $n \times n$  permutation matrices  $P_1, P_2$ , and  $Q$  such that  $P_1QAQ^T$  and  $P_2QBQ^T$  each have the form (4), where the corresponding blocks  $A_i$  on the main diagonals are identical.*

**4. Maximal subgroups of  $S_n$ .** We now apply the results of the last two sections to obtain a new proof of the theorem of Schwartz [10] that the maximal subgroups of  $S_n$  are isomorphic to full symmetric groups. Since the maximal subgroups containing the idempotents  $E$  and  $PEP^T$ , for any permutation matrix  $P$ , are isomorphic, it is sufficient to consider only those maximal subgroups  $H_E$  where  $E$  is a canonical idempotent.

**Theorem 4.1.** *Let  $E$  be a canonical idempotent in  $S_n$  with rank  $k$ , and let  $A \in S_n$ . Then  $A \mathcal{H} E$  if and only if  $A$  has the form*

$$(5) \quad A = \begin{pmatrix} M_{11} & \cdots & M_{1k} & 0 \\ \cdots & \cdots & \cdots & \cdots \\ M_{k1} & \cdots & M_{kk} & 0 \\ W_1 & \cdots & W_k & 0 \end{pmatrix}$$

where each  $M_{ij}$  is  $n_i \times n_j$ ; for each  $i$  and each  $j$  there is exactly one  $M_{ij}$  which is nonzero, and each row of this  $M_{ij}$  is a row of  $E_j$ ; and where  $W_j = D_i M'_{ij}$ , where  $M'_{ij}$  consists of  $n_{k+1}$  rows of the nonzero block  $M_{ij}$ .

*Proof.* Assume  $A \mathcal{H} E$ . Then  $A \mathcal{L} E$  and  $A \mathcal{R} E$ , so there exist  $n \times n$  permutation matrices  $P$  and  $Q$  such that  $AP$  has the form (3) and  $QA$  has the form (4). Partition  $A$  according to (5) where, for the moment, each  $M_{ij}$  is  $n_i \times n_j$ , and observe that the last  $n_{k+1}$  columns of  $A$  must be zero since  $AE = A$ . In the form (3), the block  $A_1$  is positive, so

$$(M_{11}, \dots, M_{1k}, 0) = (A_1, 0, \dots, 0) P^T$$

has nonzero columns. Let  $M_{1j}$  be a block having a nonzero column. Then

$$QM_j = Q \begin{pmatrix} M_{1j} \\ \cdots \\ M_{kj} \\ W_j \end{pmatrix}$$

is the  $j$ -th column of blocks in the form (4). Since  $QM_j E_j = QM_j$ ,  $M_{1j} E_j = M_{1j}$ , so each row of  $M_{1j}$  is a row of  $E_j$ . Thus  $M_{1i} = 0$  for all  $i \neq j$ .

Similarly, for each  $i = 1, \dots, k$ , exactly one block  $M_{ij}$  of  $(M_{i1}, \dots, M_{ik}, 0)$  is positive, and all the rest are zero blocks. Further, each row of this  $M_{ij}$  is a row of  $E_j$ . If, for some  $t \neq i$ ,  $M_{tj}$  is also positive, then  $(M_{i1}, \dots, M_{ik}, 0) P$  and  $(M_{t1}, \dots, M_{tk}, 0) P$  both have nonzero blocks in the  $j$ -th position of the block form (3), which is impossible. Hence, for each  $i$  and each  $j$ , exactly one  $M_{ij}$  is nonzero.

Moreover, since  $AP$  has the form (3), if  $M_{ij}$  is positive, then  $M_{ij}$  is the block  $A_i$  in  $AP$ , so

$$W_j = U_i = D_i A'_i = D_i M'_{ij}$$

where  $M'_{ij}$  consists of  $n_{k+1}$  rows of  $M_{ij}$ .

Conversely suppose  $A$  has the form (5). Let  $P = (P_{ij})$ ,  $i, j = 1, \dots, k + 1$ , where the size of the block  $P_{ij}$  is determined as follows:  $P_{ij}$  is  $n_i \times n_i$  whenever  $M_{ji} \neq 0$ , for  $i = 1, \dots, k$ , and  $P_{k+1,j}$  is  $n_{k+1} \times n_j$  for  $j = 1, \dots, k + 1$ . Let  $P_{ij} = I$  if  $P_{ij}$  is square. Otherwise set  $P_{ij} = 0$ . Then  $P$  is an  $n \times n$  permutation matrix such that  $AP$  has the form (3). Thus  $A \mathcal{R} E$ . Similarly there exists an  $n \times n$  permutation matrix  $Q$  such that  $QA$  has the form (4), so  $A \mathcal{L} E$ . Thus  $A \mathcal{H} E$ , which completes the proof.

For each  $i = 1, \dots, k$ , the mapping  $i \rightarrow j$  if  $M_{ij} \neq 0$  is a permutation of the set  $\{1, \dots, k\}$ . Such a mapping may be defined for each  $A \in H_E$ , and there are precisely  $k!$  such mappings. Thus we obtain the following corollary.

**Corollary 4.2.** (Schwarz) *Let  $E$  be an idempotent in  $S_n$  with rank  $k$ . Then the maximal subgroup  $H_E$  of  $S_n$  having  $E$  as identity element is isomorphic to the full symmetric group on  $k$  letters.*

**5. Green's relations for regular elements of  $D_n$ .** In this section Theorems 2.1 and 3.1 are used to characterize Green's relations for regular elements of  $D_n$ . These results have been obtained in another way by MONTAGUE and PLEMMONS [7] who were able to remove the restriction of regularity. The section concludes with several necessary and sufficient conditions for an element of  $D_n$  to be regular.

The first two lemmas, which are dual, characterize  $\mathcal{R}$ -classes and  $\mathcal{L}$ -classes of  $D_n$  containing canonical idempotents.

**Lemma 5.1.** *Let  $E$  be a canonical idempotent in  $D_n$  with rank  $k$ , and let  $A \in D_n$ . Then  $A \mathcal{R} E$  if and only if there exists an  $n \times n$  permutation matrix  $P$  such that  $AP = E$ .*

*Proof.* Assume  $A \mathcal{R} E$ . Since  $E$  is a canonical idempotent in  $D_n$ ,  $E$  is also a canonical idempotent in  $S_n$ . Thus, by Theorem 2.1, there exists an  $n \times n$  permutation matrix  $P$  such that

$$AP = A_1 \oplus \dots \oplus A_k$$

where each  $A_i$  is an  $n_i \times t_i$  stochastic matrix of rank 1. But each  $A_i$  is doubly stochastic, so  $n_i = t_i$  and  $A_i = E_i$ . Thus  $AP = E$ . The converse is obvious.

**Lemma 5.2.** *Let  $E$  be a canonical idempotent in  $D_n$  with rank  $k$ , and let  $A \in D_n$ . Then  $A \mathcal{L} E$  if and only if there exists an  $n \times n$  permutation matrix  $Q$  such that  $QA = E$ .*

As a result of the next three theorems, we obtain a characterization of the relations  $\mathcal{R}$  and  $\mathcal{L}$  for regular elements of  $D_n$ . The characterizations of the relations  $\mathcal{H}$  and  $\mathcal{D}$  follow as corollaries.

**Theorem 5.3.** *Let  $A$  be a regular element of  $D_n$ . Then there exists a unique idempotent  $E$  in  $D_n$  such that  $AP = E$  for some  $n \times n$  permutation matrix  $P$ .*

Proof. Since  $A$  is regular,  $A \mathcal{R} E$  for some idempotent  $E$  in  $D_n$ . If also  $A \mathcal{R} F$  and  $F^2 = F$ , then  $EF = F$  and  $FE = E$  since an idempotent is a left identity for its  $\mathcal{R}$ -class. But then  $E = E^T = (FE)^T = E^T F^T = EF = F$ , so  $E$  is unique. Now  $G = QEQ^T$  is a canonical idempotent for some  $n \times n$  permutation matrix  $Q$ . Also  $QAQ^T \mathcal{R} G$ , so, by Lemma 5.1, there exists an  $n \times n$  permutation matrix  $P_1$  such that  $QAQ^T P_1 = G = QEQ^T$ . Thus  $AQ^T P_1 Q = E$ , so  $AP = E$  where  $P = Q^T P_1 Q$ .

**Theorem 5.4.** *Let  $A$  be a regular element of  $D_n$ . Then there exists a unique idempotent  $F$  in  $D_n$  such that  $QA = F$  for some  $n \times n$  permutation matrix  $Q$ .*

**Theorem 5.5.** *Let  $A, B$  be regular elements of  $D_n$ . Then  $A \mathcal{R} B[A \mathcal{L} B]$  if and only if  $B = AP[B = QA]$  for some  $n \times n$  permutation matrix  $P[Q]$ .*

Proof. Assume  $A \mathcal{R} B$ . By Theorem 5.3 there exists a unique idempotent  $E$  in  $D_n$  such that  $AP_1 = E$  for some  $n \times n$  permutation matrix  $P_1$ . Now  $B \mathcal{R} E$ , so, by Lemma 5.1, there exists an  $n \times n$  permutation matrix  $P_2$  such that  $BP_2 = E$ . Thus  $B = EP_2^T = AP_1 P_2^T = AP$ , where  $P = P_1 P_2^T$ . The converse is obvious.

**Corollary 5.6.** *Let  $A, B$  be regular elements of  $D_n$ . Then  $A \mathcal{H} B$  if and only if  $B = QA = AP$  for some  $n \times n$  permutation matrices  $P$  and  $Q$ .*

Proof. The corollary is an immediate consequence of Theorem 5.5 and the definition of  $\mathcal{H}$ .

**Corollary 5.7.** *Let  $A, B$  be regular elements of  $D_n$ . Then  $A \mathcal{D} B$  if and only if  $B = QAP$  for some  $n \times n$  permutation matrices  $P$  and  $Q$ .*

Proof. Assume  $A \mathcal{D} B$ . Then there exists  $X$  in  $D_n$  such that  $A \mathcal{R} X$  and  $X \mathcal{L} B$  [2, Lemma 2.1]. Now  $X = AP = Q^T B$  for some  $n \times n$  permutation matrices  $P$  and  $Q$ , so  $B = QAP$ . The converse is obtained by reversing steps.

We conclude this section with several necessary and sufficient conditions for an element of  $D_n$  to be regular. A number of other conditions may be found in [7].

**Theorem 5.8.** *Let  $A \in D_n$ . Then the following statements are equivalent.*

- (1)  $A$  is regular.
- (2)  $QAP$  is idempotent for some  $n \times n$  permutation matrices  $P$  and  $Q$ .
- (3)  $Q_1 A$  is idempotent for some  $n \times n$  permutation matrix  $Q_1$ .
- (4)  $AP_1$  is idempotent for some  $n \times n$  permutation matrix  $P_1$ .
- (5)  $AA^T \mathcal{R} A$ ; that is,  $AA^T = AP$  for some  $n \times n$  permutation matrix  $P$ .
- (6)  $A^T A \mathcal{L} A$ ; that is,  $A^T A = QA$  for some  $n \times n$  permutation matrix  $Q$ .

**Proof.** The equivalence of the first four statements follows from Theorems 5.3 and 5.4 and Corollary 5.7. If (4) holds, then  $A = EP_1^T$ , where  $E^2 = E$ , so  $AA^T = EP_1^T P_1 E = E = AP_1$ , so  $AA^T \mathcal{R} A$ , proving (5). We now show (5) implies (1), which will complete the proof since (6) is dual to (5). Assume  $AA^T = AP$ . Now  $AA^T = (AA^T)^T = (AP)^T = P^T A^T$ , so  $A = AA^T P^T = P^T A^T P^T$ . Thus

$$AA^T A = APA = APP^T A^T P^T = AA^T P^T = A,$$

so  $A$  is regular.

**6. Maximal subgroups of  $D_n$ .** In this section we give a new proof of the characterization of the maximal subgroups of  $D_n$  which was first found independently by Schwarz [11] and FARAHAT [4]. Every regular  $\mathcal{D}$ -class of  $D_n$  contains a canonical idempotent, and every idempotent element of  $D_n$  is permutationally similar to a canonical idempotent. Thus it suffices to consider only those maximal subgroups  $H_E$  of  $D_n$  having a canonical idempotent  $E$  as identity element, since all other maximal subgroups will be isomorphic to one of these.

**Theorem 6.1.** (Schwarz; Farahat) *Let  $E$  be the canonical idempotent in  $D_n$  corresponding to  $n = n_1 + \dots + n_k$ , with  $n_1 \geq n_2 \geq \dots \geq n_k \geq 1$ . Let  $m_1, \dots, m_t$  be the distinct members of the sequence  $n_1, \dots, n_k$ , with  $m_1 = n_1, m_t = n_k$ , and  $m_1 > m_2 > \dots > m_t$ . For  $i = 1, \dots, t$ , let  $p_i$  be the multiplicity of  $m_i$  in the sequence  $n_1, \dots, n_k$ . Then the maximal subgroup  $H_E$  of  $D_n$  is isomorphic to the group  $S(p_1) \oplus \oplus S(p_2) \oplus \dots \oplus S(p_t)$  where, for  $i = 1, \dots, t$ ,  $S(p_i)$  is the full symmetric group on  $p_i$  letters.*

**Proof.** Let  $A \in D_n$ . Then, by Corollary 5.6,  $A \in H_E$  if and only if there exist  $n \times n$  permutation matrices  $P$  and  $Q$  such that  $A = QE = EP$ ; that is,  $E = Q^T EP$ . Thus  $A \in H_E$  if and only if  $A$  can be obtained from  $E$  by a permutation of rows (or columns) which permutes the  $p_i$  blocks of  $E$  which have each entry equal to  $1/m_i$ ,  $i = 1, \dots, t$ , and which does not interchange blocks with entries  $1/m_i$  and  $1/m_j$  whenever  $i \neq j$ . For each  $i = 1, \dots, t$ , the set of all such permutations of blocks of  $E$  is isomorphic to the group  $S(p_i)$ . Thus  $H_E$  is isomorphic to the group  $S(p_1) \oplus \dots \oplus S(p_t)$ .

**7. Generalized matrix inverses in  $D_n$ .** Given any matrix  $A$  over the complex field, PENROSE [8] shows that there exists a unique complex matrix  $X$  such that  $AXA = A$ ,  $XAX = X$ , and each of  $AX$  and  $XA$  is Hermitian. The matrix  $X$  is called the Moore-Penrose inverse of  $A$ , and is denoted by  $A^+$ . It has applications in approximation theory and in numerical analysis; see, e.g., [1], [5]. A semi-inverse of  $A$  is a matrix  $X$  satisfying the equations  $AXA = A$  and  $XAX = X$ .

In this section we give a new proof of the result of PLEMMONS and CLINE [9] concerning the semi-inverse and the Moore-Penrose inverse of a regular doubly stochastic matrix. The stochastic case is considered in [12].

**Theorem 7.1.** (PLEMMONS and CLINE) *A regular element  $A$  of  $D_n$  has  $A^T$  as its unique semi-inverse in  $D_n$ . Moreover,  $A^T$  is the Moore-Penrose inverse of  $A$ .*

*Proof.* Let  $A$  be a regular element of  $D_n$ . By Theorem 5.3 there exists a unique idempotent  $E$  in  $D_n$  such that  $A \mathcal{R} E$ , so  $A = EP$  for some  $n \times n$  permutation matrix  $P$ . Now  $A^T = P^T E$ , so

$$AA^T A = EPP^T EEP = EP = A$$

and

$$A^T AA^T = P^T EEP P^T E = P^T E = A^T,$$

so  $A^T$  is a semi-inverse of  $A$  in  $D_n$ . Since  $E$  is the only idempotent such that  $A \mathcal{R} E$  it follows [2, Theorem 2.18] that  $A^T$  is the unique semi-inverse of  $A$  in  $D_n$ . Since doubly stochastic idempotents are symmetric,  $A^T$  must be the Moore-Penrose inverse of  $A$ .

#### References

- [1] *A. Ben-Israel* and *A. Charnes*, "Contributions to the theory of generalized inverses", *J. Soc. Indust. Appl. Math.*, *XI* (1963), 667–699.
- [2] *A. H. Clifford* and *G. B. Preston*, *The Algebraic Theory of Semigroups*, Vol. *I*, Math. Surveys No. 7, Amer. Math. Soc., Providence, 1961.
- [3] *J. L. Doob*, "Topics in the theory of Markov chains", *Trans. Amer. Math. Soc.*, *52* (1942), 37–64.
- [4] *H. K. Farahat*, "The semigroup of doubly-stochastic matrices", *Proc. Glasgow Math. Assoc.*, *7* (1966), 178–183.
- [5] *T. N. E. Greville*, "Some applications of the pseudoinverse of a matrix", *SIAM Rev.*, *II* (1960), 15–22.
- [6] *K. H. Hofmann* and *P. S. Mostert*, *Elements of Compact Semigroups*, Charles E. Merrill, Columbus, 1966.
- [7] *J. S. Montague* and *R. J. Plemmons*, "Convex matrix equations", *Bull. Amer. Math. Soc.*, *78* (1972), 965–968.
- [8] *R. Penrose*, "A generalized inverse for matrices", *Proc. Cambridge Phil. Soc.*, *51* (1955), 406–413.
- [9] *R. J. Plemmons* and *R. E. Cline*, "The generalized inverse of a nonnegative matrix", *Proc. Amer. Math. Soc.*, *31* (1972), 46–50.
- [10] *Š. Schwarz*, "On the structure of the semigroup of stochastic matrices", *Magyar Tud. Akad. Kutato Int. Kozl.*, (A) *9* (1964), 297–311.
- [11] *Š. Schwarz*, "A note on the structure of the semigroup of doubly stochastic matrices", *Mat. Časopis Sloven. Akad. Vied.*, *17* (1967), 308–316.
- [12] *J. R. Wall*, "Generalized inverses of stochastic matrices", *Lin. Alg. Appl.*, to appear.

*Author's address:* Mathematics Department, Auburn University, Auburn, Alabama 36830, U.S.A.