

F. Burton Jones

Connected simple graphs and a selection problem

Czechoslovak Mathematical Journal, Vol. 25 (1975), No. 2, 300–301

Persistent URL: <http://dml.cz/dmlcz/101321>

Terms of use:

© Institute of Mathematics AS CR, 1975

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

CONNECTED SIMPLE GRAPHS AND A SELECTION PROBLEM

F. B. JONES, Riverside

(Received April 8, 1974)

A problem left open by CEDER and BRUCKNER in [1] is the following (as modified by PAUL ERDŐS):

Suppose that M is a connected subset of the plane such that each vertical line intersects M in exactly two points. Does M contain a connected subset M_1 such that every vertical line intersects M_1 in exactly one point? In other words, must there exist a function (from R to R) whose graph lies in M and is connected? The example given below shows that such a selection is not always possible.

Let M be the union of the following plane sets:


{the graph of $y = 0$ for $x \leq 1$ };

{the graph of $y = 1$ for $x \geq 1$ };

{the graph of $y = x$ for $x \leq 0$ and $x \geq 1$ };

and

the closed straight line intervals from $(1 - 1/2^n, 1)$
to $(1 - 1/2^{n+1}, 0)$ for $n = 0, 1, 2, 3, \dots$



Clearly, M is connected. Also it is clear that if one is to make a selection of one point for each vertical line which is to produce a connected set, one must choose only those points of M on the x -axis between $x = 0$ and $x = 1$. This is true because, roughly speaking, these points separate M between $-\infty$ and $+\infty$. But having done this, there is no way to connect to the points in M for $x \geq 1$ and $y \geq 1$.

References

- [1] *A. M. Bruckner and J. Ceder, On jumping functions by connected sets, Czechoslovak Math. J., 22 (1972), 435–448: in particular, Question 1, p. 442.*

Author's address: Department of Mathematics, University of California, Riverside, CA. 92502, U.S.A.