

Jorge Martinez

Torsion theory for lattice-ordered groups Part II: Homogeneous  $l$ -groups

*Czechoslovak Mathematical Journal*, Vol. 26 (1976), No. 1, 93–100

Persistent URL: <http://dml.cz/dmlcz/101376>

## Terms of use:

© Institute of Mathematics AS CR, 1976

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

TORSION THEORY FOR LATTICE-ORDERED GROUPS  
PART II: HOMOGENEOUS  $l$ -GROUPS

JORGE MARTINEZ, Gainesville

(Received April 8, 1974)

**Introduction.** This note pursues in greater detail some of the discussion initiated in [6]; we use the same terminology and notation of [6]. We shall assume that the reader is familiar with CONRAD [3], and FUCHS [4], and we shall systematically treat such material as standard theory of lattice-ordered groups (henceforth:  $l$ -groups). At the risk of sounding pretentious, we will also assume the reader has the good sense to read [6] before taking this on, although it is by no means a prerequisite.

We write all  $l$ -groups additively without regard to commutativity or the lack of it. If  $A$  and  $B$  are subsets of a set  $X$ , we denote (proper) containment by  $(A \subset B) A \subseteq B$ ;  $A \setminus B$  stands for the complement of  $B$  in  $A$ .

Let us start by reviewing the notion of a torsion class from [6]. A class  $\mathcal{T}$  of  $l$ -groups will be called a *torsion class* if it is closed with respect to taking 1) convex  $l$ -subgroups, 2)  $l$ -homomorphic images and 3) joins of convex  $l$ -subgroups in  $\mathcal{T}$ . With each torsion class we associate a radical (also denoted by  $\mathcal{T}$ ), so that if  $G$  is an  $l$ -group then  $\mathcal{T}(G)$  is the join of all the convex  $l$ -subgroups of  $G$  belonging to  $\mathcal{T}$ . If  $\mathcal{T}$  is a torsion class then: a)  $\mathcal{T}(A) = A \cap \mathcal{T}(G)$  for each convex  $l$ -subgroup  $A$  of  $G$ , b)  $[\mathcal{T}(G)] \Phi \subseteq \mathcal{T}(H)$  for each  $l$ -epimorphism  $\Phi : G \rightarrow H$  (proposition 1.1 in [6]). Conversely, any radical satisfying a) and b) gives rise to a torsion class of which it is the associated torsion radical (proposition 1.2 in [6]). Finally, we say that a torsion class is *complete* if it closed under extensions.

In § 4 of [6] we introduced the notion of a homogeneous  $l$ -group:  $G$  is *homogeneous* if for each torsion class  $\mathcal{T}$  either  $G \in \mathcal{T}$  or else  $\mathcal{T}(G) = 0$ . Since torsion radicals are characteristic  $l$ -ideals it follows that all characteristically simple  $l$ -groups are homogeneous. In this category fall the free abelian  $l$ -groups, cardinal sums of reals, rationals or integers, periodic real sequences, etc.

We also developed two criteria for telling when an  $l$ -group  $G$  is homogeneous. Let  $\mathcal{X}_G$  and  $\mathcal{X}^G$  denote respectively the torsion class generated by  $G$ , and the largest torsion class relative to having  $\mathcal{T}(G) = 0$ . The two criteria are as follows:

**Theorem** (4.1 in [6]): *If  $G$  is homogeneous then  $\mathcal{X}^G$  is complete and meet irreducible in the lattice of all torsion classes. Conversely, if  $\mathcal{X}^G$  is meet irreducible then  $G$  has a nontrivial homogeneous  $l$ -ideal. On the other hand, if  $\mathcal{X}$  is any complete, meet irreducible torsion class, there is a homogeneous  $l$ -group  $H$  such that  $\mathcal{X} = \mathcal{X}^H$ .*

**Theorem** (4.2 in [6]): *If  $G$  is homogeneous  $\mathcal{X}_G$  is join irreducible in the lattice of all torsion classes. Conversely, if  $\mathcal{U}$  is a join irreducible torsion class and it covers  $\mathcal{U}^\sim$ , and there is an  $l$ -group  $H \in \mathcal{U}$  so that  $\mathcal{U}^\sim(H) = 0$ , then  $H$  is homogeneous,  $\mathcal{X}_H = \mathcal{U}$  and  $\mathcal{X}^H$  is the largest torsion class satisfying  $\mathcal{T} \cap \mathcal{U} = \mathcal{U}^\sim$ .*

Note. We use meet and join irreducibility relative to arbitrary meets and joins respectively.

We conclude this introduction with some additional basic facts about homogeneous  $l$ -groups:

1. If  $G$  is homogeneous so is every convex  $l$ -subgroup.
2. If  $G = \boxplus \{G_\gamma \mid \gamma \in \Gamma\}$ , a cardinal sum of  $l$ -groups  $G_\gamma$ , and  $G_\gamma \simeq G_\delta$  for  $\gamma, \delta \in \Gamma$  and each  $G_\gamma$  is homogeneous, then  $G$  is homogeneous.
3. If  $G$  is homogeneous and  $A$  is a non-trivial convex  $l$ -subgroup of  $G$  then  $\mathcal{X}_G = \mathcal{X}_A$ .

The above are easy to prove and the reader is invited to try them.

## 1. FINITE VALUED, HOMOGENEOUS $l$ -GROUPS

Before launching ahead we need a general lemma concerning the principal torsion classes  $\mathcal{X}_G$ .

**1.1 Lemma.** *Let  $G$  and  $H$  be two  $l$ -groups;  $H \in \mathcal{X}_G$  if and only if  $H = \bigvee_{i \in I} H_i$ , where each  $H_i$  is a convex  $l$ -subgroup of  $H$ , and for each  $i \in I$  there exist convex  $l$ -subgroups  $N_i \subseteq D_i$  of  $G$  with  $N_i$  normal in  $D_i$ , such that  $D_i/N_i \simeq H_i$ .*

*Proof.* It is clear that if  $H$  is a join of convex  $l$ -subgroups as described in the lemma then  $H \in \mathcal{X}_G$ . So all that's needed is to show that the class  $\mathcal{T} = \{H \mid H = \bigvee_{i \in I} H_i, H_i \text{ a convex } l\text{-subgroup of } H \text{ and a quotient of a convex } l\text{-subgroup of } G\}$  is a torsion class. Obviously,  $\mathcal{T}$  is closed under joins of convex  $l$ -subgroups in  $\mathcal{T}$ . Next, suppose  $H \in \mathcal{T}$  and  $K$  is a convex  $l$ -subgroup of  $H$ , write  $H = \bigvee_{i \in I} H_i$ , each  $H_i$  a convex  $l$ -subgroup of  $H$  isomorphic to  $D_i/N_i$ , where  $N_i \subseteq D_i$  are convex  $l$ -subgroups of  $G$  and  $N_i$  is normal in  $D_i$ . Now  $K = \bigvee_{i \in I} K \cap H_i$  and  $K \cap H_i \simeq D_i^*/N_i$ , where  $D_i^* \subseteq D_i$  is a convex  $l$ -subgroup of  $G$ ; thus  $K \in \mathcal{T}$ .

Finally, suppose  $\phi : H \rightarrow L$  is an  $l$ -homomorphism of  $H \in \mathcal{T}$  onto the  $l$ -group  $L$ . If  $H = \bigvee_{i \in I} H_i$  as before, then  $L = \bigvee_{i \in I} H_i \phi$  and  $H_i \phi$  is a quotient of a quotient of

a convex  $l$ -subgroup of  $G$ . Thus  $H;\phi$  is itself a quotient of a convex  $l$ -subgroup of  $G$ , and we are able to conclude that  $L \in \mathcal{T}$ . Hence  $\mathcal{T}$  is a torsion class,  $\mathcal{T} = \mathcal{X}_G$  and the lemma is proved.

It will be useful to make the following definition now: if  $G$  is an  $l$ -group,  $N \subseteq D$  are convex  $l$ -subgroups of  $G$  with  $N$  normal in  $D$ , we call  $D/N$  a *subquotient* of  $G$ . If  $T$  is a well ordered set and  $\{G_t \mid t \in T\}$  is a family of convex  $l$ -subgroups of  $G$ , so that  $G_s \subseteq G_t$  if  $s < t$ , and  $\bigcup G_t = G$ , we call  $G$  an *inductive limit* of the  $G_t$ .

**1.2 Lemma.** *Let  $H$  be a finite valued  $l$ -group;  $H \in \mathcal{X}_G$  if and only if  $H$  is a cardinal sum of inductive limits of subquotients of  $G$ .*

(Note. Here and for the remainder of the section we assume  $G$  is a finite valued  $l$ -group.)

*Proof.* Every finite valued  $l$ -group  $H$  is a cardinal sum of cardinally indecomposable  $l$ -groups. (This can be seen as follows: the root system of regular subgroups is a disjoint union of indecomposable – and hence directed – root systems. The indecomposable summands of  $H$  are then obtained by considering the  $l$ -ideals generated by the special elements “living on” a fixed root system component. It should be clear of course, that if the root system of regular subgroups of  $H$  is indecomposable to start with, then  $H$  is cardinally indecomposable.)

So we will lose no generality if we prove that whenever  $H \in \mathcal{X}_G$  and cardinally indecomposable, then  $H$  is isomorphic to an inductive limit of subquotients of  $G$ .

Assuming this, the proof from here on breaks down into two parts: a) if  $0 < x \in H$  is special then  $H(x)$  is isomorphic to a subquotient of  $G$ ; b)  $H = \bigcup \{H(x_j) \mid j \in J\}$  where  $J$  is a well ordered set, and each  $0 < x_j \in H$  is special, so that  $H(x_j) \subset H(x_k)$  if  $j < k$ .

a) If  $0 < x \in H$  is special then  $H(x)$  is a lexicographic extension of an  $l$ -ideal  $M$  of  $H(x)$  so that  $H(x)/M$  is a subgroup of  $\mathbf{R}$ , the additive real with the usual order.  $H(x) \in \mathcal{X}_G$  and join irreducible in the lattice of convex  $l$ -subgroups of  $H$ , and must therefore be a subquotient of  $G$ .

b) We select a root (root  $\equiv$  maximal chain) out of the root system of regular subgroups of  $H$ ; since  $H$  is indecomposable we can state that if we pick for each regular subgroup  $M$  on that root, a special element  $x_M > 0$  having its value at  $M$ , then  $H = \bigcup_{x_M > 0} H(x_M)$ . By choosing a suitable cofinal, well ordered subset of the root, we obtain the desired well ordered chain of principal convex  $l$ -subgroups.

The referee of this note has provided the author with an example to show that the words “inductive limit of” cannot be deleted in lemma 1.2. Consider the  $o$ -group  $A_n$  of the lexicographic product of  $n$  copies of the additive reals, ordered from left to right. Let  $G$  be the cardinal sum of the  $A_n$  ( $n = 1, 2, \dots$ ). Let  $H$  be the direct sum of copies of the additive reals  $R_n$  ( $n = 1, 2, \dots$ ), lexicographically ordered from left to right. Then  $H$  is the inductive limit of convex subgroups which are quotients of  $G$ , and hence  $H \in \mathcal{X}_G$ . But  $H$  has infinite sequences  $a_1 \gg a_2 \gg a_3 \gg \dots$  (where  $a \gg b$

means that  $a \geq kb$  for each positive integer  $k$ ), and no totally ordered subquotient of  $G$  has this property. Thus  $H$  is not a subquotient of  $G$ . (Note also that  $H$  is indecomposable, as it is an  $o$ -group.)

We are now ready for the main theorem of this section; we give it in two stages.

**1.3 Theorem.** *Suppose  $G$  is an indecomposable  $l$ -group;  $G$  is homogeneous if and only if for each  $0 < a \in G$ ,  $G$  is isomorphic to an inductive limit of subquotients of  $G(a)$ .*

*Proof.* If  $G$  is isomorphic to an inductive limit of subquotients of  $G(a)$  for each  $0 < a \in G$ , then  $G$  is clearly homogeneous.

Conversely, suppose  $0 < x \in G$  and  $G$  is homogeneous.  $G(x)$  and  $G$  generate the same torsion class  $\mathcal{X}$ . As  $G \in \mathcal{X}$ , by lemma 1.2,  $G$  is isomorphic to an inductive limit of subquotients of  $G(x)$ , and we're done.

**1.4 Proposition.**  *$G$  is homogeneous if and only if it is a cardinal sum of indecomposable, homogeneous  $l$ -groups, each one being an inductive limit of subquotients of any other in the decomposition.*

*Proof.* Sufficiency. Suppose  $G = \boxplus \{G_\gamma \mid \gamma \in \Gamma\}$  where each  $G_\gamma$  is indecomposable and homogeneous, and further  $G_\gamma$  is an inductive limit of subquotients of  $G_\delta$ , for  $\gamma, \delta \in \Gamma$ . If  $\mathcal{T}$  is a torsion class and  $\mathcal{T}(G) \neq 0$ , then  $\mathcal{T}(G_\gamma) \neq 0$  for some  $\gamma \in \Gamma$ , since  $\mathcal{T}(G) = \boxplus \mathcal{T}(G_\gamma)$ ; (proposition 1.3 in [6]). Thus  $G_\gamma \in \mathcal{T}$  and hence each  $G_\delta \in \mathcal{T}$  since  $G_\delta$  is an inductive limit of subquotients of  $G_\gamma$ . It follows that  $G \in \mathcal{T}$  and  $G$  is homogeneous.

Necessity. If  $G$  is homogeneous, write  $G = \boxplus_{\gamma \in \Gamma} G_\gamma$ , where each  $G_\gamma$  is indecomposable. Each  $G_\gamma$  is homogeneous, and  $\mathcal{X}_G = \mathcal{X}_{G_\gamma}$  for each  $\gamma \in \Gamma$ . By lemma 1.2  $G_\gamma$  is an inductive limit of subquotients of  $G_\delta$  for all  $\gamma, \delta \in \Gamma$ .

Recall that an  $l$ -group  $G$  has *property (F)* if each  $0 < g \in G$  exceeds at most finitely many pairwise disjoint elements.

**1.4.1 Corollary.** *Suppose  $G$  has property (F);  $G$  is homogeneous if and only if  $G$  is a cardinal sum of homogeneous  $o$ -groups, any two of which are inductive limits of subquotients of each other.*

Before going on to the non-finite valued case we should point out the following offshoot of lemma 1.2.

**1.5 Proposition.** *Let  $\mathcal{T}$  be a class of finite valued  $l$ -groups closed under taking convex  $l$ -subgroups and quotients. Then  $\mathcal{T}$  is a torsion class if and only if  $\mathcal{T}$  is closed under cardinal sums and unions of chains of convex  $l$ -subgroups in  $\mathcal{T}$ .*

*Proof.* Necessity is obvious, so we move right on to the sufficiency. Let  $H = \bigvee_{i \in I} H_i$ , where each  $H_i$  is a convex  $l$ -subgroup of  $H$  belonging to  $\mathcal{T}$ . In the spirit

of lemma 1.2 we may assume without loss of generality that  $H$  is indecomposable, and write  $H = \bigcup\{H(x_j) \mid j \in J\}$ , where  $J$  is well ordered, each  $x_j > 0$  is special, and  $H(x_j) \subset H(x_k)$  for  $j < k$ . Clearly, each  $H(x_j) \subseteq H_{i(j)}$ , for a suitable  $i(j) \in I$ . So each  $H(x_j) \in \mathcal{T}$  and hence  $H \in \mathcal{T}$ , which shows that  $\mathcal{T}$  is a torsion class.

## 2. NOW INFINITELY MANY VALUES

Before saying anything really intelligent about homogeneous  $l$ -groups which are not finite valued, we must present some new torsion classes; let  $\alpha$  be an infinite cardinal number, and  $V_\alpha$  be the class of all  $l$ -groups in which all non-zero elements have at most  $\alpha$  values.

**2.1 Lemma.**  $V_\alpha$  is a torsion class for each infinite cardinal number  $\alpha$ .

*Proof.* Suppose  $G \in V_\alpha$  and  $C$  is a convex  $l$ -subgroup of  $G$ ; if  $0 < c \in C$  and  $N$  is a value of  $c$  in  $C$  then there is a regular subgroup  $M$  of  $G$  so that  $M \cap C = N$ , and of course  $M$  is a value of  $c$  in  $G$ . This suffices to show that  $c$  has at most  $\alpha$  values in  $C$ ,  $V_\alpha$  is therefore closed relative to convex  $l$ -subgroups.

Next, suppose  $G \in V_\alpha$  and  $K$  is an  $l$ -ideal of  $G$ . If  $0 \neq g + K$  its values in  $G/K$  are in one to one correspondence with the values of  $g$  in  $G$  that exceed  $K$ . Hence  $g + K$  has at most  $\alpha$  values, and  $G/K \in V_\alpha$ .

Finally, suppose  $G = \bigvee_{i \in I} G_i$ , where each  $G_i$  is a convex  $l$ -subgroup of  $G$  belonging to  $V_\alpha$ .  $G$  is the subgroup generated by the  $G_i$ , and if  $0 < g \in G$  then  $g = g_{i_1} + \dots + g_{i_n}$ , where  $0 < g_{i_\lambda}$  and  $g_{i_\lambda} \in G_{i_\lambda}$  ( $\lambda = 1, 2, \dots, n$ ). If  $M$  is a value of  $g$  in  $G$  then  $M$  is a value of at least one of the  $g_{i_\lambda}$ . The values of  $g_{i_\lambda}$  in  $G$  are in one to one correspondence with its values in  $G_{i_\lambda}$ , whence  $g$  has at most  $\alpha$  values in  $G$ . (Note: it is here that we need the assumption that  $\alpha$  be an infinite cardinal number.)

This completes the proof of the lemma.

**2.2 Theorem.** Suppose  $G$  is a homogeneous, hyper-archimedean  $l$ -group. Then either  $G$  is finite valued, in which case it is a cardinal sum of copies of a fixed subgroup of  $\mathbf{R}$ , or else there is an infinite cardinal number  $\alpha$  so that each non-zero element of  $G$  has precisely  $\alpha$  values.

*Proof.* Since  $G$  is hyper-archimedean all regular subgroups are maximal, and so it follows that if  $0 < g \in G$  has at most  $\beta$  values, then any positive element below  $g$  has the same property. We ignore the finite valued case, the conclusion being obvious by now. Thus we assume there is an infinite cardinal number  $\alpha$ , and an element  $0 < a \in G$  having  $\alpha$  values. By our remark here, and because  $G$  is homogeneous  $G \in V_\alpha$ .

If some  $h \in G$  has fewer values, say  $\gamma$  of them, with  $\gamma < \alpha$ , then  $V_\gamma(G) \neq 0$ , which implies that  $G \in V_\gamma$ , a contradiction. We must conclude then that each non-zero element of  $G$  has precisely  $\alpha$  values.

For each infinite cardinal number  $\alpha$  we can construct a homogeneous, hyper-archimedean  $l$ -groups  $B_\alpha$  in which each non-zero element has precisely  $\alpha$  values. Let  $I$  be a set of cardinality  $\alpha$ , and let  $B$  be the  $l$ -group of bounded, integer valued functions on  $I$ . Let  $J$  be the  $l$ -ideal of functions whose supports have cardinality less than  $\alpha$ ; finally, set  $B_\alpha = B/J$ .

Since  $B$  is hyper-archimedean so is  $B_\alpha$ ; in fact  $B$  is an  $S$ -group; that is, it is generated by its singular elements. (Recall that an element  $0 < s$  is *singular* if  $0 \leq g \leq s$  implies that  $g \wedge (s - g) = 0$ .) The  $S$ -groups form a torsion class, so  $B_\alpha$  is also an  $S$ -group. Define  $u \in B$  to be the constant function 1;  $u$  is a strong order unit for  $B$ , so that  $u + J$  is a strong order unit for  $B_\alpha$ .

Suppose  $0 < g \in B \setminus J$  and let  $I_g$  be the support of  $g$ . Since  $I$  and  $I_g$  have the same cardinality there is a bijection between them; it extends to an isomorphism  $\phi$  from  $B$  onto  $B(g)$  so that  $u\phi = g$  and  $J$  is mapped onto the  $l$ -ideal  $J_g$  of all functions in  $B(g)$  whose supports have cardinality less than  $\alpha$ . Noting that  $J_g = J \cap B(g)$  we have,  $B_\alpha = B/J \simeq B(g)/J_g = B(g)/B(g) \cap J \simeq B(g) + J/J$ , and the last one is the  $l$ -ideal of  $B_\alpha$  generated by  $g + J$ .


We've shown then that every principal convex  $l$ -subgroup of  $B_\alpha$  is isomorphic to  $B_\alpha$ . This clearly suffices to establish that  $B_\alpha$  is homogeneous. Next, partition  $I$  into  $\alpha$  subsets of cardinality  $\alpha$  each. This gives rise to  $\alpha$  pairwise disjoint singular elements whose supports all have cardinality  $\alpha$ , by passing to the appropriate characteristic functions. This should convince the reader that  $u + J$  has  $\alpha$  values, and hence that all non-zero elements of  $B_\alpha$  have  $\alpha$  values.

We give two examples in closing this section:

1. First, theorem 2.2 only gives a necessary condition for homogeneity. For example let  $G$  be the  $l$ -group of periodic real sequences with integers in all odd components;  $G$  is not homogeneous, it is hyper-archimedean, and each non-zero element of  $G$  has countably many values.

2. An  $l$ -group may be homogeneous, not finite valued, yet contain some finite valued elements. Consider the  $v$ -group  $V = V(A, \mathbf{R})$  over the root system  $A$  pictured below.  $V$  is not finite valued; for example there is the function  $f \in V$  defined by

$$f(n) = \begin{cases} 1 & \text{if } n = 2^k \\ 0 & \text{otherwise.} \end{cases}$$



Yet  $V$  is homogeneous, because each  $0 < x \in V$  exceeds a special element  $0 < y \in V$  such that  $V(y) \simeq V$ .

### 3. VOIDS AND CLOSING COMMENTS

Obviously this note fails to answer certain basic questions about homogeneous  $l$ -groups. Unquestionably, a defining condition is needed that circumvents torsion classes; we have it only in the finite valued case.

One interesting question involves homogeneous  $o$ -groups: we know now that they are characterized by saying that each positive element “dominates” a subquotient isomorphic to the given group. The question is whether *subquotient* may be replaced by *convex subgroup*. The author has tried in vain to concoct a Hahn group whose skeletal chain  $A$  is such that for each  $\lambda \in A$  there is a convex subset  $\Gamma_\lambda$  of  $A$ , satisfying  $\mu \leq \lambda$  for each  $\mu \in \Gamma_\lambda$ , whose corresponding Hahn group is isomorphic to the original  $o$ -group, yet  $\Gamma_\lambda$  cannot always be chosen to be an order ideal of  $A$ . The author blithely believes that such examples exist.

We conclude with two examples; the first answers a conjecture of [6], the second raises some new intriguing questions.

a) Let  $G = C([0, 1])$ ; it is shown in [1], corollary 6.11 that  $K = \{f \in G \mid f \text{ vanishes off an open interval in } [0, 1]\}$ , is the smallest non-trivial characteristic  $l$ -ideal of  $G$ . We proved in [6], § 4, that  $K$  could not be a torsion radical, and wondered whether  $G$  is not actually homogeneous.

Let us see that it is homogeneous: suppose  $0 < f \in G$ ; then there exist  $a, b \in (0, 1)$ ,  $a < b$ , so that  $f(t) > 0$  for each  $t \in (a, b)$ . Pick two points  $c$  and  $d$  so that  $a < c < d < b$ , and let  $L$  be the  $l$ -ideal of functions that vanish off  $(a, b)$ . We can define an  $l$ -homomorphism  $\phi$  from  $L$  onto  $C([c, d])$  by restriction; (it is reasonably obvious that  $\phi$  is indeed onto.) What we’ve proved is that for each  $0 < f \in G$  there is a subquotient of  $G(f)$  isomorphic to  $G$ ; it follows that  $G$  is homogeneous.

b) Let  $G = \mathbf{Z} \coprod \mathbf{Z}$ , the free product as abelian  $l$ -groups of the additive integers with themselves; it can be seen from the discussion of [5] that  $G$  is the  $l$ -subgroup of  $C([0, 1])$  generated by  $f(t) = t$  and  $g(t) = 1 - t$ . This says that  $G$  is the full group of piecewise linear and continuous functions on  $[0, 1]$ , (finitely many pieces), with integral slopes everywhere. From here one can develop an argument very similar to — but more delicate than — the one in a) to show that  $G$  is homogeneous.

The free abelian  $l$ -group on any number of generators is a free product of copies of  $\mathbf{Z} \boxplus \mathbf{Z}$ , and is homogeneous. In view of this and b) above one ought to wonder whether the free (abelian) product of any number of copies of a (homogeneous) abelian  $l$ -group is homogeneous.

#### Bibliography

- [1] Richard Byrd, Paul Conrad & J. T. Lloyd: Characteristic subgroups of lattice ordered groups; Trans. Amer. Math. Soc., 158 (1971), 339–371.



- [2] *Paul Conrad*: Epi-archimedean groups; Czech. Math. Journal, 24 (99), (1974) 192–218.
- [3] *Paul Conrad*: Lattice-ordered Groups; Tulane University, 1970.
- [4] *Laszlo Fuchs*: Partially Ordered Algebraic Systems; Pergamon Press, Addison Wesley, New York 1963.
- [5] *Jorge Martinez*: Free products of abelian  $l$ -groups; Czech. Math. Journal, 23 (98), (1973), 349–361.
- [6] *Jorge Martinez*: Torsion theory for lattice-ordered groups; Czech. Math. Journal 25 (100), (1975), 284–299.

*Author's address*: Department of Mathematics, University of Florida, Gainesville, Florida 32611, U.S.A.