

Zbigniew Sawoń

Some remarks about the three spaces problem in Banach algebras

Czechoslovak Mathematical Journal, Vol. 28 (1978), No. 1, 56–58

Persistent URL: <http://dml.cz/dmlcz/101513>

Terms of use:

© Institute of Mathematics AS CR, 1978

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SOME REMARKS
ABOUT THE THREE SPACES PROBLEM IN BANACH ALGEBRAS

ZBIGNIEW SAWOŃ, Warszawa

(Received November 26, 1975)

Let X be a Banach algebra, not necessarily commutative, with a unit element, and let I be a closed two-sided ideal in X . Thus X/I is also a Banach algebra with unit. Suppose (α) is some property of Banach algebras; we shall say that (α) is a regular property if the fact that I and X/I have property (α) implies that X has property (α) . In the present paper we shall show the following properties to be regular:

- (α_1) X is a B^* -equivalent algebra;
- (α_2) X is a function algebra.

1. Let X be a Banach algebra with involution and a unit element. X is a B^* -equivalent algebra if it admits a norm (defining the correct topology) which is submultiplicative and for which

$$|x^*x| = |x|^2 \quad \text{for all } x \in X.$$

It is well-known [1] that a Banach $*$ -algebra X is B^* -equivalent if and only if there exists a constant $C > 0$ such that

$$\|x^*x\| \geq C\|x\|^2$$

for all $x \in X$, in the original norm. Thus, in a B^* -equivalent algebra, $x_n^*x_n \rightarrow 0$ implies $x_n \rightarrow 0$.

Theorem 1.1. *Let X be a Banach algebra with a unit element and involution, let I be a closed $*$ -ideal in X . If I and X/I are both B^* -equivalent algebras (being in a natural way Banach algebras with involution), then X is a B^* -equivalent algebra.*

Proof. Suppose on the contrary that X is not B^* -equivalent. Then there is a sequence (y_n) in X such that $\|y_n\| = 1$ and $y_n^*y_n \rightarrow 0$.

Let $\varkappa : X \rightarrow X/I$ be the quotient map. One has

$$\varkappa(y_n^* y_n) = \varkappa(y_n^*) \varkappa(y_n) = \varkappa(y_n)^* \varkappa(y_n) \rightarrow 0.$$

Hence, since X/I is a B^* -equivalent algebra, $\varkappa(y_n) \rightarrow 0$. Therefore there is a sequence (i_n) in I such that $y_n + i_n^* \rightarrow 0$ in X . Certainly the sequence (i_n^*) must be bounded and, therefore, (i_n) is bounded since the involution is continuous on I . Hence

$$(y_n i_n)^* (y_n i_n) = i_n^* y_n^* y_n i_n \rightarrow 0.$$

Since I is a B^* -equivalent algebra, $y_n i_n \rightarrow 0$. But now

$$i_n^* i_n = (y_n + i_n^*) i_n - y_n i_n \rightarrow 0$$

(for (i_n) is bounded and $y_n + i_n^* \rightarrow 0$). Therefore, $i_n \rightarrow 0$, since I is a B^* -equivalent algebra.

However, it is impossible that $i_n^* \rightarrow 0$ because $\|y_n\| = 1$ and $y_n + i_n^* \rightarrow 0$. So it must be the case that X is a B^* -equivalent algebra, which proves Theorem 1.1.

2. Now let X be a commutative Banach algebra with a unit element. X is called a *function algebra* if the norm in X is equivalent to the spectral norm $\|\cdot\|_{\text{sp}}$, defined by

$$\|x\|_{\text{sp}} = \lim_{n \rightarrow \infty} \|x^n\|^{1/n} = \sup_{f \in \mathfrak{M}(X)} |f(x)|$$

(where $\mathfrak{M}(X)$ is the structure space of X , i.e. the space of all nonzero multiplicative linear functionals).

Lemma 2.1. *The following conditions are equivalent:*

- (i) X is a function algebra,
- (ii) if $x_n \in X$, $n = 1, 2, \dots$, and $x_n^2 \rightarrow 0$, then $x_n \rightarrow 0$,
- (iii) if $x_n \in X$, $n = 1, 2, \dots$, and $x_n^2 \rightarrow 0$, then (x_n) is bounded,
- (iv) there exists $C > 0$ such that $\|x\|^2 \leq C \|x^2\|$ for every $x \in X$.

Proof. The implications (i) \Rightarrow (ii) and (ii) \Rightarrow (iii) are trivial. Suppose that the condition (iii) holds. If (iv) is not true, then there exists a sequence (x_n) in X such that $\|x_n\| = 1$ and $\|x_n\|^2 > n^3 \|x_n^2\|$, $n = 1, 2, \dots$. Then $\|x_n^2 \cdot n^2 / \|x_n\|^2\| < 1/n$ and $(x_n \cdot n / \|x_n\|)^2 \rightarrow 0$, but the sequence $(x_n \cdot n / \|x_n\|)$ is not bounded. It remains to show (iv) \Rightarrow (i), but this is obvious.

Theorem 2.2. *Let X be a commutative*) Banach algebra with a unit element and I a closed ideal in X . If I and X/I are function algebras, so is X . (In other words, the property of being a function algebra is regular.)*

*) Reviewer's remark: this assumption is superfluous.

Proof. We show that the condition (iii) of Lemma 2.1. holds. Suppose instead that $x_n^2 \rightarrow 0$ and the sequence (x_n) is not bounded. We may assume that $\|x_n\| \rightarrow \infty$, and let $y_n = x_n/\|x_n\|$, ($n = 1, 2, \dots$). Then certainly $y_n^2 \rightarrow 0$. If $\varkappa : X \rightarrow X/I$ is the quotient homomorphism, then

$$\varkappa(y_n^2) = (\varkappa(y_n))^2 \rightarrow 0.$$

Therefore, since X/I is a function algebra, $\varkappa(y_n) \rightarrow 0$. Consequently, there exists a sequence (i_n) in I such that $y_n + i_n \rightarrow 0$, and so necessarily (i_n) is bounded. Thus $(i_n y_n)^2 = i_n^2 y_n^2 \rightarrow 0$, and since I is a function algebra, $i_n y_n \rightarrow 0$. Hence

$$i_n^2 = i_n(y_n + i_n) - i_n y_n \rightarrow 0 \quad \text{and} \quad i_n \rightarrow 0.$$

But this is absurd, as $y_n + i_n \rightarrow 0$ and $\|y_n\| = 1$ for all n .

I would like to thank J. ZEMÁNEK for his help in preparation of this paper.

Reference

- [1] *V. Pták*: Banach algebras with involution, *Manuscripta math.* 6 (1972), p. 245—290.

Author's address: 00-901 Warszawa, Pałac Kultury i Nauki, 9 p., Polska (Instytut Matematyki Uniwersytetu Warszawskiego).