

Mary Ellen Rudin
 \aleph -Dowker spaces

Czechoslovak Mathematical Journal, Vol. 28 (1978), No. 2, 324–326

Persistent URL: <http://dml.cz/dmlcz/101534>

Terms of use:

© Institute of Mathematics AS CR, 1978

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

κ -DOWKER SPACES

MARY ELLEN RUDIN, Madison

(Received October 5, 1976)

In a written communication to the Prague Topology Symposium of 1976, K. MORITA proposed the following:

Conjecture 1. *If a Hausdorff space Y has the property that $X \times Y$ is normal for all normal Hausdorff spaces X , then Y is discrete.*

In an abstract and talk at this symposium M. ATSUJI pointed out that Morita's conjecture follows from:

Conjecture 2. *For each infinite cardinal κ , there is a normal Hausdorff space X_κ which has a decreasing family $\{D_\alpha\}_{\alpha < \kappa}$ of closed sets such that $\bigcap_{\alpha < \kappa} D_\alpha = \emptyset$ and, if $\{U_\alpha\}_{\alpha < \kappa}$ is a family of open sets with $D_\alpha \subset U_\alpha$ for each α , then $\bigcap_{\alpha < \kappa} U_\alpha \neq \emptyset$.*

A space X_κ having the properties described in Atsugi's conjecture could be called a κ -Dowker space since X_ω would be an ordinary Dowker space. The purpose of this note is to prove that there are κ -Dowker spaces for all cardinals κ , thus proving conjectures (1) and (2).

I. Assume that κ is an infinite cardinal; we construct X_κ by simply generalizing the construction given in [1] of an ordinary Dowker space.

We begin by choosing an increasing family $\{\lambda_\alpha\}_{\alpha < \kappa}$ of regular cardinals such that $\lambda_0 < \kappa^+$ and $\lambda_\alpha = \lambda_\alpha^\alpha$. Let $\lambda = \sup\{\lambda_\alpha \mid \alpha < \kappa\}$.

Let $F = \{f : \kappa \rightarrow \lambda \mid f(\alpha) \leq \lambda_\alpha \text{ for all } \alpha < \kappa\}$.

Let $G = \{g \in F \mid g(\alpha) < \lambda_\alpha \text{ for all } \alpha < \kappa\}$.

Let $X = X_\kappa = \{f \in F \mid \exists \beta < \kappa \text{ such that } \kappa^+ \leq \text{cf}(f(\alpha)) \leq \lambda_\beta \text{ for all } \alpha < \kappa\}$.

If f and g belong to F we say $g < f$ if $g(\alpha) < f(\alpha)$ for all α , and we say $g \leq f$ if $g(\alpha) \leq f(\alpha)$ for all α . If $g < f$, define $U_{gf} = \{h \in X \mid g < h \leq f\}$. We topologize X by using $\{U_{gf} \mid g < f \text{ in } F\}$ as a basis.

To check that $X_\kappa = X$ has the properties desired in Conjecture (2), the reader acquainted with [1] will probably have no difficulty. Significant changes needed other than replacing ω by κ are indicated below.

II. For each $\alpha < \kappa$, let $D_\alpha = \{f \in X \mid f(\beta) = \lambda_\beta \text{ for all } \beta \leq \alpha\}$. Clearly $\bigcap_{\alpha < \kappa} D_\alpha = \emptyset$ and each D_α is closed. Assume that for each $\alpha < \kappa$, $D_\alpha \subset U_\alpha$ which is open. We want to prove:

Lemma (3). $\bigcap_{\alpha < \kappa} U_\alpha \neq \emptyset$.

As in [1] it would suffice to prove:

Lemma (2). *Suppose that $\alpha < \kappa$. There is a term f of G such that $\{g \in G \cap \bigcap X \mid f < g\} \subset U_\alpha$.*

Since $\{\beta < \alpha\}$ need not be finite if $\kappa \neq \omega$, we need a different proof of Lemma (2) than that given in [1]. This is again relevant in the proof of Lemma (6). In both proofs we make use of $\lambda_\alpha^\kappa = \lambda_\alpha$.

Proof of Lemma (2). For each $\delta < \lambda_\alpha$, we choose $h_\delta \in \prod_{\beta \leq \alpha} \lambda_\beta$ in such a way that $\delta < \lambda_\alpha$ and $h \in \prod_{\beta \leq \alpha} \lambda_\beta$ imply that there is a $\gamma < \lambda_\alpha$ with $\delta < \gamma$ and $h_\gamma = h$. This is possible since $\{\lambda_\beta\}_{\beta \leq \alpha}$ are increasing and $\lambda_\alpha = \lambda_\alpha^\kappa$ imply that λ_α^κ and λ_α^α and $\prod_{\beta \leq \alpha} \lambda_\beta$ all have cardinality λ_α .

Assuming there is no f satisfying Lemma (2), we define terms g_δ and f_δ of G for all $\delta < \lambda_\alpha$ by induction on δ . If f_γ has been defined for all $\gamma < \delta$, define g_δ by $g_\delta(\beta) = h_\delta(\beta)$ for $\beta \leq \alpha$ and $g_\delta(\beta) = \sup \{f_\gamma(\beta) \mid \gamma < \delta\}$ for $\alpha < \beta < \kappa$. Then choose $f_\delta \in (X \cap G) - U_\alpha$ with $g_\delta < f_\delta$ as guaranteed by assumption. Let f be the term of F with $f(\beta) = \lambda_\beta$ for $\beta \leq \alpha$ and $f(\beta) = \sup \{f_\delta(\beta) \mid \delta < \lambda_\alpha\}$ for $\alpha < \beta < \kappa$. Since $f \in D_\alpha, f \in U_\alpha$ and there is $g < f$ with $U_{g,f} \subset U_\alpha$. For $\alpha < \beta < \kappa$, $\{f_\delta(\beta)\}_{\delta < \lambda_\alpha}$ is strictly increasing. So there is a $\delta < \lambda_\alpha$ such that $f_\delta(\beta) > g(\beta)$ for all $\alpha < \beta < \kappa$. Thus there is a $\gamma < \lambda_\alpha$ with $\delta < \gamma$ and $h_\gamma(\beta) = g(\beta)$ for all $\beta \leq \alpha$. But then $f_\gamma \in U_{g,f}$ contradicting $f_\gamma \notin U_\alpha$.

III. It remains to prove that X is normal. We might as well prove that X is collectionwise normal. So assume that \mathcal{H} is a closed discrete family of closed sets. By exactly the same proof given in [1], we can find disjoint open sets separating the members of \mathcal{H} provided we can prove:

Lemma (4). *The intersection of any family of less than κ open sets is open.*

Lemma (5). *Suppose that $t \in F$ and $\kappa^+ \leq \text{cf}(t(\alpha))$ for all $\alpha < \kappa$. There is an $f \in F$ such that $f < t$ and $\{h \in X \mid f < h \leq t\}$ intersects at most one member of \mathcal{H} .*

Lemma (4) is proved exactly as in [1]; and, as in [1], Lemma (5) follows from:

Lemma (6). *Suppose that $\alpha < \kappa$ and define $U_\alpha = \{h \in X \mid h \leq t \text{ and } \text{cf}(h(\beta)) \leq \lambda_\alpha \text{ for all } \beta < \alpha\}$. There is an $f \in F$ such that $f < t$ and $\{h \in U_\alpha \mid f < h\}$ intersects at most one term of \mathcal{H} .*

Proof of Lemma (6). Let

$$N = \{\beta < \kappa \mid \text{cf}(t(\beta)) \leq \lambda_\alpha\} \quad \text{and} \quad M = \{\beta < \kappa \mid \text{cf}(t(\beta)) > \lambda_\alpha\}.$$

As in the proof of Lemma (2), for each $\delta < \lambda_\alpha$ we choose $g_\delta \in \prod_{\beta \in N} t(\beta)$ in such a way that $\delta < \lambda_\alpha$ and $g \in \prod_{\beta \in N} t(\beta)$ imply there is a $\gamma < \lambda_\alpha$ with $\delta < \gamma$ and $g < g_\gamma$. The fact that the cardinality of N is at most κ and λ_α^* is λ_α makes this possible.

Assuming there is no f satisfying Lemma (6), we define $h_\delta \in U_\alpha$, $k_\delta \in U_\alpha$, and $f_\delta \in F$ for each $\delta < \lambda_\alpha$ by induction on δ . Assuming that h_γ and k_γ have been defined for all $\gamma < \delta$, define f_δ by $f_\delta(\beta) = g_\delta(\beta)$ for $\beta \in N$ and $f_\delta(\beta) = \sup(\{h_\gamma(\beta)\}_{\gamma < \delta} \cup \{k_\gamma(\beta)\}_{\gamma < \delta})$ for $\beta \in M$. Then choose h_δ and k_δ to be terms U_α with $f_\delta < h_\delta$ and $f_\delta < k_\delta$ belonging to *different* terms of \mathcal{H} .

Let $f \in F$ be defined by $f(\beta) = t(\beta)$ for $\beta \in N$ and $f(\beta) = \sup\{f_\delta(\beta) \mid \delta < \lambda_\alpha\}$ for $\beta \in M$. There is $g < f$ such that U_{gf} intersects at most one term of \mathcal{H} . Also there is a $\delta < \lambda_\alpha$ with $f_\delta(\beta) > g(\beta)$ for all $\beta \in M$. So there is $\gamma < \lambda_\alpha$ with $\delta < \gamma$ and $f_\gamma(\beta) > g_\gamma(\beta)$ for all $\beta \in N$. Hence $f_\gamma \in U_{gf}$. But $f_\gamma < h_\gamma$ and $f_\gamma < k_\gamma$ and thus $h_\gamma \in U_{gf}$ and $k_\gamma \in U_{gf}$. However this contradicts U_{gf} intersecting only one term of \mathcal{H} .

Bibliography

- [1] *M. E. Rudin*: A normal space X for which $X \times I$ is not normal, *Fund. Math.* **LXXIII** (1971), 179–186.

Author's address: Mathematics Department, University of Wisconsin-Madison, Van Vleck Hall, 480 Lincoln Drive, Madison, Wisconsin 53706, U.S.A.