

Ladislav Nebeský

A note on upper embeddable graphs

Czechoslovak Mathematical Journal, Vol. 33 (1983), No. 1, 37–40

Persistent URL: <http://dml.cz/dmlcz/101853>

Terms of use:

© Institute of Mathematics AS CR, 1983

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON UPPER EMBEDDABLE GRAPHS

LADISLAV NEBESKÝ, Praha

(Received August 20, 1980)

In the present note only graphs in the sense of the book [1] are considered (multiple edges or loops are not allowed). Let G be a graph. Its vertex set, its edge set, and the number of its components will be denoted by $V(G)$, $E(G)$, and $c(G)$, respectively. If U is a nonempty subset of $V(G)$, then $\langle U \rangle_G$ denotes the subgraph of G induced by U , and $E(U, G)$ denotes the set of edges $e \in E(G)$ with the property that e is incident with exactly one vertex of U . Define $\beta(G) = |E(G)| - |V(G)| + c(G)$.

0. Let G be a connected graph. As was shown in [4], for no integer $n > \lceil \beta(G)/2 \rceil$, there exists a 2-cell (or cellular [7]) embedding of G onto the orientable surface of genus n . G is said to be *upper embeddable* if there exists a 2-cell embedding of G onto the orientable surface of genus $\lceil \beta(G)/2 \rceil$. (For various details concerning the concept of upper embeddability and related subjects the reader is referred to [6]).

If H is a graph, then we denote by $b(H)$ the number of components F of H with the property that $\beta(F)$ is odd.

We now state two characterizations of upper embeddable graphs:

Theorem 0. *Let G be a connected graph. Then the following statements are equivalent:*

- (I) G is upper embeddable;
 - (II) there exists a spanning tree T of G such that for at most one component F of $G - E(T)$, $|E(F)|$ is odd;
 - (III) for every $A \subseteq E(G)$,
- $$(*) \quad b(G - A) + c(G - A) - 2 \leq |A|.$$

The equivalence (I) \Leftrightarrow (II) was found independently by Jungerman [2] and Xuong [8]. The equivalence (II) \Leftrightarrow (III) follows immediately from the results proved in [3].

In the present note two results will be deduced from the equivalence (I) \Leftrightarrow (III).

1. Let G be a graph, and let n be a positive integer. We shall say that G is *oddly n -edge-connected* if it is connected, and for every nonempty proper subset U of $V(G)$ with the properties that $\langle U \rangle_G$ is connected, $\beta(\langle U \rangle_G)$ is odd, and no component of $G - E(U, G)$ is a tree, it holds that $|E(U, G)| \geq n$.

Theorem 1. *Every oddly 4-edge-connected graph is upper embeddable.*

Proof. On the contrary, we assume that there exists an oddly 4-edge-connected graph G which is not upper embeddable. It follows from the implication (III) \Rightarrow (I) that there exists $A \subseteq E(G)$ such that (*) does not hold, and that A is minimal in the sense that for every $A' \subseteq E(G)$, if $|A'| < |A|$, then $b(G - A') + c(G - A') - 2 \leq |A'|$. This implies that $c(G - A) = b(G - A)$, and that every component of $G - A$ is an induced subgraph of G . Moreover, we get that $c(G - A) \geq 2$.

Consider an arbitrary component F of $G - A$. It is clear that F is a component of $G - E(V(F), G)$. Since $c(G - A) = b(G - A)$, we have that $\beta(F)$ is odd and that no component of $G - E(U, G)$ is a tree. Since G is oddly 4-edge-connected, $|E(V(F), G)| \geq 4$.

This implies that $2|A| \geq 4b(G - A)$, and thus (*) holds, which is a contradiction. Hence, the theorem follows.

We say that a graph G is cyclically n -edge-connected ($n \geq 1$) if it is connected, and for every nonempty proper subset U of $V(G)$ with the property that neither of the graphs $\langle U \rangle_G$ and $\langle V(G) - U \rangle_G$ is a forest, it holds that $|E(U, G)| \geq n$. It is clear that every cyclically n -edge-connected graph ($n \geq 1$) is oddly n -edge-connected.

Corollary (Payan and Xuong [5]). *If a graph is cyclically 4-edge-connected, then it is upper embeddable.*

Note that Payan and Xuong [5] proved the result in the corollary without utilizing the implication (III) \Rightarrow (I), but their proof is rather difficult. Theorem 1 is stronger than the corollary: the graphs in Figs. 1 and 2 can serve as examples of oddly 4-edge-connected graphs which are not cyclically 4-edge-connected.

2. We shall say that a connected graph G is *absolutely upper embeddable* if every graph which is spanned by G is upper embeddable. According to the definition, every absolutely upper embeddable graph is upper embeddable. The trees of diameter ≥ 5 and the graph in Fig. 1 can serve as examples of upper embeddable graphs which are not absolutely upper embeddable.

Let H be a graph. We denote by $i(H)$ the number of components F of H with the property that either $\beta(F)$ is odd or F is a non-complete graph. Obviously, $i(H) \geq b(H)$.

The following theorem gives a characterization of absolutely upper embeddable graphs:

Theorem 2. *A connected graph G is absolutely upper embeddable if and only if*

$$(**) \quad i(G - A) + c(G - A) - 2 \leq |A|,$$

for every $A \subseteq E(G)$.

Proof. (1) We first assume that there exists $A \subseteq E(G)$ such that (**) does not hold. We shall assume that A is minimal in the sense that for every $A_0 \subseteq E(G)$, if $|A_0| < |A|$, then $i(G - A_0) + c(G - A_0) - 2 \leq |A_0|$. This implies that every component of $G - A$ is an induced subgraph of G . We wish to prove that G is not absolutely upper embeddable.

Consider a graph H obtained from G in such a way that one new edge is inserted into each component F of $G - A$ with the property that F is non-complete and $\beta(F)$ is even. Since every component of $G - A$ is an induced subgraph of G , no new edge of H belongs to A . Since $b(H - A) = i(G - A)$ and $c(H - A) = c(G - A)$, we have that $b(H - A) + c(H - A) - 2 > |A|$. According to the implication (I) \Rightarrow (III), H is not upper embeddable. The desired result follows.




Fig. 1.




Fig. 2.

(2) We now assume that G is not absolutely upper embeddable. We wish to prove that there exists $A \subseteq E(G)$ such that $(**)$ does not hold. There exists a graph H' which is spanned by G and which is not upper embeddable. According to the implication (III) \Rightarrow (I), there exists $A' \subseteq E(H')$ such that $b(H' - A') + c(H' - A') - 2 > |A'|$. Put $A = A' \cap E(G)$.

Consider an arbitrary component F' of H' . Obviously, $b(F') + c(F') \leq 2$. Denote $F = \langle V(F') \rangle_G$. If $c(F) = 1$, then $i(F) \geq i(F') \geq b(F')$, and thus $i(F) + c(F) \geq b(F') + c(F')$. If $c(F) \geq 2$, then $i(F) + c(F) \geq c(F) \geq b(F') + c(F')$.

This observation implies that $(**)$ does not hold, which completes the proof.

It can be easily deduced from Theorem 2 that the graph in Fig. 2 is absolutely upper embeddable.

References

- [1] M. Behzad, G. Chartrand and L. Lesniak-Forster: *Graphs & Digraphs*. Prindle, Weber & Schmidt, Boston 1979.

- [2] *M. Jungerman*: A characterization of upper embeddable graphs. *Trans. Amer. Math. Soc.* 241 (1978), 401—406.
- [3] *L. Nebeský*: A new characterization of the maximum genus of a graph. *Czechoslovak Math. J.* 31 (106) (1981), 604—613.
- [4] *E. A. Nordhaus, B. M. Stewart and A. T. White*: On the maximum genus of a graph. *J. Combinatorial Theory* 11B (1971), 258—267.
- [5] *C. Payan and N. H. Xuong*: Upper embeddability and connectivity of graphs. *Discrete Mathematics* 27 (1979), 71—80.
- [6] *R. D. Ringelsen*: Survey of results on the maximum genus of a graph. *J. Graph Theory* 3 (1979), 1—13.
- [7] *G. Ringel*: *Map Color Theorem*. Springer-Verlag, Berlin 1974.
- [8] *N. H. Xuong*: How to determine the maximum genus of a graph. *J. Combinatorial Theory* 26B (1979), 217—225.

Added in proofs. The equivalence (I) \Leftrightarrow (II) also follows immediately from the results in *N. P. Homenko, N. A. Ostroverkhy, and V. A. Kusmenko*: The maximum genus of graphs (in Ukrainian, English summary). φ -peretvorenniya grafiv (N. P. Homenko, ed.), IM AN URSR, Kiev 1973, pp. 180—210.

Author's address: 116 38 Praha 1, nám. Krasnoarmějců 2, ČSSR (Filozofická fakulta Univerzity Karlovy).