

Michal Horák; Ivan Kolář

On the higher order Poincaré-Cartan forms

*Czechoslovak Mathematical Journal*, Vol. 33 (1983), No. 3, 467–475

Persistent URL: <http://dml.cz/dmlcz/101896>

## Terms of use:

© Institute of Mathematics AS CR, 1983

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE HIGHER ORDER POINCARÉ-CARTAN FORMS

MICHAL HORÁK and IVAN KOLÁŘ, Brno

(Received April 20, 1982)

This paper is intended to clarify some basic geometric structures related with the higher order Hamilton formalism in arbitrary fibred manifolds. Using a suitable generalization of the formal exterior differentiation, we show that any  $r$ -th order Lagrangian determines a family of the Poincaré-Cartan forms, which is reduced to a single form only if  $r \leq 2$  or the base manifold  $X$  is one-dimensional. For  $\dim X = 1$ , we then derive a relation generalizing the so-called basic theorem of the first order Hamilton formalism in fibred manifolds deduced by Goldschmidt and Sternberg, [2]. — Our consideration is in the category  $C^\infty$ .

**1. A general form of the variational formula.** Given any fibred manifold  $\pi: Y \rightarrow X$ , we denote by  $\pi_r: J^r Y \rightarrow X$  its  $r$ -th jet prolongation and by  $\pi_r^s: J^r Y \rightarrow J^s Y$ ,  $0 \leq s < r$ , ( $J^0 Y = Y$ ) the jet projections. All morphisms are assumed to be base-preserving.

For any morphism  $\varphi: J^r Y \rightarrow \wedge^k T^* X$ , one defines its formal exterior differential  $D\varphi: J^{r+1} Y \rightarrow \wedge^{k+1} T^* X$  by

$$(1) \quad D\varphi(j^{r+1}s) = d(\varphi \circ j^r s)$$

for every section  $s$  of  $Y$ , [3], [8]. If

$$x^i, y^p, i, j, \dots = 1, \dots, n = \dim X, \quad p = 1, \dots, m = \dim Y - \dim X,$$

are some local fibre coordinates on  $Y$ ,  $y_j^p, \dots, y_{j_1 \dots j_r}^p$  are the induced coordinates on  $J^r Y$  and the coordinate expression of  $\varphi$  is

$$(2) \quad \varphi \equiv a_{i_1 \dots i_k}(x^i, y^p, y_j^p, \dots, y_{j_1 \dots j_r}^p) dx^{i_1} \wedge \dots \wedge dx^{i_k},$$

then

$$(3) \quad D\varphi \equiv D_l a_{i_1 \dots i_k} dx^l \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k},$$

provided

$$(4) \quad D_l f = \partial_l f + (\partial_p f) y_l^p + \dots + (\partial_p^{j_1 \dots j_r} f) y_{j_1 \dots j_r}^p$$

with  $\partial_l = \partial/\partial x^l$ ,  $\partial_p = \partial/\partial y^p$ ,  $\dots$ ,  $\partial_p^{j_1 \dots j_r} = \partial/\partial y_{j_1 \dots j_r}^p$ , means the formal (or total) derivative of a function  $f: J^r Y \rightarrow \mathbf{R}$ . Clearly,  $DD\varphi = 0$ . It is well-known that

$J^{r+1}Y \rightarrow J^rY$  is an affine bundle, whose associated vector bundle is the pullback of  $VY \otimes S^{r+1}T^*X$  over  $J^rY$ , where  $VY$  means the vertical tangent bundle of  $Y$  and  $S$  denotes the symmetric tensor product. By (3) and (4),  $D\varphi: J^{r+1}Y \rightarrow \Lambda^{k+1}T^*X$  is an affine morphism for every  $\varphi$ .

Given a projectable vector field  $\eta$  on  $Y$ , we denote by  $J^r\eta$  the vector field on  $J^rY$  induced by means of flows

$$(5) \quad \exp(tJ^r\eta) = J^r(\exp t\eta).$$

If  $\eta \equiv \eta^p(x, y) \partial_p$  is a vertical vector field, then the coordinate expression of  $J^r\eta$  is

$$(6) \quad J^r\eta \equiv \eta^p \partial_p + D_j \eta^p \partial_p^j + \dots + D_{j_1 \dots j_r} \eta^p \partial_p^{j_1 \dots j_r}.$$

Using (3) and (6), one verifies easily

**Lemma 1.** *For every morphism  $A: J^rY \rightarrow V^*J^qY \otimes \Lambda^k T^*X$  over the identity of  $J^qY$ ,  $q \leq r$ , there exists a unique morphism  $DA: J^{r+1}Y \rightarrow V^*J^{q+1}Y \otimes \Lambda^{k+1}T^*X$  satisfying*

$$(7) \quad \langle DA, J^{q+1}\eta \rangle = D(\langle A, J^q\eta \rangle)$$

for every vertical vector field  $\eta$  on  $Y$ .

In coordinates, if

$$(8) \quad A \equiv (a_{p_i_1 \dots i_k} dy^p + \dots + a_{p_i_1 \dots i_k}^{j_1 \dots j_q} dy_{j_1 \dots j_q}^p) \otimes dx^{i_1} \wedge \dots \wedge dx^{i_k},$$

then

$$(9) \quad DA = (D_l a_{p_i_1 \dots i_k} dy^p + a_{p_i_1 \dots i_k} dy_l^p + \dots + D_l a_{p_i_1 \dots i_k}^{j_1 \dots j_q} dy_{j_1 \dots j_q}^p + a_{p_i_1 \dots i_k}^{j_1 \dots j_q} dy_{j_1 \dots j_q}^l) \otimes dx^l \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

Obviously,  $DA$  is an affine morphism and it holds  $DDA = 0$ .

Define  $K_q^s$  by an exact sequence

$$(10) \quad 0 \rightarrow K_q^s \rightarrow VJ^qY \xrightarrow{V\pi_q^s} VJ^sY \rightarrow 0, \quad s \leq q,$$

where  $V\pi_q^s$  means the vertical tangent map to  $\pi_q^s$ . It is well-known that  $K_q^{q-1}$  is the pullback of  $VY \otimes S^q T^*X$  over  $J^qY$ . We have a sequence of inclusions

$$(11) \quad K_q^{q-1} \rightarrow K_q^{q-2} \rightarrow \dots \rightarrow K_q^0 \rightarrow VJ^qY,$$

which induces the dual sequence of epimorphisms

$$(12) \quad V^*J^qY \rightarrow K_q^{0*} \rightarrow \dots \rightarrow K_q^{q-2*} \rightarrow K_q^{q-1*}.$$

A morphism  $A: J^rY \rightarrow V^*J^qY \otimes \Lambda^k T^*X$  will be called graded if there exist  $A_1, \dots, A_q$  such that the following diagram commutes

$$(13) \quad \begin{array}{ccc} J^r Y & \longrightarrow^{A} & V^* J^q Y \otimes \wedge^k T^* X \\ \downarrow \pi_r^{r-1} & & \downarrow \\ J^{r-1} Y & \longrightarrow^{A_1} & K_q^{0*} \otimes \wedge^k T^* X \\ \vdots & \vdots & \vdots \\ J^{r-q-1} Y & \longrightarrow^{A_{q-1}} & K_q^{q-2*} \otimes \wedge^k T^* X \\ \downarrow \pi_{r-q-1}^{r-q} & & \downarrow \\ J^{r-q} Y & \longrightarrow^{A_q} & K_q^{q-1*} \otimes \wedge^k T^* X \end{array}$$

where the arrows in the second column are the tensor products of (12) and the identity of  $\wedge^k T^* X$ . Clearly, if  $A$  is graded, then  $DA$  is also graded.

We define an  $r$ -th order Lagrangian  $\lambda$  on  $Y$  as a morphism  $\lambda: J^r Y \rightarrow \wedge^n T^* X$ , [3], [4], [8]. In coordinates,

$$\lambda \equiv L(x^i, y^p, y_j^p, \dots, y_{j_1 \dots j_r}^p) dx^1 \wedge \dots \wedge dx^n.$$

Its vertical differential  $VJ^r Y \rightarrow \wedge^n T^* X$  will be interpreted as a map  $\delta\lambda: J^r Y \rightarrow V^* J^r Y \otimes \wedge^n T^* X$ .

**Proposition 1.** *For any  $r$ -th order Lagrangian  $\lambda$  on  $Y$ , there exist a graded morphism  $M: J^{2r-1} Y \rightarrow V^* J^{r-1} Y \otimes \wedge^{n-1} T^* X$  and a unique Euler morphism  $E: J^{2r} Y \rightarrow V^* Y \otimes \wedge^n T^* X$  such that*

$$(14) \quad \delta\lambda = DM + E.$$

*For  $r = 1$  or  $n = 1$ ,  $M$  is uniquely determined. If  $r \geq 2$  and  $n \geq 2$ , any other morphism with this property is of the form  $M + DC$ , where  $C$  is any graded morphism  $C: J^{2r-2} Y \rightarrow V^* J^{r-2} Y \otimes \wedge^{n-2} T^* X$ .*

*Proof.* Write  $\omega_i = (\partial/\partial x^i) \lrcorner dx^1 \wedge \dots \wedge dx^n$  and

$$(15) \quad M = (b_p^i dy^p + b_p^{j_1} dy_{j_1}^p + \dots + b_p^{j_1 \dots j_{r-1}} dy_{j_1 \dots j_{r-1}}^p) \otimes \omega_i,$$

$$(16) \quad E = (e_p dy^p) \otimes dx^1 \wedge \dots \wedge dx^n,$$

so that (14) is equivalent to

$$\begin{aligned} \partial_p^{j_1 \dots j_r} L &= b_p^{(j_1 \dots j_r)} \\ &\vdots \\ \partial_p^{j_1 \dots j_k} L &= D_j b_p^{j_1 \dots j_k j} + b_p^{(j_1 \dots j_k)} \\ &\vdots \\ \partial_p L &= D_j b_p^j + e_p \end{aligned}$$

where the round bracket denotes symmetrization. Therefore,

$$\begin{aligned} b_p^{j_1 \dots j_r} &= \partial_p^{j_1 \dots j_r} L + c_p^{j_1 \dots j_r}, \quad c_p^{j_1 \dots (j_{r-1} j_r)} = 0, \\ &\vdots \\ b_p^{j_1 \dots j_k} &= \partial_p^{j_1 \dots j_k} L - D_j b_p^{j_1 \dots j_k j} + c_p^{j_1 \dots j_k}, \quad c_p^{j_1 \dots (j_{k-1} j_k)} = 0, \\ &\vdots \\ e_p &= \partial_p L - D_j b_p^j, \end{aligned}$$

where  $c_p^{j_1 \dots j_k}$  are any functions on  $J^{2r-k}Y$  antisymmetric in  $j_{k-1}$  and  $j_k$ . Hence  $D_{ij} c_p^{j_1 \dots j_{k-2} ij} = 0$ , which implies that

$$(17) \quad e_p = \partial_p L - D_j \partial_p^j L + \dots + (-1)^r D_{j_1 \dots j_r} \partial_p^{j_1 \dots j_r} L$$

is uniquely determined. Further, the space of all  $c_p^{j_1 \dots j_r}$  is the pullback over  $J^r Y$  of the following vector bundle

$$(18) \quad V^*Y \otimes (S^{q-1}TX \otimes TX \cap S^{q-2}TX \otimes \Lambda^2 TX) \otimes \Lambda^n T^*X$$

with  $q = r$ . Take a global section  $c_r$  of the latter vector bundle and apply induction. By the induction hypothesis, for any  $i = 0, 1, \dots, r - k - 1$  we have considered an affine subbundle of the pullback of  $(K_r^{r-i-1})^* \otimes \Lambda^n T^*X$  over  $J^{r+i}Y$ , the associated vector bundle being the pullback of (18) with  $q = r - i$  over  $J^{r+i}Y$ , and we have constructed a section  $c_{r-i}$  of the latter bundle. In this situation, the space of all  $c_p^{j_1 \dots j_k}$  is an affine subbundle of the pullback of  $(K_r^{k-1})^* \otimes \Lambda^n T^*X$  over  $J^{r+k}Y$ , the associated vector bundle being the pullback of (18) with  $q = k$  over  $J^{r+k}Y$ . Indeed, since the values of  $b_p^{j_1 \dots j_r}, \dots, b_p^{j_1 \dots j_{k+1}}$  are already fixed by means of  $c_r, \dots, c_{k+1}$ , the differences  $c_p^{j_1 \dots j_k}$  lie in the subspace  $(K_k^{k-1})^* \otimes \Lambda^n T^*X \subset (K_r^{k-1})^* \otimes \Lambda^n T^*X$  determined by means of the dual map to the epimorphism  $V\pi_r^k: VJ^r Y \rightarrow VJ^k Y$ . Hence we can construct a global section  $c_k$  of the affine bundle in question and continue in our induction procedure. Finally, we obtain a graded morphism  $M: J^{2r-1}Y \rightarrow V^*J^{r-1}Y \otimes \Lambda^{n-1}T^*X$  satisfying (14). Analyzing this construction, we find easily that any other graded morphism with the same property is of the form  $M + DC$  mentioned in our Proposition, QED.

Any morphism  $M: J^{2r-1}Y \rightarrow V^*J^{r-1}Y \otimes \Lambda^{n-1}T^*X$  such that there is an  $E$  satisfying (14) will be called a morphism associated to  $\lambda$ . From the proof of Proposition 1 we obtain a somewhat stronger result: if  $M_i, i = 1, 2$  are two morphisms associated to  $\lambda$  (not necessarily graded) and if  $E_i$  satisfies (14), then  $E_1 = E_2$ . In other words, the Euler morphism is uniquely determined even if  $M$  is not graded.

**Remark 1.** If we take any vertical vector field  $\eta$  on  $Y, \langle \delta \lambda, J^r \eta \rangle =: \delta_\eta \lambda$  is the classical variation of  $\lambda$  with respect to  $\eta$ . Then (14) implies

$$(19) \quad \delta_\eta \lambda = D \langle M, J^{r-1} \eta \rangle + \langle E, \eta \rangle,$$

which is a variational formula of classical type.

**2. Poincaré-Cartan morphisms.** We shall distinguish a special class of the associated morphisms. For this purpose, we need a modification of our operation  $D$ .

Consider a morphism  $A: J^r Y \rightarrow (K_q^s)^* \otimes \Lambda^k T^* X$  with a coordinate expression

$$A \equiv (a_{p_{i_1 \dots i_k}^{j_1 \dots j_{s+1}}} dy_{j_1 \dots j_{s+1}}^p + \dots + a_{p_{i_1 \dots i_k}^{j_1 \dots j_q}} dy_{j_1 \dots j_q}^p) \otimes dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

As we have no inclusion  $(K_q^s)^* \subset V^* J^q Y$ , we cannot construct  $DA$ . However, one can consider a map  $B: J^r Y \rightarrow V^* J^q Y \otimes \Lambda^k T^* X$  covering  $A$ ,

$$B \equiv (b_{p_{i_1 \dots i_k}^{j_1 \dots j_s}} dy_{j_1 \dots j_s}^p + \dots + a_{p_{i_1 \dots i_k}^{j_1 \dots j_{s+1}}} dy_{j_1 \dots j_{s+1}}^p + a_{p_{i_1 \dots i_k}^{j_1 \dots j_q}} dy_{j_1 \dots j_q}^p) \otimes dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

Denote by  $(DB)_s$  the canonical projection of  $DB$  into  $(K_{q+1}^s)^* \otimes \Lambda^{k+1} T^* X$ , so that

$$(20) \quad (DB)_s \equiv (b_{p_{i_1 \dots i_k}^{j_1 \dots j_s}} dy_{j_1 \dots j_s}^p + D_l a_{p_{i_1 \dots i_k}^{j_1 \dots j_{s+1}}} dy_{j_1 \dots j_{s+1}}^p + \dots + a_{p_{i_1 \dots i_k}^{j_1 \dots j_q}} dy_{j_1 \dots j_q}^p) dx^l \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

This depends on the choice of  $B$  only by the first term. But (20) is an affine map and  $b_{p_{i_1 \dots i_k}^{j_1 \dots j_s}}$  belong to its ‘‘absolute’’ part, so that the linear map associated to (20) is completely determined by  $A$ . The latter map will be denoted by  $\overline{DA}$ .

Consider a graded morphism  $M$  associated to  $\lambda$ . Since  $M$  is graded, the values of  $\overline{D}(M_k)$  lie in the subspace  $(K_k^{k-1})^* \otimes \Lambda^n T^* X = V^* Y \otimes S^k TX \otimes \Lambda^n T^* X$ . Let  $\lrcorner: S^q TX \otimes \Lambda^k T^* X \rightarrow S^{q-1} TX \otimes \Lambda^{k-1} T^* X$  be the standard map, [7]. We shall say that  $M$  is a Poincaré-Cartan (in short: P.-C.) morphism of  $\lambda$ , if it holds

- (i)  $(\text{id} \otimes \lrcorner) \circ M_{r-1} = 0$ ,
- (ii)  $M_{k-1}$  is affine and its associated linear morphism coincides with  $-(\text{id} \otimes \lrcorner) \circ \overline{D}(M_k)$  for all  $k = r-1, \dots, 2$ .

The coordinate meaning of these additional conditions is  $c_p^{j_1 \dots j_r} = 0$  and  $c_p^{j_1 \dots j_k}$  are some functions defined on  $J^{2r-k-1} Y$  (and not on  $J^{2r-k} Y$  as in the general case). Quite similarly to Proposition 1, one deduces

**Proposition 2.** *For any  $r$ -th order Lagrangian  $\lambda$  on  $Y$  there exists a P.-C. morphism  $M$ . This morphism is unique if  $r = 1, 2$  or  $n = 1$ . If  $r \geq 3$  and  $n \geq 2$ , any other P.-C. morphism of  $\lambda$  is of the form  $M + DC$ , where  $C$  is any graded morphism  $C: J^{2r-3} Y \rightarrow V^* J^{r-3} Y \otimes \Lambda^{n-2} T^* X$ .*

**Remark 2.** If the base manifold  $X$  is an affine space, we have a map  $a: VJ^{r-1} Y \otimes T^* X \rightarrow VJ^r Y$  defined as follows. Any  $b \in T_x^* X$ ,  $x \in X$ , determines a unique affine map  $f: X \rightarrow R$  such that  $b = j_x^1 f$ . Using the well-known identification  $VJ^{r-1} Y \approx J^{r-1} VY$ , [2], we can express any  $u \in (VJ^{r-1} Y)_x$  as  $u = j_x^{r-1} \sigma$ , where  $\sigma$  is a local section of  $VY$ . Then  $j_x^r(f \cdot \sigma) \in J^r VY \approx VJ^r Y$  is completely determined by  $b$  and  $u$ . This map is bilinear and induces  $a$ . Having any manifold  $Q$  and any map  $F: Q \rightarrow$

$\rightarrow V^*J^rY \otimes \wedge^{n-1}T^*X$ , we can now construct the following maps

$$\begin{array}{c} Q \rightarrow^F V^*J^rY \otimes \wedge^{n-1}T^*X \\ \downarrow a^* \otimes \text{id} \\ V^*J^{r-1}Y \otimes TX \otimes \wedge^{n-1}T^*X \\ \downarrow \text{id} \otimes \lrcorner \\ V^*J^{r-1}Y \otimes \wedge^{n-2}T^*X. \end{array}$$

If the resulting map vanishes,  $F$  will be called quasi-symmetric. Having a coordinate expression of  $F$  of the form (15),  $F$  is quasi-symmetric iff all  $b$ 's are symmetric in all superscripts. Analyzing the proof of Proposition 1, we deduce easily that there is a unique quasi-symmetric P.-C. morphism of  $\lambda$ , which will be called the affine P.-C. morphism of  $\lambda$ . Its coordinate expression is

$$(21) \quad [(\partial_p^{j_1 \dots j_{r-1} L}) dy_{j_1 \dots j_{r-1}}^p + (\partial_p^{j_1 \dots j_{r-2} L} - D_j \partial_p^{j_1 \dots j_{r-2} j L}) dy_{j_1 \dots j_{r-2}}^p + \dots \\ \dots + (\partial_p^j L - D_j \partial_p^j L + \dots + (-1)^{r-1} D_{j_1 \dots j_{r-1}} \partial_p^{j_1 \dots j_{r-1} L}) dy^p] \otimes \omega_l.$$

**3. Transfer to exterior forms.** We recall that an exterior  $k$ -form  $\omega$  on  $J^rY$  is called contact, if  $(J^r s)^* \omega = 0$  for every section of  $Y$ . A contact  $k$ -form  $\omega$  is said to be 2-contact, if  $\zeta \lrcorner \omega$  is a contact  $(k-1)$ -form for every vertical vector field  $\zeta$  on  $J^rY$ .

Any morphism  $\varphi: J^rY \rightarrow \wedge^k T^*X$  can be canonically interpreted as an exterior  $k$ -form on  $J^rY$ , which will be denoted by  $\tilde{\varphi}$ . By the very definition of the formal exterior differentiation,

$$(22) \quad \Delta\varphi := d(\tilde{\varphi}) - (D\varphi)^\sim$$

is a contact  $(k+1)$ -form on  $J^{r+1}Y$ . In particular, any function  $f: J^rY \rightarrow \mathbf{R}$  determines a contact 1-form  $\Delta f$  on  $J^{r+1}Y$ .

**Lemma 2.** For every morphism  $A: J^rY \rightarrow V^*J^qY \otimes \wedge^k T^*X$  and every vertical vector field  $\zeta$  on  $J^qY$ ,

$$(23) \quad (D\langle A, \zeta \rangle)^\sim - (d(\langle A, \zeta \rangle))^\sim$$

is a contact form.

*Proof.* If  $\zeta^p \partial_p + \dots + \zeta_{j_1 \dots j_q}^p \partial_p^{j_1 \dots j_q}$  is the coordinate expression of  $\zeta$ , then the coordinate expression of (23) is

$$\begin{aligned} & [(\Delta a_{p i_1 \dots i_k}) \zeta^p + a_{p i_1 \dots i_k} \Delta \zeta^p + \dots + (\Delta a_{p i_1 \dots i_k}^{j_1 \dots j_q}) \zeta_{j_1 \dots j_q}^p + \\ & + a_{p i_1 \dots i_k}^{j_1 \dots j_q} \Delta \zeta_{j_1 \dots j_q}^p] \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}. \end{aligned}$$

Consider the canonical map  $\psi_q: TJ^{q+1}Y \rightarrow VJ^qY$  (called the structure form of  $TJ^{q+1}Y$ ), [1], [2]. The coordinate expression of  $\psi_q$  is

$$(\Delta y^p) \partial_p + \dots + (\Delta y_{j_1 \dots j_q}^p) \partial_p^{j_1 \dots j_q}.$$

For every  $A: J^rY \rightarrow V^*J^qY \otimes \wedge^k T^*X$ ,  $r > q$ , we define a  $(k+1)$ -form  $\psi_q \bar{\wedge} A$  on  $J^rY$  by the natural combination of the contraction with respect to  $VJ^qY$  and alter-

nation. The coordinate expression of  $\psi_q \bar{\wedge} A$  is

$$(24) \quad (a_{p_1 \dots i_k} \Delta y^p + \dots + a_{p_1 \dots i_k}^{j_1 \dots j_q} \Delta y_{j_1 \dots j_q}^p) \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

Obviously, it holds

$$(25) \quad \zeta \lrcorner (\psi_q \bar{\wedge} A) = (\langle A, \zeta \rangle)^\sim$$

for every vertical vector field  $\zeta$  on  $J^q Y$ . (As  $\psi_q \bar{\wedge} A$  is a  $\pi_r^q$ -horizontal form,  $\zeta \lrcorner (\psi_q \bar{\wedge} A)$  has a well-defined meaning even though  $\zeta$  is a vector field on  $J^q Y$  and not on  $J^r Y$ .)

**Lemma 3.** *The following form is 2-contact*

$$(26) \quad (\psi_{q+1} \bar{\wedge} DA) + d(\psi_q \bar{\wedge} A).$$

Proof. The coordinate expression of (26) is

$$(\Delta y^p \wedge \Delta a_{p_1 \dots i_k} + \dots + \Delta y_{j_1 \dots j_q}^p \wedge \Delta a_{p_1 \dots i_k}^{j_1 \dots j_q}) \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

Set  $\varepsilon = \psi_1 \bar{\wedge} E$ , which is an  $(n+1)$ -form on  $J^{2r} Y$ .

**Proposition 3.** *For any morphism  $M$  associated to  $\lambda$ , any vertical vector field  $\eta$  on  $Y$  and any section  $s$  of  $Y$ , it holds*

$$(27) \quad (j^r s)^* (J^r \eta \lrcorner d\tilde{\lambda}) = (j^{2r-1} s)^* d(J^{r-1} \eta \lrcorner (\psi_{r-1} \bar{\wedge} M)) + (j^{2r} s)^* (\eta \lrcorner \varepsilon).$$

Proof. Let us start from (19). Obviously,  $\langle \delta\lambda, J^r \eta \rangle = J^r \eta \lrcorner d\tilde{\lambda}$  and  $(\langle E, \eta \rangle)^\sim = \eta \lrcorner \varepsilon$ . By Lemma 2,  $(j^{2r-1} s)^* (\langle D\langle M, J^{r-1} \eta \rangle \rangle)^\sim = (j^{2r-1} s)^* (d\langle M, J^{r-1} \eta \rangle)^\sim$ . Using (25), we obtain  $(\langle M, J^{r-1} \eta \rangle)^\sim = J^{r-1} \eta \lrcorner (\psi_{r-1} \bar{\wedge} M)$ , which proves (27).

For any morphism  $M$  associated to  $\lambda$ , the  $n$ -form  $\Theta = \tilde{\lambda} + \psi_{r-1} \bar{\wedge} M$  will be called a form associated to  $\lambda$ .

**Proposition 4.** *Let  $\Theta$  be a form associated to  $\lambda$ . Then  $\zeta \lrcorner d\Theta$  is a contact form for any  $\pi_{2r-1}^0$ -vertical vector field  $\zeta$  on  $J^{2r-1} Y$ .*

Proof. By (14),  $\psi_r \bar{\wedge} \delta\lambda = \psi_r \bar{\wedge} DM + \psi_r \bar{\wedge} E$ . Hence  $(\zeta \lrcorner \delta\lambda)^\sim - (\zeta \lrcorner DM)^\sim = (\zeta \lrcorner E)^\sim = 0$  as  $\zeta$  is  $\pi_{2r-1}^0$ -vertical. By Lemma 3, the form  $\psi_r \bar{\wedge} DM + d(\psi_{r-1} \bar{\wedge} M)$  is 2-contact, so that  $(\zeta \lrcorner DM)^\sim + \zeta \lrcorner d(\psi_{r-1} \bar{\wedge} M)$  is a contact form. Since  $(\zeta \lrcorner \delta\lambda)^\sim = \zeta \lrcorner d\tilde{\lambda}$ , the form  $\zeta \lrcorner d\Theta$  is also contact, QED.

If  $M$  is a P.-C. morphism of  $\lambda$ , then the corresponding form  $\Theta = \tilde{\lambda} + \psi_{r-1} \bar{\wedge} M$  will be called a P.-C. form of  $\lambda$ . If the base manifold is an affine space, then the P.-C. form corresponding to the affine P.-C. morphism will be also said to be affine. Such a form was considered by Krupka, [5]. However, we remark that for  $r \geq 3$  and  $n \geq 2$ , the coordinate expressions corresponding to (21), which are used in [5], have no intrinsic meaning in the case of an arbitrary fibred manifold.



**4. On the higher order Hamilton formalism.** Let  $L_r(Y)$  denote the pullback of  $V^*Y \otimes S^rTX \otimes \wedge^r T^*X$  over  $J^{r-1}Y$ . This vector bundle will be said to be the  $r$ -th Legendre bundle of  $Y$ . The restriction of  $\delta\lambda$  to  $K_r^{r-1}$  can be interpreted as a map  $\sigma : J^rY \rightarrow L_r(Y)$ , which will be called the Legendre transformation of  $\lambda$ . If  $w_p^{j_1 \dots j_r}$  are the natural fibre coordinates on  $L_r(Y)$ , then the equations of  $\sigma$  are

$$w_p^{j_1 \dots j_r} = \partial_p^{j_1 \dots j_r} L.$$

A Lagrangian will be called regular, if its Legendre transformation is a local diffeomorphism. (The general idea of a regular higher order Lagrangian is due to D. Krupka and M. Francaviglia, [9].) Hence  $\lambda$  is regular iff  $\partial_p^{j_1 \dots j_r} \partial_q^{k_1 \dots k_r} L$  is a regular matrix.

**Lemma 4.** For  $\dim X = 1$ , if  $\Theta$  is the P.-C. form of a regular Lagrangian and a section  $u : X \rightarrow J^{2r-1}Y$  satisfies

$$(28) \quad u^*(\zeta \lrcorner d\Theta) = 0$$

for every  $\pi_{2r-1}^0$ -vertical vector field  $\zeta$  on  $J^{2r-1}Y$ , then  $u$  is holonomic, i.e. there is a section  $s : X \rightarrow Y$  such that  $u = j^{2r-1}s$ . If  $r = 1$  the same result holds for any  $\dim X$ .

*Proof.* We shall use local coordinates. For  $\dim X = 1$ , we shall write  $y_{j_1 \dots j_r}^p = y_{(r)}^p$  and  $L_{pq} = \partial_p^{j_1 \dots j_r} \partial_q^{k_1 \dots k_r} L$ . Let  $\zeta = \zeta_{(1)}^p \partial_p^{(1)} + \dots + \zeta_{(2r-1)}^p \partial_p^{(2r-1)}$ . Then (28) leads to the following sequence of equations

$$\begin{aligned} u^*(L_{pq} \zeta_{(2r-1)}^p \Delta y^q) &= 0, \quad \text{which gives} \quad u^* \Delta y^q = 0, \\ &\vdots \\ u^*(L_{pq} \zeta_{(1)}^p \Delta y_{(2r-2)}^q) &= 0, \quad \text{which gives} \quad u^* \Delta y_{(2r-2)}^q = 0. \end{aligned}$$

Thus,  $u$  is holonomic. If  $r = 1$ , the same result is deduced for any  $\dim X$  in [3], QED.

**Lemma 5.** For any contact  $k$ -form  $\omega$  on  $J^rY$ , any projectable vector field  $\eta$  on  $Y$  and any section  $s$  of  $Y$ , it holds

$$(29) \quad (j^r s)^* d(J^r \eta \lrcorner \omega) = -(j^r s)^* (J^r \eta \lrcorner d\omega).$$

*Proof.* Since  $\omega$  vanishes on the  $r$ -jet prolongation of any section of  $Y$  and the flow of  $J^r \eta$  transforms any  $r$ -jet prolongation of a section into an  $r$ -jet prolongation of a section, the Lie derivative  $L_{J^r \eta} \omega$  is also a contact form. Using the standard formula for the Lie derivative

$$L_{J^r \eta} \omega = J^r \eta \lrcorner d\omega + d(J^r \eta \lrcorner \omega),$$

we find (29), QED.

**Proposition 5.** If  $\Theta$  is any form associated to  $\lambda$  and  $s : X \rightarrow Y$  is a section, then the

condition  $(j^{2r-1}s)^*(\zeta \lrcorner d\Theta) = 0$  for any  $\pi_{2r-1}$ -vertical vector field  $\zeta$  on  $J^{2r-1}Y$  is equivalent to  $(j^{2r}s)^*E = 0$ .

*Proof.* Consider first a section  $s$  of  $Y$  and set  $\zeta_s = \zeta|_{j^{2r-1}s}$ . Then the projection  $\eta_s = T\pi_{2r-1}^0(\zeta_s)$  can be extended to a vertical vector field  $\eta$  on  $Y$ . By construction,  $(J^{2r-1}\eta - \zeta)|_{j^{2r-1}s}$  is  $\pi_{2r-1}^0$ -vertical, so that  $(j^{2r-1}s)^*(\zeta \lrcorner d\Theta) = (j^{2r-1}s)^* \cdot (J^{2r-1}\eta \lrcorner d\Theta)$  by Proposition 4. According to Lemma 5 and Proposition 3,  $(j^{2r-1}s)^*(J^{2r-1}\eta \lrcorner d\Theta) = (j^{2r}s)^*(\eta \lrcorner \varepsilon)$ . Hence  $(j^{2r}s)^*E = 0$  is equivalent to  $(j^{2r-1}s)^*(\zeta \lrcorner d\Theta) = 0$ , QED.

The following result generalizes the so-called basic theorem of the first order Hamilton formalism [2], [3], (Another geometrical treatment of the first order case can be found in a paper by Ragionieri and Ricci, [6].)

**Theorem.** For  $\dim X = 1$ , let  $\Theta$  be the Poincaré-Cartan form of a regular  $r$ -th order Lagrangian on  $Y$  and  $u: X \rightarrow J^{2r-1}Y$  any section. Then the equation  $u^*(\zeta \lrcorner d\Theta) = 0$  for every  $\pi_{2r-1}$ -vertical vector field  $\zeta$  on  $J^{2r-1}Y$  is equivalent to a pair of conditions

$$u = j^{2r-1}s \text{ for a section } s \text{ of } Y \text{ and } (j^{2r}s)^*E = 0.$$

The same result holds for  $r = 1$  and any  $\dim X$ .

*Proof.* By Lemma 4,  $u^*(\zeta \lrcorner d\Theta)$  implies  $u = j^{2r-1}s$ . Then our Theorem follows from Proposition 5, QED.

#### References

- [1] P. L. Garcia: The Poincaré-Cartan invariant in the calculus of variations, Istituto Nazionale di Alta Matematica, Symposia Matematica XIV, Roma 1974, 219–246.
- [2] H. Goldschmidt, S. Sternberg: The Hamilton formalism in the calculus of variations, Ann. Inst. Fourier (Grenoble), 23 (1973), 203–267.
- [3] I. Kolář: On the Hamilton formalism in fibered manifolds, Scripta Fac. Sci. Nat. UJEP Brunensis, Physica 3–4, 5 (1975), 249–254.
- [4] I. Kolář: Lie derivatives and higher order Lagrangians, Proceedings of the conference on differential geometry and its applications, Universita Karlova Praha 1982, 117–123.
- [5] D. Krupka: Natural Lagrangian structures, to appear in Banach Center Publications, Warsaw.
- [6] R. Ragionieri, R. Ricci: Hamiltonian formalism in the calculus of variations, Bolletino U.M.I. (5) 18-B(1981), 119–130.
- [7] S. Sternberg: Lectures on differential geometry, Prentice-Hall, New Jersey, 1964.
- [8] A. Trautman: Invariance of Lagrangian systems, General Relativity, Papers in honour J. L. Synge, Clarendon Press, Oxford 1972, 85–99.
- [9] M. Francaviglia, D. Krupka: The Hamilton formalism in higher order variational problems, to appear in Ann. Inst. H. Poincaré.

*Authors address:* 603 00 Brno, Mendlovo nám. 1, ČSSR (Matematický ústav ČSAV, branch Brno).