

Ivan Chajda; Bohdan Zelinka

Directly decomposable tolerances on direct products of lattices and semilattices

*Czechoslovak Mathematical Journal*, Vol. 33 (1983), No. 4, 519–521

Persistent URL: <http://dml.cz/dmlcz/101907>

## Terms of use:

© Institute of Mathematics AS CR, 1983

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

DIRECTLY DECOMPOSABLE TOLERANCES  
ON DIRECT PRODUCTS OF LATTICES AND SEMILATTICES

IVAN CHAJDA, Přešov, and BOHDAN ZELINKA, Liberec

(Received May 8, 1970)

L 8

A *compatible tolerance* on an algebra  $\mathfrak{A} = \langle A, \mathcal{F} \rangle$  is a reflexive and symmetric binary relation on  $A$  having the Substitution Property with respect to all operations of  $\mathcal{F}$  with non-zero arities. In particular, if  $L$  is a lattice and  $T$  is a reflexive and symmetric relation on the support of  $L$ , then  $T$  is a compatible tolerance on  $L$  if and only if  $(a, b) \in T, (c, d) \in T$  always imply  $(a \vee c, b \vee d) \in T$  and  $(a \wedge c, b \wedge d) \in T$ .

Let  $L_\gamma$  be lattices for  $\gamma \in \Gamma$  and  $L = \prod_{\gamma \in \Gamma} L_\gamma$ , let  $T$  be a compatible tolerance on  $L$ . The tolerance  $T$  is called *directly decomposable* if there exist compatible tolerances  $T_\gamma$  on  $L_\gamma$  (for each  $\gamma \in \Gamma$ ) such that  $T = \prod_{\gamma \in \Gamma} T_\gamma$  (this means that  $(a, b) \in T$  if and only if  $(pr_\gamma a, pr_\gamma b) \in T_\gamma$  for each  $\gamma \in \Gamma$ , where  $pr_\gamma$  means the projection onto the  $\gamma$ -th direct factor of  $L$ ).

Some conditions for the direct decomposability of compatible tolerances were investigated in [4]. The paper [3] contains the complete solution of the problem of direct decomposability of compatible tolerances on lattices in the case of  $\Gamma$  finite. If  $\mathfrak{A}_\gamma \in \mathcal{V}$  for finite  $\Gamma$  and a given variety  $\mathcal{V}$ , the problem is completely solved in [2]. However, the case of  $\Gamma$  infinite has still remained open.

An analogous problem for congruences on (infinite) direct products of lattices was partially solved in [1] and [5]. We shall use the methods from [1] and [5] to obtain a similar result in the case of tolerances. Let us recall some notions from [5]. Suppose  $L = \prod_{\gamma \in \Gamma} L_\gamma$  and  $x \in L, y \in L$ . Denote  $x(\gamma) = pr_\gamma x$ . Further denote by  $f(x, y, \gamma)$  the element of  $L$  defined by

$$\begin{aligned} f(x, y, \gamma)(\gamma) &= x(\gamma), \\ f(x, y, \gamma)(\delta) &= y(\delta) \quad \text{for } \delta \in \Gamma, \delta \neq \gamma. \end{aligned}$$

The following lemma is a generalization of that in [5]; the application of transitivity in that proof is avoided.

**Lemma.** *Let  $T$  be a compatible tolerance on the lattice  $L = \prod_{\gamma \in \Gamma} L_\gamma$ , let  $x \in L, y \in L, (x, y) \in T$ . Then  $(f(x, z, \gamma), f(y, z, \gamma)) \in T$  for each  $z \in L$  and each  $\gamma \in \Gamma$ .*

**Proof.** Clearly  $x \wedge y \leq f(y, x, \gamma) \leq x \vee y$ . Hence  $(x, y) \in T$  implies  $(x \wedge y, x \vee y) \in T$  and  $(x, f(y, x, \gamma)) \in T$  in virtue of the convexity of  $T$ . By the Substitution Property we have

$$(x \wedge f(x, z, \gamma), f(y, x, \gamma) \wedge f(x, z, \gamma)) \in T.$$

Since

$$f(a_1, b_1, \gamma) \wedge f(a_2, b_2, \gamma) = f(a_1 \wedge a_2, b_1 \wedge b_2, \gamma),$$

we obtain

$$(f(x, z \wedge x, \gamma), f(x \wedge y, z \wedge x, \gamma)) \in T.$$

By using the operation  $\vee$  and the pair  $(f(y, z, \gamma), f(y, z, \gamma)) \in T$ , we obtain  $(f(x \vee y, z, \gamma), f(y, z, \gamma)) \in T$ . Analogously we can prove  $(f(x, z, \gamma), f(x \vee y, z, \gamma)) \in T$ . Using the operation  $\wedge$ , we obtain  $(f(x, z, \gamma), f(y, z, \gamma)) \in T$ , which was to be proved.

Let  $T$  be a compatible tolerance on a lattice  $L$  and let  $m$  be a given infinite cardinal number. The tolerance  $T$  is called *conditionally  $\vee$ - $m$ -complete*, if  $(a_\delta, b_\delta) \in T$  for  $\delta \in \Delta$  with  $|\Delta| = m$  imply  $(\bigvee_{\delta \in \Delta} a_\delta, \bigvee_{\delta \in \Delta} b_\delta) \in T$  provided that both  $\bigvee_{\delta \in \Delta} a_\delta$  and  $\bigvee_{\delta \in \Delta} b_\delta$  exist in  $L$ . Dually we can define a *conditionally  $\wedge$ - $m$ -complete tolerance*.

**Theorem 1.** *Each conditionally  $\vee$ - $m$ -complete tolerance on the lattice  $L = \prod_{\gamma \in \Gamma} L_\gamma$  with  $|\Gamma| = m$  is directly decomposable. Each conditionally  $\wedge$ - $m$ -complete tolerance on  $L$  is directly decomposable.*

**Proof.** Put

$$T_\gamma = \{(x_\gamma, y_\gamma) \mid x_\gamma = x(\gamma), y_\gamma = y(\gamma) \text{ for some } (x, y) \in T\},$$

where  $T$  is a conditionally  $\vee$ - $m$ -complete tolerance on  $L = \prod_{\gamma \in \Gamma} L_\gamma$  with  $|\Gamma| = m$ . Clearly  $T_\gamma$  is a compatible tolerance on  $L_\gamma$  for each  $\gamma \in \Gamma$  and

$$T \subseteq \prod_{\gamma \in \Gamma} T_\gamma.$$

We prove the converse inclusion. Let  $(x, y) \in \prod_{\gamma \in \Gamma} T_\gamma$ . With respect to the convexity of compatible tolerances it suffices to consider only the case  $x \leq y$ . Then  $(x(\gamma), y(\gamma)) \in T$  for each  $\gamma \in \Gamma$ , i.e. there exist elements  $a$  and  $b$  of  $L$  such that  $(f(x, a, \gamma), f(y, b, \gamma)) \in T$ . By Lemma, this implies

$$(f(f(x, a, \gamma), x, \gamma), f(f(y, b, \gamma), x, \gamma)) \in T.$$

Since  $f(f(x, a, \gamma), x, \gamma) = x$  and  $f(f(y, b, \gamma), x, \gamma) = f(y, x, \gamma)$ , we infer  $(x, f(y, x, \gamma)) \in T$ . As  $x \leq y$ , we conclude  $y = \bigvee_{\gamma \in \Gamma} f(y, x, \gamma)$ . Since  $T$  is conditionally  $\vee$ - $m$ -complete, we obtain  $(x, y) \in T$ , which was to be proved. Dually we can prove the assertion for conditionally  $\wedge$ - $m$ -complete tolerances.

**Corollary** (cf. [3]). *Each compatible tolerance on the lattice  $L = L_1 \times \dots \times L_n$  is directly decomposable.*

Now we shall turn our attention to *semilattices*. The operation on a semilattice will be denoted by  $\otimes$ .

Consider a semilattice  $S = \prod_{\gamma \in \Gamma} S_\gamma$ , where  $|\Gamma| = m \geq \aleph_0$ . A *conditionally m-complete tolerance* can be defined analogously to the above defined similar concepts for lattices.

**Theorem 2.** *Let  $S = \prod_{\gamma \in \Gamma} S_\gamma$ , where  $S_\gamma$  are semilattices with zero elements and  $|\Gamma| = m \geq \aleph_0$ . Then there exists a conditionally m-complete tolerance which is not directly decomposable.*

*Proof.* For each  $\gamma \in \Gamma$  let the zero element of the semilattice  $S_\gamma$  be denoted by  $z_\gamma$ . Let  $T$  be the tolerance on  $S$  defined so that  $(a, b) \in T$  if and only if either  $a = b$ , or there exists an infinite subset  $\Gamma(a, b)$  of  $\Gamma$  such that  $a(\gamma) = b(\gamma) = z_\gamma$  for each  $\gamma \in \Gamma(a, b)$ . We shall prove that  $T$  is a conditionally m-complete tolerance on  $S$ . Let  $(a_\delta, b_\delta) \in T$  for  $\delta \in \Delta$ , where  $|\Delta| = m$ . If  $a_\delta = b_\delta$  for each  $\delta \in \Delta$ , then  $\otimes_{\delta \in \Delta} a_\delta = \otimes_{\delta \in \Delta} b_\delta$  and  $(\otimes_{\delta \in \Delta} a_\delta, \otimes_{\delta \in \Delta} b_\delta) \in T$ . If there exists  $\varepsilon \in \Delta$  such that  $a_\varepsilon \neq b_\varepsilon$ , then there exists an infinite subset  $\Gamma(a_\varepsilon, b_\varepsilon)$  of  $\Gamma$  such that  $a_\varepsilon(\gamma) = b_\varepsilon(\gamma) = z_\gamma$  for each  $\gamma \in \Gamma(a_\varepsilon, b_\varepsilon)$ . Now  $\otimes_{\delta \in \Delta} a_\delta(\gamma) = a_\varepsilon(\gamma) \otimes_{\delta \in \Delta - \{\varepsilon\}} a_\delta(\gamma) = z_\gamma \otimes_{\delta \in \Delta - \{\varepsilon\}} a_\delta(\gamma) = z_\gamma$ ,  $\otimes_{\delta \in \Delta} b_\delta(\gamma) = b_\varepsilon(\gamma) \otimes_{\delta \in \Delta - \{\varepsilon\}} b_\delta(\gamma) = z_\gamma \otimes_{\delta \in \Delta - \{\varepsilon\}} b_\delta(\gamma) = z_\gamma$  for each  $\gamma \in \Gamma(a_\varepsilon, b_\varepsilon)$  and  $(\otimes_{\delta \in \Delta} a_\delta, \otimes_{\delta \in \Delta} b_\delta) \in T$ . Now define again  $T_\gamma = \{(x_\gamma, y_\gamma) \mid x_\gamma = x(\gamma), y_\gamma = y(\gamma) \text{ for some } (x, y) \in T\}$ . For each  $\gamma \in \Gamma$  let  $a_\gamma, b_\gamma$  be two arbitrary elements of  $S_\gamma$ . Let  $c_\gamma, d_\gamma$  be the elements of  $S$  such that  $c_\gamma(\gamma) = a_\gamma, d_\gamma(\gamma) = b_\gamma, c_\gamma(\delta) = d_\gamma(\delta) = z_\gamma$  for each  $\delta \in \Gamma - \{\gamma\}$ . Clearly  $(c_\gamma, d_\gamma) \in T$ , hence  $(a_\gamma, b_\gamma) \in T_\gamma$ . As  $a_\gamma, b_\gamma$  were chosen arbitrarily,  $T_\gamma$  is the universal binary relation on  $S_\gamma$  for each  $\gamma \in \Gamma$ . Therefore  $\prod_{\gamma \in \Gamma} T_\gamma$  is the universal binary relation on  $S$ , while  $T$  is not and  $T \neq \prod_{\gamma \in \Gamma} T_\gamma$ .

#### References

- [1] *Chajda, I.*: Congruence factorizations on distributive lattices. *Math. Slovaca* 28 (1978), 343–347.
- [2] *Chajda, I.*: Varieties with directly decomposable diagonal subalgebras. *Algebra Univ.* (to appear).
- [3] *Chajda, I. - Nieminen, J.*: Direct decomposability of tolerances on lattices, semilattices and quasilattices. *Czech. Math. J.* (to appear).
- [4] *Chajda, I. - Zelinka, B.*: Tolerance relations on direct products. *Glasnik Mat. (Zagreb)* 14 (1979), 11–16.
- [5] *Pócs, J.*: Congruence relations on direct products of lattices. *Math. Slovaca* (to appear).

*Authors' addresses:* I. Chajda, tř. Lidových milicí 22, 750 00 Přerov, ČSSR; B. Zelinka, Felberova 2, 460 01 Liberec 1, ČSSR (katedra matematiky VŠST).