

Alessandra Cherubini; Ada Varisco

Semigroups and rings whose proper one-sided ideals are power joined

*Czechoslovak Mathematical Journal*, Vol. 34 (1984), No. 1, 121–125

Persistent URL: <http://dml.cz/dmlcz/101931>

## Terms of use:

© Institute of Mathematics AS CR, 1984

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## SEMIGROUPS AND RINGS WHOSE PROPER ONE-SIDED IDEALS ARE POWER JOINED

A. CHERUBINI and A. VARISCO, Milano

(Received February 9, 1983)

The main result of the present paper consists in a characterization of semigroups whose proper one-sided ideals are power joined (Th. 1.6). A theorem of Pondělíček on uniform semigroups, contained in [3], is found again as a corollary of this result. Moreover, with regard to the fact that every one-sided ideal is a biideal (biideal of a semigroup  $S$  is a subsemigroup  $B$  of  $S$  such that  $BSB \subseteq B$ ), Th. 1.6 provides also an answer to the question put in the Mathematical Reviews (82g; 20097) by the reviewer of the note [3]. The second section of the work contains a characterization of rings whose proper one-sided ideals are multiplicatively power joined semigroups.

1. We start by proving some lemmas which will enable us to establish the main theorem. We remember that by a *power joined semigroup* we mean a semigroup  $S$  such that  $a^h = b^k$  for every  $a, b \in S$  and  $h, k$  positive integers (see [4], II.7.7).

**Lemma 1.1.** *A semigroup  $S$  whose proper left (right) ideals of the form  $Sb$  ( $bS$ ) ( $b \in S$ ) are power joined is a semilattice of archimedean semigroups.*

*Proof.* Let  $a, b \in S$  with  $a = xby$  ( $x, y \in S^1$ ). Then  $a^2 = (xbyx)by$  with  $z = xbyx \in S$ . If  $Sb = S$ , it follows that  $z = wb$  for some  $w \in S$ , and therefore  $a^2 = zby = wb^2y$ . If on the contrary,  $Sb \subset S$ , since  $Sb$  is power joined, there are two positive integers  $h, k$  such that  $(yzb)^h = b^{2k}$ , and it results that  $a^{2(h+1)} = (zby)^{h+1} = zb(yzb)^h y = zb^{2k+1}y$ . Thus in any case  $b^2$  divides a power of  $a$ , which suffices to conclude that  $S$  is a semilattice of archimedean semigroups (see [5], Th. 2.1).

**Lemma 1.2.** *A non archimedean semigroup  $S$  whose proper one-sided ideals are power joined is a semilattice of two semigroups  $M$  and  $S \setminus M$ , where  $M$  is power joined and coincides with the greatest ideal of  $S$ , and  $S \setminus M$  is a group. Moreover,  $S$  has an identity, which is the identity of  $S \setminus M$ .*

*Proof.* By Lemma 1.1,  $S$  is a semilattice of archimedean semigroups, so it is Putcha's  $\mathcal{Q}$ -semigroup (see [1], Definitions 1.1 and 1.4). Therefore it follows from Corollary 1.5 of [1] that  $S$  is a semilattice of two semigroups:  $M$  which is power

joined and the greatest ideal of  $S$ , and  $S \setminus M$ . It remains to prove that  $S \setminus M$  is a group, whose identity is the identity of  $S$ . In fact, if  $L$  is a proper left ideal of  $S \setminus M$ , it is immediate to verify that  $L \cup M$  is a proper left ideal of  $S$ , and therefore a power joined semigroup. But this is a contradiction, since power joined semigroups cannot be disjoint unions of proper subsemigroups. Hence  $S \setminus M$  is left simple. In the same way we find that  $S \setminus M$  is right simple. Thus  $S \setminus M$  is a group. Now, if  $u$  is the identity of  $S \setminus M$ ,  $Su \subset S$  implies that  $Su$  is power joined. Then, for every  $x \in M$ , there exists a positive integer  $m$  such that  $(xu)^m = u$ , but this implies  $u \in M$ , a contradiction. Thus  $Su = S$ , and analogously  $uS = S$ , which means that  $u$  is the identity of  $S$ .

**Lemma 1.3.** *A non simple archimedean semigroup whose proper one-sided ideals are power joined is power joined.*

Proof is immediate.

**Lemma 1.4.** *A simple semigroup  $S$  whose proper one-sided ideals are power-joined has at least an idempotent.*

Proof. The lemma is obvious if  $S$  is a group. Otherwise, we may suppose that  $S$  is not left simple. Therefore, there exists  $a \in S$  such that  $Sa \subset S$ . Moreover,  $S$  being simple, we have  $Sa^2S = S$ , hence  $a = xa^2y$  for some  $x, y \in S$ . Since  $Sa$  is power joined, there are two positive integers  $h, k$  such that  $(xa)^h = a^{2k}$ , whence  $a = (xa)ay = (xa)^h ay^h = a^{2k+1}y^h$ . Hence  $a^2 = a^{2k-1}a^2y^ha$  with  $y^ha \in Sa$ . Then there exist two positive integers  $m, n$  such that  $(y^ha)^m = a^{2n}$ , and consequently,  $a^2 = (a^{2k-1})^m a^2(y^ha)^m = a^{(2k-1)m+2+2n}$ . So  $S$  has an idempotent.

**Lemma 1.5.** *A simple semigroup  $S$  whose proper one-sided ideals are power-joined is either a group or a left (right) zero-semigroup of two periodic groups.*

Proof. Since a simple semigroup with a unique idempotent is a group, we may suppose that  $S$  contains two idempotents  $e, f$  with  $e \neq f$  (Lemma 1.4). First we remark that  $Se \subset S$  implies  $e = (fe)^h$  for some positive integer  $h$ , since  $Se$  is a power joined semigroup containing  $e$  and  $fe$ . Hence  $e = fe$ . Analogously  $fS \subset S$  implies  $f = fe$ . Since  $e \neq f$ , we have either  $Se = S$  or  $fS = S$ . In the same way we find that necessarily either  $eS = S$  or  $Sf = S$ .

If  $Se = eS = S$  or  $Sf = fS = S$ , the semigroup  $S$  has an identity. If, on the contrary,  $Se = Sf = S$  ( $eS = fS = S$ ),  $S$  has two right (left) identities. It is immediate to verify that in the first case  $S$  cannot have a third idempotent different from  $e$  and  $f$ . In the second case, every other idempotent is a right (left) identity. In both cases one of the idempotents of  $S$  is primitive (since their number is finite when  $S$  has an identity<sup>1)</sup>); since they are not comparable in the other case), so  $S$  is completely simple, i.e.  $S$  is a rectangular band of groups. This leads to a contradiction when  $S$

<sup>1)</sup>  $S$ , being simple with order greater than 1, cannot have a zero.

has an identity, so all idempotents of  $S$  are right (left) identities. Now, let  $G_\alpha$  and  $G_\beta$  be two maximal subgroups of  $S$  with  $G_\alpha \neq G_\beta$ . From the fact that the idempotents of  $S$  are right (left) identities it follows that  $S$  is a left (right) zero-semigroup of groups, whence  $(G_\alpha \cup G_\beta)S \subseteq G_\alpha \cup G_\beta$  ( $S(G_\alpha \cup G_\beta) \subseteq G_\alpha \cup G_\beta$ ). Then, if  $G_\alpha \cup G_\beta \subset S$ ,  $G_\alpha \cup G_\beta$  has to be power joined, a contradiction. Thus  $G_\alpha \cup G_\beta = S$  and  $S$  is a left (right) zero-semigroup of two groups  $G_\alpha, G_\beta$ . Finally,  $G_\alpha$  and  $G_\beta$  are periodic since they are proper right (left) ideals of  $S$  and consequently, power joined groups.

Now we are able to state the following result:

**Theorem 1.6.** *Let  $S$  be a semigroup whose proper one-sided ideals are power joined. Then  $S$  satisfies one of the following conditions:*

- i)  $S$  is power joined,
- ii)  $S$  is a group,
- iii)  $S$  is a left (right) zero-semigroup of two periodic groups,
- iv)  $S$  is a semilattice of two semigroups  $M$  and  $S \setminus M$ , where  $M$  is power joined and coincides with the greatest ideal of  $S$ , and  $S \setminus M$  is a group. Moreover, the identity of  $S \setminus M$  is the identity for  $S$ .

*Conversely, if  $S$  is a semigroup of type i), ii), iii) or iv), every proper one-sided ideal of  $S$  is power joined.*

**Proof.** The first part of the statement follows from Lemmas 1.2, 1.3 and 1.5. The converse is immediate.

We recall that a semigroup  $S$  is said to be *uniform* if every two left ideals of  $S$  and every two right ideals of  $S$  have a non-empty intersection (see [3], p. 331). According to this definition, we may find again Th. 1 of [3] as a corollary of the above Th. 1.6. In fact, when  $S$  is uniform, the case iii) of the statement of Th. 1.6 can not occur, since the components of a left (right) zero-semigroup of groups are disjoint right (left) ideals, and the converse is obvious.

**Addendum.** From the above Th. 1.6 and from Th. 4 of [7] the following Theorem can be immediately deduced: “ $S$  is a semigroup whose proper subsemigroups are power joined if and only if  $S$  is either power joined or a band of order two”. This result extends Th. 2 of [8], which has come to the authors’ knowledge when the manuscript was already sent to the Editor.

**2.** In this section we shall prove a theorem for rings analogous to Th. 1.6. In the sequel we shall denote by  $(R, \cdot)$  the multiplicative semigroup of a ring  $R$ .

**Theorem 2.1.** *Let  $R$  be a ring whose proper one-sided ideals are multiplicatively power joined semigroups. Then  $R$  satisfies one of the following conditions:*

- i)  $R$  is a nilring,

ii)  $R$  is a ring with identity and  $(R, \cdot)$  is a semilattice of two semigroups  $M$  and  $R \setminus M$ , where  $M$  is a nilring and coincides with the greatest ideal of  $R$ , and  $R \setminus M$  is a group.

Conversely, if  $R$  is a ring of type i) or ii), every proper one-sided ideal of  $R$  is a multiplicatively power joined semigroup.

*Proof.* Let  $R$  be a ring whose proper one-sided ideals are multiplicatively power joined semigroups. If  $(R, \cdot)$  is archimedean, it has to be a nilsemigroup, since it contains the zero, and in this case  $R$  is a nilring. Then, let us suppose that  $(R, \cdot)$  is not archimedean. We note that every left ideal of  $(R, \cdot)$  of the form  $Rb$  ( $b \in R$ ) is an additive subgroup of  $R$ , hence it is a left ideal of  $R$ . Analogously, every right ideal of  $(R, \cdot)$  of the form  $aR$  ( $a \in R$ ) is a right ideal of  $R$ . Therefore, by Lemma 1.1,  $(R, \cdot)$  is a semilattice of archimedean semigroups. Since  $(R, \cdot)$  is not archimedean, it contains a proper prime ideal  $M$ <sup>2)</sup> (see [1], Th. 1.3). For every  $a, b \in M$ , we have  $aR \subseteq M$  and  $Rb \subseteq M$ ; hence  $aR$  and  $Rb$  are proper one-sided ideals of  $R$ , so they are multiplicatively power joined semigroups. Since  $ab \in aR \cap Rb$ , there are three positive integers  $h, k, l$  such that  $a^{2h} = (ab)^k = b^{2l}$ . Thus  $M$  is power joined and, containing zero, it has to be a nilsemigroup. Then, let  $a \in M$ ,  $c \in R \setminus M$ . If  $Rc \subset R$ , since  $Rc$  and  $aR$  are power joined and  $ac \in aR \cap Rc$ , we may conclude as above that  $a^{2h} = c^{2l}$  for some positive integers  $h, l$ . This is a contradiction, since  $a$  and  $c$  are in disjoint semigroups. Thus, for every  $c \in R \setminus M$  we have  $Rc = R$ . Analogously we find  $cR = R$ . This implies that for every  $c, d \in R \setminus M$  there exist  $x, y \in R$  such that  $d = xc = cy$ . Since  $x \in M$  ( $y \in M$ ) implies  $d \in M$ , a contradiction, we have  $x, y \in R \setminus M$ , which enables us to conclude that  $R \setminus M$  is a group. If  $u$  is the identity of  $R \setminus M$ , the relations  $Ru = uR = R$  imply that  $u$  is the identity of  $R$ . Thus it remains to prove that  $M$  is the greatest ideal of  $R$ . In fact, since  $(R, \cdot)$  is a semilattice of archimedean semigroups, such that every element has a power in a subgroup, the set of all nilpotents of  $R$  is an ideal of  $R$  (see [6], Th. 8), which obviously coincides with  $M$ . The maximality of  $M$  is guaranteed by the fact that  $R \setminus M$  is a group.

The converse is immediate.

**Remark 2.2.** The class of rings of type ii) in the statement of Th. 2.1 contains all division rings; nevertheless it may be interesting to note that this class is wider than that of division rings. This is proved by the following example. Let  $R$  be the set of square matrices of the form

$$\begin{bmatrix} a & b \\ -b & a \end{bmatrix} + h \begin{bmatrix} 1 & i \\ i & -1 \end{bmatrix}$$

( $a, b$  real numbers;  $h$  complex;  $i = \sqrt{-1}$ ). It is a routine verification to prove that  $R$  is a ring with identity with respect to the usual sum and product of matrices. Moreover, it is immediate to show that the subset  $M$  of  $R$  containing the matrices

<sup>2)</sup> Here and in [1] "prime ideal" means, following Clifford, an ideal  $I$  of a semigroup  $S$  such that  $S \setminus I$  is a subsemigroup. Such ideals are called "completely prime" by Petrich.

$$h \begin{bmatrix} 1 & i \\ i & -1 \end{bmatrix}$$

is a nilring of order greater than 1 and it is the greatest ideal of  $R$ , while the subset  $R \setminus M$  is a multiplicative group. So  $R$  is a ring of type ii) and it is not a division ring.

#### References

- [1] *Cherubini, A.* and *A. Varisco*: On Putcha's  $Q$ -semigroups, *Semigroup Forum* 18 (1979) 313–317.
- [2] *Clifford, A. H.* and *G. B. Preston*: The algebraic theory of semigroups, Amer. Math. Soc., Vol. I (1961).
- [3] *Pondělíček, B.*: Uniform semigroups whose proper quasi-ideals are power joined, *Semigroup Forum* 22 (1981) 331–337.
- [4] *Petrich, M.*: Introduction to semigroups, Merrill, Columbus (1973).
- [5] *Putcha, M. S.*: Band of  $t$ -archimedean semigroups, *Semigroup Forum* 6 (1973) 232–239.
- [6] *Putcha, M. S.*: Rings which are semilattices of archimedean semigroups, *Semigroup Forum* 23 (1981) 1–5.
- [7] *Cherubini, A.* e *A. Varisco*: Sui semigrupperi i cui sottosemigrupperi propri sono  $t$ -archimedei, *Istituto Lombardo (Rend. Sc.)* 112 (1978) 91–98.
- [8] *Nagy, A.*: Semigroups whose proper two-sided ideals are power joined, *Semigroup Forum* 25 (1982) 325–329.

*Authors' address*: Dipartimento di Matematica, Politecnico di Milano, 20133, Milano, Italia.