

Ivan Chajda; Bohdan Zelinka

A characterization of tolerance-distributive tree semilattices

Czechoslovak Mathematical Journal, Vol. 37 (1987), No. 2, 175–180

Persistent URL: <http://dml.cz/dmlcz/102146>

Terms of use:

© Institute of Mathematics AS CR, 1987

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A CHARACTERIZATION OF TOLERANCE-DISTRIBUTIVE
TREE SEMILATTICES

IVAN CHAJDA, Přerov, and BOHDAN ZELINKA, Liberec

(Received May 28, 1982)

A tolerance on an algebra $\mathfrak{A} = \langle A, \mathcal{F} \rangle$ is a reflexive and symmetric binary relation T on A which has the Substitution Property with respect to \mathcal{F} , i.e., $(a_1, b_1) \in T, \dots, (a_n, b_n) \in T$ implies $(f(a_1, \dots, a_n), f(b_1, \dots, b_n)) \in T$ for each n -ary operation $f \in \mathcal{F}$ and any elements $a_1, \dots, a_n, b_1, \dots, b_n$ of A . The set of all tolerances on \mathfrak{A} forms an algebraic lattice $LT(\mathfrak{A})$ with respect to the set inclusion (see [4], [5]). Basic properties of this lattice were investigated in [4] and, especially for semilattices, in [5] and [6].

An algebra \mathfrak{A} is called *congruence-distributive*, if the congruence lattice $Con(\mathfrak{A})$ is distributive. It is well-known that lattices and semilattices are congruence-distributive. Although tolerances are a generalization of congruences, the situation with them is quite different. We shall call an algebra \mathfrak{A} *tolerance-distributive* (or *tolerance-modular*), if $LT(\mathfrak{A})$ is distributive (or modular, respectively). A class \mathcal{K} of algebras is tolerance-distributive (or tolerance-modular), if each $\mathfrak{A} \in \mathcal{K}$ has this property.

It was proved in [2] and [3] that the variety \mathcal{D} of all distributive lattices is the only non-trivial tolerance-distributive lattice variety. A variety of semilattices is tolerance-modular if and only if it is trivial, see [2]. The variety \mathcal{D} of distributive lattices is the only non-trivial tolerance-modular variety [1]. H.-J. Bandelt [1] has investigated a weaker condition: a lattice L with the least element O is O -modular, if it does not contain a minimal non-modular sublattice containing the least element O . He has proved that every lattice L is tolerance- O -modular (i.e., $LT(L)$ is O -modular).

Our first results for tolerance lattices of semilattices were presented in [5]:

- (i) The class of all semilattices is not tolerance- O -modular.
- (ii) The class of all tree semilattices is tolerance- O -modular.

Recall that a semilattice S is a tree semilattice, if each interval of S (in the induced ordering) is a chain. The result (ii) has motivated our effort to characterize tolerance-modular or tolerance-distributive semilattices among tree semilattices.

Let S be a semilattice. Its operation will be denoted by the symbol \vee and the induced ordering of S will be defined by: $x \leq y$ if and only if $x \vee y = y$.

Let us introduce the relation $C(S)$ of comparability on S . We define $C(S) = \{(x, y) \mid x \vee y \in \{x, y\}\}$. Clearly, $C(S)$ is a reflexive and symmetric binary relation on S . It plays the key role in our investigation of tolerance-distributivity of S . First we ask whether $C(S)$ is a tolerance, i.e., whether $C(S) \in LT(S)$ and whether at least the intersection of $T \in LT(S)$ with $C(S)$ is in $LT(S)$.

Theorem 1. *Let S be a semilattice. The following conditions are equivalent:*

- (1) S is a tree semilattice.
- (2) $T \cap C(S) \in LT(S)$ for each $T \in LT(S)$.

Proof. (1) \Rightarrow (2) Clearly, $T \cap C(S)$ is a reflexive and symmetric binary relation on S , since both T and $C(S)$ have these properties. It remains to prove the Substitution Property of $T \cap C(S)$. Let $(a, b) \in T \cap C(S)$, $(c, d) \in T \cap C(S)$. Then the Substitution Property of T implies $(a \vee c, b \vee d) \in T$. We only need to prove that also $(a \vee c, b \vee d) \in C(S)$. As $(a, b) \in C(S)$, $(c, d) \in C(S)$, we have four possibilities:

$$\begin{aligned} a &\leq b, & c &\leq d; \\ a &\geq b, & c &\geq d; \\ a &\leq b, & c &\geq d; \\ a &\geq b, & c &\leq d. \end{aligned}$$

The first two of them imply trivially the comparability of $a \vee c$ and $b \vee d$, and the fourth is analogous to the third. Without loss of generality it suffices to study the third case. Then

$$a \leq a \vee c \leq b \vee c \quad \text{and} \quad a \leq b \leq b \vee d \leq b \vee c,$$

thus both $a \vee c, b \vee d$ lie in the interval $[a, b \vee c]$. Since S is a tree semilattice, the interval is a chain and hence $a \vee c$ and $b \vee d$ are comparable, which proves (2).

(2) \Rightarrow (1). If S is not a tree semilattice, then it contains a subsemilattice $\{x, y, z, x \vee y\}$ with the diagram in Fig. 1. Then $(x, z) \in C(S)$, $(z, y) \in C(S)$. Let T be the least tolerance of $LT(S)$ containing the pairs (x, z) and (z, y) . Then $(x, z) \in T \cap C(S)$, $(z, y) \in T \cap C(S)$ and $(x \vee z, z \vee y) = (x, y) \in T$. But x and y are not comparable, i.e. $(x, y) \notin C(S)$ thus $T \cap C(S)$ has not the Substitution Property and $T \cap C(S) \notin LT(S)$.

Corollary 1. *If S is a tree semilattice, then $C(S) \in LT(S)$.*

Proof. The assertion follows directly from (2) of Theorem 1 by putting $T = S \times S$. First we shall study those tolerances T of $LT(S)$ for which $T \subseteq C(S)$.

Theorem 2. *Let S be a tree semilattice. Let $T_1 \in LT(S)$, $T_2 \in LT(S)$, $T_1 \subseteq C(S)$, $T_2 \subseteq C(S)$. Then $T_1 \vee T_2 = T_1 \cup T_2$.*

Proof. Let $T = T_1 \cup T_2$. Let $(a, b) \in T$, $(c, d) \in T$; then each of the pairs (a, b) , (c, d) belongs to T_1 or to T_2 . If $(a, b) \in T_1$, $(c, d) \in T_1$, then $(a \vee c, b \vee d) \in T_1 \subseteq T$. If $(a, b) \in T_2$, $(c, d) \in T_2$, then $(a \vee c, b \vee d) \in T_2 \subseteq T$. Suppose that $(a, b) \in T_1$,

$(c, d) \in T_2$. As $T_1 \in C(S)$, $T_2 \in C(S)$, the elements a, b are comparable and so are c, d . Let $a \leq b$, $c \leq d$. We have $a \leq a \vee c \leq b \vee d$, $a \leq b \leq b \vee d$. As S is a tree semilattice, the interval $[a, b \vee d]$ is a chain and thus $a \vee c$ and b are comparable. Analogously $c \leq a \vee c \leq b \vee d$, $c \leq d \leq b \vee d$, and $a \vee c$ and d are comparable as well. If $a \vee c \leq b$, $a \vee c \leq d$, then both b, d are in the interval $[a \vee c, b \vee d]$ and they are comparable. If $b \leq d$, then $b \vee d = d$ and $(a \vee c, b \vee d) = (a \vee c, d) = (c \vee a, d \vee a) \in T_2 \subseteq T$, because $(c, d) \in T_2$, $(a, a) \in T_2$. If $b \geq d$, then analogously $(a \vee c, b \vee d) \in T_1 \subseteq T$. If $a \vee c \geq b$, $a \vee c \geq d$, then $a \vee c \geq b \vee d$; as $a \leq b$, $c \leq d$, we have also $a \vee c \leq b \vee d$ and thus $a \vee c = b \vee d$. This implies $(a \vee c, b \vee d) \in \Delta \subseteq T$. If $b \leq a \vee c \leq d$, then $b \vee d = d$ and $(a \vee c, b \vee d) = (a \vee c, d) = (c \vee a, d \vee a) \in T_2 \subseteq T$. If $d \leq a \vee c \leq b$, then $b \vee d = b$ and $(a \vee c, b \vee d) = (a \vee c, b) = (a \vee c, b \vee c) \in T_1 \subseteq T$. Now let $a \leq b$, $c \geq d$. We have $a \leq a \vee c \leq b \vee c$, $a \leq b \leq b \vee d \leq b \vee c$, thus $a \vee c$ and $b \vee d$ are comparable. If $a \vee c \leq b \vee d$, then $b \leq b \vee d$ and $a \leq b$ imply $b \vee c = (b \vee a) \vee c = b \vee (a \vee c) \leq b \vee (b \vee d) = b \vee d$. As $c \geq d$ gives $b \vee c \geq b \vee d$, we have $b \vee c = b \vee d$, thus $(a \vee c, b \vee d) = (a \vee c, b \vee c) \in T_1 \subseteq T$. If $a \vee c \geq b \vee d$, then $b \leq a \vee c$, $c \leq a \vee c$ and $b \vee c \leq a \vee c$; since also $a \vee c \leq b \vee c$, we have $a \vee c = b \vee c$ and $(a \vee c, b \vee d) = (c \vee b, d \vee b) \in T_2 \subseteq T$. The cases $a \geq b$, $c \geq d$ and $a \geq b$, $c \leq d$ are analogous. We have proved that $T \in LT(S)$ and thus $T = T_1 \cup T_2 = T_1 \vee T_2$.

Theorem 3. *Let S be a tree semilattice. Then $C(S)$ is a distributive element in the lattice $LT(S)$, i.e.*

$$C(S) \wedge (T_1 \vee T_2) = (C(S) \wedge T_1) \vee (C(S) \wedge T_2)$$

for any $T_1 \in LT(S)$, $T_2 \in LT(S)$.

Proof. We have $C(S) \wedge T_1 \subseteq C(S)$, $C(S) \wedge T_2 \subseteq C(S)$ and thus, according to Theorem 2, $(C(S) \wedge T_1) \vee (C(S) \wedge T_2) = (C(S) \cap T_1) \cup (C(S) \cap T_2) = C(S) \cap (T_1 \cup T_2)$. On the other hand, suppose that there exist elements x, y of S such that $(x, y) \in C(S) \wedge (T_1 \vee T_2)$, but $(x, y) \notin C(S) \cap (T_1 \cup T_2)$. Then (as the operation \vee on S is commutative and associative) there exist elements x_1, x_2, y_1, y_2 of S such that $x = x_1 \vee x_2$, $y = y_1 \vee y_2$, $(x_1, y_1) \in T_1$, $(x_2, y_2) \in T_2$. As $(x, y) \in C(S)$, without loss of generality we may suppose that $x \geq y$. Consider the elements $x_1 \vee y$, $x_2 \vee y$. As they are both greater than or equal to y , they are comparable. As $(x_1 \vee y) \vee (x_2 \vee y) = x_1 \vee x_2 \vee y = x \vee y = x$, we have either $x_1 \vee y = x$, or $x_2 \vee y = x$. Suppose that $x_1 \vee y = x$. Then $(x_1, y_1) \in T_1$, $(y, y) \in T_1$ imply $(x_1 \vee y, y_1 \vee y) = (x, y) \in T_1$. Analogously $x_2 \vee y = x$ implies $(x, y) \in T_2$. In both cases we have a contradiction with the assumption that $(x, y) \in C(S) \cap (T_1 \cup T_2)$. Hence $C(S) \wedge (T_1 \vee T_2) = C(S) \cap (T_1 \cup T_2) = (C(S) \wedge T_1) \vee (C(S) \wedge T_2)$, which was to be proved.

Theorem 4. *Let S be a tree semilattice. The tolerances from $LT(S)$ which are*

contained in $C(S)$ form a sublattice $L_0(S)$ of $LT(S)$. The mapping $\varphi: T \mapsto T \cap C(S)$ is a homomorphism of $LT(S)$ onto $L_0(S)$.

Proof. We have $C(S) \in LT(S)$ according to Corollary 1. The lattice $L_0(S)$ is the ideal of $LT(S)$ with the greatest element $C(S)$. If T_1, T_2 are in $LT(S)$, then obviously $\varphi(T_1) \wedge \varphi(T_2) = (T_1 \cap C(S)) \cap (T_2 \cap C(S)) = (T_1 \cap T_2) \cap C(S) = \varphi(T_1 \wedge T_2)$. Now consider $T_1 \vee T_2$. According to Theorems 2 and 3 we have $\varphi(T_1) \vee \varphi(T_2) = \varphi(T_1 \cup \varphi(T_2)) = (T_1 \cap C(S)) \cup (T_2 \cap C(S)) = (T_1 \cup T_2) \cap C(S) = (T_1 \wedge T_2) \vee \vee C(S) = \varphi(T_1 \vee T_2)$. Thus $\varphi(T_1 \vee T_2) = \varphi(T_1) \vee \varphi(T_2)$, and φ is a homomorphism. Finally, for each $T \in L_0(S)$ we have $\varphi(T) = T$, thus φ is a mapping of $LT(S)$ onto $L_0(S)$.

Corollary 2. For a tree semilattice S the lattice $L_0(S)$ is a sublattice of the lattice of all subsets of $S \times S$ and hence it is distributive.

Let S be a tree semilattice, let $T_0 \in L_0(S)$. By $L^*(T_0)$ we denote the set of all tolerances $T \in LT(S)$ such that $T \cap C(S) = T_0$.

Theorem 5. Let S be a tree semilattice, let $T_0 \in L_0(S)$. The set $L^*(T_0)$ is a sublattice of $LT(S)$; its least element is T_0 , its greatest element is $\tilde{T}_0 = \{(x, y) \mid (x, x \vee y) \in T_0 \ \& \ (y, x \vee y) \in T_0\}$.

Proof. Let $T_1 \in L^*(T_0), T_2 \in L^*(T_0)$. Consider the homomorphism φ from Theorem 3. We have $\varphi(T_1 \vee T_2) = \varphi(T_1) \vee \varphi(T_2) = T_0 \vee T_0 = T_0$, $\varphi(T_1 \wedge T_2) = \varphi(T_1) \wedge \varphi(T_2) = T_0 \wedge T_0 = T_0$ and thus $T_1 \vee T_2 \in L^*(T_0), T_1 \wedge T_2 \in L^*(T_0)$ and $L^*(T_0)$ is a sublattice of $LT(S)$. Obviously $T_0 \in L^*(T_0)$ and $T_0 \subseteq T$ for each $T \in L^*(T_0)$, hence T_0 is the least element of $L^*(T_0)$. Now consider \tilde{T}_0 . Let $(a, b) \in \tilde{T}_0 \cap C(S)$. Then a, b are comparable; without loss of generality let $a \leq b$. As $(a, b) \in \tilde{T}_0$, we have $(a, b) = (a, a \vee b) \in T_0$; thus $\tilde{T}_0 \cap C(S) = T_0$ and $\tilde{T}_0 \in L^*(T_0)$. Let $T \in L^*(T_0)$ and let $(c, d) \in T$. As T has the Substitution Property, we have $(c, c \vee d) \in T, (d, c \vee d) \in T$. As $c \leq c \vee d, d \leq c \vee d$, we have $(c, c \vee d) \in C(S), (d, c \vee d) \in C(S)$, hence $(c, c \vee d) \in T \cap C(S)$, and so $(d, c \vee d) \in T_0$. As T and (c, d) were chosen arbitrarily, we have $T \subseteq \tilde{T}_0$ for each $T \in L^*(T_0)$ and \tilde{T}_0 is the greatest element of $L^*(T_0)$.

The tolerance $T \in LT(S)$ for which $T \subseteq C(S)$ is easily recognized.

Theorem 6. Let S be a tree semilattice, let T be a tolerance on S . Then the following two assertions are equivalent:

- (i) $T \in L_0(S)$.
- (ii) $T \subseteq C(S)$ and, if a, b, x are elements of S such that $a \leq x \leq b$ and $(a, b) \in T$, then $(x, b) \in T$.

Proof. (i) \Rightarrow (ii). Suppose that (i) holds. Obviously $T \in C(S)$. As $(a, b) \in T, (x, x) \in T$ and $T \in L_0(S) \subseteq LT(S)$, we have $(a \vee x, b \vee x) = (x, b) \in T$.

(ii) \Rightarrow (i). Suppose that (ii) holds. Let $(a, b) \in T, (c, d) \in T$. If b and d are incomparable, then $a \vee c = b \vee d$, because S is a tree semilattice, and thus $(a \vee c,$

$b \vee d) \in \Delta \subseteq T$. If $b \leq d$, then $b \vee d = d$ and $c \leq a \vee c \leq b \vee d = d$. But then $(c, d) \in T$ implies $(a \vee c, b \vee d) = (a \vee c, d) \in T$ according to (ii). The case $d \leq b$ is analogous. We have proved that $T \in LT(S)$. As $T \subseteq C(S)$, we have $T \in L_0(S)$.

Now we are ready to prove our main theorem characterizing tolerance-distributive and tolerance-modular tree semilattices.

Theorem 7. *Let S be a tree semilattice. Then the following three assertions are equivalent:*

(i) *S is a chain or S contains a maximal chain S_0 and an element $z \in S_0$ such that each element of $S - S_0$ is covered by z .*

(ii) *$LT(S)$ is distributive.*

(iii) *$LT(S)$ is modular.*

Proof. (i) \Rightarrow (ii). If S is a chain, then $LT(S) = L_0(S)$ and (ii) holds according to Corollary 2. Suppose that there exists a maximal chain S_0 in S and an element $z \in S_0$ such that each element of $S - S_0$ is covered by z . Let $T_1 \in LT(S)$, $T_2 \in LT(S)$ and suppose that $T_1 \vee T_2 \neq T_1 \cup T_2$. Let $(a, b) \in T_1 \vee T_2 - T_1 \cup T_2$. If a and b are comparable, then $(a, b) \in \varphi(T_1 \vee T_2) = \varphi(T_1) \cup \varphi(T_2) \subseteq T_1 \cup T_2$ according to Theorem 3, which is a contradiction. Thus suppose that a, b are incomparable. All elements of S which are greater than or equal to z are comparable with all elements of S ; therefore $a < z$, $b < z$ and at least one of the elements a, b belongs to $S - S_0$. Without loss of generality let $a \in S - S_0$; then a is a minimal element of S and all elements less than b (if any) form a chain. There exist pairs $(c, d) \in T_1$, $(e, f) \in T_2$ such that $c \vee e = a$, $d \vee f = b$. As a is a minimal element of S , we have $c = e = a$. As all elements less than b form a chain, at least one of the elements d, f is equal to b . Hence at least one of the pairs (c, d) , (e, f) is equal to (a, b) , and $(a, b) \in T_1 \cup T_2$, which is a contradiction. As T_1, T_2 , (a, b) were chosen arbitrarily, we have proved that $T_1 \vee T_2 = T_1 \cup T_2$ for any two tolerances T_1, T_2 from $LT(S)$. Since also $T_1 \wedge T_2 = T_1 \cap T_2$, the lattice $LT(S)$ is a sublattice of the lattice of all subsets of $S \times S$, and it is distributive.

(ii) \Rightarrow (iii). This is obvious.

(iii) \Rightarrow (ii). Suppose that (i) does not hold. Let S_0 be a maximal chain in S . Suppose that there exist elements x, x', y, y' such that $x \in S_0$, $y \in S_0$, $x \neq y$, $x' \in S - S_0$, $y' \in S - S_0$, x is the least element of S_0 greater than x' and y is the least element of S_0 greater than y' . As S_0 is a maximal chain in S , the element y is not a minimal element of S and there exists an element $y'' \in S_0$ such that $y'' < y$. Let $T_1 = C(S) \cup \{(x', y'), (y', x')\}$, $T_2 = C(S) \cup \{(x', y''), (y'', x')\}$, $T_3 = C(S) \cup \{(x', y''), (y'', x'), (x', y), (y, x')\}$, $T_4 = C(S) \cup \{(x', y), (y, x'), (x', y'), (y', x'), (x', y''), (y'', x')\}$. Each of these tolerances is in $LT(S)$ (the proof is left to the reader) and together with $C(S)$ they form a sublattice of $LT(S)$ whose diagram is in Fig. 2. Hence $LT(S)$ is not modular. We have proved that if $LT(S)$ is modular, then at most one element of the maximal chain S_0 of S may have the property that it is the least

element of S_0 greater than an element of $S - S_0$. If this is fulfilled and (i) does not hold, then there are elements x, x', y, y' of S such that $x < x' < z, y < y' < z$, and each of the elements x, x' is incomparable with each of the elements y, y' . Let $T_1 = C(S) \cup \{(x, y'), (y', x)\}$, $T_2 = C(S) \cup \{(x', y), (y, x')\}$, $T_3 = C(S) \cup \{(x', y), (y, x'), (x', y'), (y', x')\}$, $T_4 = C(S) \cup \{(x, y'), (y', x), (x', y), (y, x'), (x', y'), (y', x')\}$. These tolerances are in $LT(S)$ and together with $C(S)$ they form a sublattice of $LT(S)$ whose diagram is in Fig. 2. Hence $LT(S)$ is not modular. The only possibility for S to be modular is to fulfill (i).




Fig. 1




Fig. 2

Remark. In the terminology of [5] we may say that a tree semilattice S is tolerance-modular and tolerance-distributive if and only if it is either a chain, or a star semilattice, or the union of a chain and a star semilattice.

References

- [1] Bandelt, H.-J.: Tolerance relations on lattices. Bull. Austral. Math. Soc. 23 (1981), 367—381.
- [2] Chajda, I.: Distributivity and modularity of lattice of tolerance relations. Algebra Univ. 12 (1981), 247—255.
- [3] Chajda, I.: A characterization of distributive lattices by tolerance relations. Arch. Math. Brno 15 (1979), 203—204.
- [4] Chajda, I. - Zelinka, B.: Lattices of tolerances. Časop. přest. mat. 102 (1977), 10—24.
- [5] Chajda, I. - Zelinka, B.: Tolerance O -modular semilattices. Glasnik Mat. Zagreb (to appear).
- [6] Zelinka, B.: Tolerance relations on semilattices. Comment. Math. Univ. Carol. 16 (1975), 333—338.

Authors' addresses: I. Chajda, 750 00 Přerov, tř. Lidových milicí 22, Czechoslovakia; B. Zelinka, 460 01 Liberec 1, Studentská 1292, Czechoslovakia (katedra tváření a plastů VŠST).