

Alejandro Balbás de la Corte; Pedro Jiménez Guerra
Representation of operators by bilinear integrals

Czechoslovak Mathematical Journal, Vol. 37 (1987), No. 4, 551–558

Persistent URL: <http://dml.cz/dmlcz/102183>

Terms of use:

© Institute of Mathematics AS CR, 1987

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

REPRESENTATION OF OPERATORS BY BILINEAR INTEGRALS

A. BALBAS and P. JIMENEZ GUERRA, Madrid

(Received May 6, 1985)

Let us consider a locally compact Hausdorff topological space T and two complete locally convex Hausdorff spaces X and Z . Denote by \mathcal{B} the family of all non empty bounded closed balanced and convex subsets of X , and by X_B the linear subspace of X generated by B ($B \in \mathcal{B}$) equipped with the Minkowski functional q_B of B . The problem to be solved here is the following: Let $\mathcal{C}_B = \mathcal{C}_B(T, X_B)$ ($B \in \mathcal{B}$) be the space of all continuous functions tending to zero at infinity $f: T \rightarrow X_B$ endowed with the usual supremum norm

$$(1) \quad \|f\|_B = \sup \{q_B[f(t)]: t \in T\},$$

$\mathcal{C} = \bigcup \{\mathcal{C}_B: B \in \mathcal{B}\}$ and $\mathcal{F}: \mathcal{C} \rightarrow Z$ a linear operator with continuous restrictions $\mathcal{F}_B = \mathcal{F} | \mathcal{C}_B$. The main object of this paper is to represent \mathcal{F} by a bilinear integral. To this end we will consider the space Y of the linear mappings from X into Z with continuous restrictions to X_B , for all $B \in \mathcal{B}$, and the evaluation from $X \times Y$ into Z will be represented by xy ($x \in X, y \in Y$).

If \mathcal{R} is a generating family of seminorms on Z , for every $r \in \mathcal{R}, B \in \mathcal{B}$ and $y \in Y$, let us set

$$(2) \quad q_{B,r}(y) = \sup \{r(xy): x \in B\}.$$

It is easily proved that $\{q_{B,r}: B \in \mathcal{B}, r \in \mathcal{R}\}$ is a saturated family of seminorms defining on Y a topology (which henceforth will be the topology supposed to be defined on Y), making the evaluation mapping $X \times Y \rightarrow Z$ hypocontinuous.

Let Σ be the Borel σ -algebra of T and $\mu: \Sigma \rightarrow Y$ a countable additive measure. We define the semivariation $\|\mu\|_{B,r}$ and the variation $|\mu|_{B,r}$ ($B \in \mathcal{B}, r \in \mathcal{R}$) in the usual way:

$$(3) \quad \|\mu\|_{B,r}(E) = \sup_{i \in \mathcal{I}} r\left(\sum x_i \mu(E_i)\right) \quad (E \in \Sigma),$$

where the supremum is taken over all finite partitions $\{E_i\}_{i \in \mathcal{I}} \subset \Sigma$ of E and all finite families $\{x_i\}_{i \in \mathcal{I}} \subset B$, and

$$(4) \quad |\mu|_{B,r}(E) = \sup_{C \in \pi} \sum q_{B,r}[\mu(C)] \quad (E \in \Sigma),$$

where the supremum is taken over all finite partitions $\pi \subset \Sigma$ of E .

A set $A \in \Sigma$ is said to be a *null set* if $\|\mu\|_{B,r}(A) = 0$ for all $B \in \mathcal{B}$ and $r \in \mathcal{R}$.

We will denote by $\mathcal{S} = \mathcal{S}(T, X)$ and $\mathcal{S}_B = \mathcal{S}(T, X_B)$ ($B \in \mathcal{B}$) the spaces of simple functions from T into X and X_B , respectively, and by $\mathcal{C} + \mathcal{S}$ or $\mathcal{C}_B + \mathcal{S}_B$ the algebraic sum of \mathcal{C} and \mathcal{S} or \mathcal{C}_B and \mathcal{S}_B , respectively.

Definition 1. Let $\{v_{B,r}: B \in \mathcal{B}, r \in \mathcal{R}\}$ be a family of positive and finite measures defined on Σ , and $\mu: \Sigma \rightarrow Y$ a countable additive measure. We say that μ is $(v_{B,r})$ -continuous if

$$(5) \quad \lim_{v_{B,r}(E) \rightarrow 0} \|\mu\|_{B,r}(E) = 0.$$

In the case of μ of bounded variation, it is easily proved that μ is $\{\|\mu\|_{B,r}: B \in \mathcal{B}, r \in \mathcal{R}\}$ -continuous.

Henceforth we will suppose to be given a fixed family $\{v_{B,r}: B \in \mathcal{B}, r \in \mathcal{R}\}$ of positive and finite measures defined on Σ .

For the spaces X, Y and Z , and the evaluation mapping, the bilinear integral used here (with analogous properties as the bilinear integral given by Sivasankara in [14]) can be defined in the following way ([14]): Let $\mu: \Sigma \rightarrow Y$ be a $(v_{B,r})$ -continuous measure.

A sequence of functions $f_n: T \rightarrow X$ is said to be B -convergent ($B \in \mathcal{B}$) to $f: T \rightarrow X$ if

$$\bigcup_{n=1}^{\infty} f_n(T) \cup f(T) \subset X_B$$

and $q_B(f_n - f) \rightarrow 0$ a.e..

A function $f: T \rightarrow X$ is said to be B -measurable ($B \in \mathcal{B}$) if $f(T) \subset X_B$ and there exists a sequence of simple functions (simple functions are defined as usual) which is B -convergent to f , and a function $g: T \rightarrow X$ is said to be measurable if it is B -measurable for some $B \in \mathcal{B}$.

We will say that a function $f: T \rightarrow X$ is B -integrable ($B \in \mathcal{B}$) if $f(T) \subset X_B$ and there exists a sequence (f_n) of simple functions which is B -convergent to f and for every $\varepsilon > 0$ and $r \in \mathcal{R}$ there exists $\delta = \delta(\varepsilon, r) > 0$ such that

$$r\left(\int_A f_n d\mu\right) < \varepsilon$$

holds for all $n \in \mathbb{N}$ and every $A \in \Sigma$ with $\|\mu\|_{B,r}(A) < \delta$ (the integral of a simple function is defined as usual). A sequence (f_n) of the above type is called an approximating sequence of f .

A function $f: T \rightarrow X$ is said to be integrable if it is B -integrable for some $B \in \mathcal{B}$. It can be proved ([14]) that if $f: T \rightarrow X$ is integrable then the limit

$$\int_A f d\mu = \lim_n \int_A f_n d\mu$$

exists for every $A \in \Sigma$ and every approximating sequence (f_n) of f , and it is independent of the choice of the approximating sequence of f .

Definition 2. A linear operator $\mathcal{F}: \mathcal{C} \rightarrow Z$ is said to be $(v_{B,r})$ -continuous if for every $B \in \mathcal{B}, r \in \mathcal{R}$ and $\varepsilon > 0$ there exists $\delta > 0$ such that for all $E \in \Sigma$ with $v_{B,r}(E) < \delta$, $r[\mathcal{F}(f)] < \varepsilon$ holds for all $f \in \mathcal{C}_B$ with $f(T) \subset B$ and $f|_{T-E} \equiv 0$.

Analogous definitions can be given for linear Z -valued operators defined on \mathcal{S} or $\mathcal{C} + \mathcal{S}$.

Proposition 3. *Let μ be a $(v_{B,r})$ -continuous measure, then all functions belonging to \mathcal{C} are μ -integrable and the linear functional $\mathcal{F}: \mathcal{C} \rightarrow Z$ defined by*

$$\mathcal{F}(f) = \int_T f \, d\mu$$

is $(v_{B,r})$ -continuous and its restrictions \mathcal{F}_B are continuous.

Proof. Let $f \in \mathcal{C}$, then there exists $B \in \mathcal{B}$ with $f \in \mathcal{C}_B$. Let us prove that f is (μ, B) -integrable (and so, integrable).

$f: T \rightarrow X_B$ has a continuous extension $\bar{f}: \bar{T} \rightarrow X_B$ (\bar{T} being Alexandroff's compactification of T) given by $\bar{f}(\infty) = 0$, and then $\bar{f}(T)$ is compact and therefore there exist $t_1, \dots, t_n \in \bar{T}$ such that

$$f(T) \subset \bar{f}(T) \subset \bigcup_1^n B(\bar{f}(t_k), \varepsilon),$$

where $B(\bar{f}(t_k), \varepsilon)$ is the closed ball with center $\bar{f}(t_k)$ and radius ε .

Consider $A_1 = B(\bar{f}(t_1), \varepsilon)$, $A_2 = B(\bar{f}(t_2), \varepsilon) - A_1, \dots, A_n = B(\bar{f}(t_n), \varepsilon) - \bigcup_{k=1}^{n-1} A_k$, $E_k = f^{-1}(A_k)$ and

$$g_\varepsilon = \sum_{k=1}^n x_k \chi_{E_k}$$

with $x_k \in A_k$. Obviously, $f(T) \subset \bigcup_{k=1}^n A_k$, and $T = \bigcup_{k=1}^n E_k$, so if $z \in T$ then there exists $k \in \{1, \dots, n\}$ such that $t \in E_k$ and $f(t) \in A_k$. Then we have $q_B(x_k - f(t)) \leq \varepsilon$ or $q_B(g_\varepsilon(t) - f(t)) \leq \varepsilon$. By taking $\varepsilon = 1/n$, for $n \in \mathbb{N}$, we obtain that f is the uniform limit of simple functions, where from it is easily deduced that f is (μ, B) -integrable.

Moreover, for $B \in \mathcal{B}$, $r \in \mathcal{R}$ and $f \in \mathcal{C}_B$ we have

$$r(\mathcal{F}_B(f)) = r(\int_T f \, d\mu) \leq \|f\|_B \|\mu\|_{B,r}(T),$$

and therefore, \mathcal{F}_B is continuous.

Finally, \mathcal{F} is $(v_{B,r})$ -continuous because for every $B \in \mathcal{B}$, $r \in \mathcal{R}$, $\varepsilon > 0$, $E \in \Sigma$ and $f \in \mathcal{C}_B$ with $f(T) \subset B$ and $f|_{T-E} \equiv 0$ we have

$$r(\int_T f \, d\mu) = r(\int_E f \, d\mu) \leq \|f\|_B \|\mu\|_{B,r}(E) \leq \|\mu\|_{B,r}(E),$$

and so, if $\delta > 0$ is such that $v_{B,r}(E) < \delta$ implies $\|\mu\|_{B,r}(E) < \varepsilon$, then

$$r(\mathcal{F}(f)) \leq \|\mu\|_{B,r}(E) < \varepsilon.$$

Proposition 4. *A linear operator $\mathcal{F}: \mathcal{C} \rightarrow Z$ with continuous restrictions \mathcal{F}_B is $(v_{B,r})$ -continuous if and only if for every $B \in \mathcal{B}$, $r \in \mathcal{R}$ and $\varepsilon > 0$ there exist $\delta > 0$, $1 \geq \delta' > 0$ such that for all $E \in \Sigma$ with $v_{B,r}(E) < \delta$, $r(\mathcal{F}(f)) < \varepsilon$ holds for all $f \in \mathcal{C}_B$ with $q_B(f(t)) \leq \delta'$ for all $t \in T - E$ ¹⁾.*

Proof. Let us suppose that \mathcal{F} is $(v_{B,r})$ -continuous, then for every $B \in \mathcal{B}$, $r \in \mathcal{R}$

¹⁾ The same result can be proved for Z -valued operators defined on \mathcal{S} .

and $\varepsilon > 0$ there exist $\delta > 0$ and $1 \geq \delta' > 0$ such that for all $E \in \Sigma$ with $v_{B,r}(E) < \delta$, $r(\mathcal{F}(f)) < \varepsilon/2$ holds for all $f \in \mathcal{C}_B$ with $f|_{T-E} \equiv 0$ or $\|f\|_B \leq \delta'$. Now let $f \in \mathcal{C}_B$ with $q_B(f(t)) \leq \delta'$ for all $t \in T-E$, then there exist $g, h \in \mathcal{C}_B$ such that $\|g\|_B \leq \delta'$, $h|_{T-E} \equiv 0$ and $f = g + h$ and therefore,

$$r(\mathcal{F}(f)) < \varepsilon.$$

Notice that if $U = \{t \in T: q_B(f(t)) \leq \delta'\}$ then we may set

$$g(t) = \begin{cases} f(t) & \text{if } t \in U \\ \frac{\delta' f(t)}{q_B(f(t))} & \text{if } t \notin U \end{cases}$$

and $h = f - g$.

Proposition 5. *Let μ and \mathcal{F} be as in Proposition 3, then there is an extension $\mathcal{F}^s: \mathcal{C} + \mathcal{S} \rightarrow Z$ of \mathcal{F} such that \mathcal{F}^s is $(v_{B,r})$ -continuous and its restrictions $\mathcal{F}_B^s: \mathcal{C}_B + \mathcal{S}_B \rightarrow Z$ are continuous for all $B \in \mathcal{B}$ (the topologies of \mathcal{C} and \mathcal{S}_B are defined by the norm (1)).*

Proof. Let us define

$$\mathcal{F}^s(f) = \int_T f \, d\mu$$

for $f \in \mathcal{C} + \mathcal{S}$, then $\mathcal{F}_B^s (B \in \mathcal{B})$ is continuous because

$$r(\int_T f \, d\mu) \leq \|f\|_B \|\mu\|_{B,r}(T)$$

for $r \in \mathcal{R}$ and $f \in \mathcal{C}_B + \mathcal{S}_B$. To prove that \mathcal{F}^s is $(v_{B,r})$ -continuous it is enough to proceed as in Proposition 3.

Theorem 6. *Let $\mathcal{F}: \mathcal{C} \rightarrow Z$ be a linear operator with continuous restrictions \mathcal{F}_B for all $B \in \mathcal{B}$. Then the following assertions are equivalent:*

6.1. *There exists a $(v_{B,r})$ -continuous countable additive measure $\mu: \Sigma \rightarrow Y$ such that*

$$\mathcal{F}(f) = \int_T f \, d\mu$$

for all $f \in \mathcal{C}$.

6.2. *There exists a $(v_{B,r})$ -continuous operator $\mathcal{F}^s: \mathcal{C} + \mathcal{S} \rightarrow Z$ with continuous restrictions $\mathcal{F}_B^s (B \in \mathcal{B})$, which extends \mathcal{F} .*

6.3. *There exists a linear $(v_{B,r})$ -continuous operator $\mathcal{G}: \mathcal{S} \rightarrow Z$ with continuous restrictions $\mathcal{G}_B (B \in \mathcal{B})$, such that for every B*

$$(6) \quad \lim_n \mathcal{G}_B(f_n) = \mathcal{F}_B(f)$$

holds for every sequence $(f_n)_n \subset \mathcal{S}_B$ which is uniformly convergent to $f \in \mathcal{C}_B$.

Proof. From Propositions 3 and 5 it is immediately deduced that 6.1 implies 6.2. Moreover, 6.2 clearly implies 6.3. Let us prove that 6.3 implies 6.1. This will be done in four steps:

i) *Construction of μ .* Let $E \in \Sigma$ and define

$$\mu(E)(x) = \mathcal{G}(x\chi_E)$$

for $x \in X$. Then $\mu(E)$ is a linear operator from X into Z such that if $B \in \mathcal{B}$, $x \in X_B$ and the sequence $(x_n)_{n \in \mathbb{N}} \subset X_B$ converges to x , then the sequence $(x_n\chi_E)$ is uniformly convergent to $x\chi_E$ and therefore

$$\mu(E)(x) = \mathcal{G}(x\chi_E) = \lim_n \mathcal{G}(x_n\chi_E) = \lim_n \mu(E)(x_n)$$

and $\mu(E) \in Y$.

ii) μ *is countably additive.* The finite additivity of μ results trivially from the linearity of \mathcal{G} . Let now $(E_n) \subset \Sigma$ be a disjoint sequence, then

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) - \sum_{i=1}^n \mu(E_i) = \mu\left(\bigcup_{i>n} E_i\right)$$

holds for every $n \in \mathbb{N}$, and given $r \in \mathcal{R}$, $B \in \mathcal{B}$ and $\varepsilon > 0$ it follows from the $(v_{B,r})$ -continuity of \mathcal{G} that there exists $n_0 \in \mathbb{N}$ such that

$$r\left[\mathcal{G}\left(x\chi_{\bigcup_{i \geq n_0} E_i}\right)\right] < \varepsilon$$

for all $x \in B$, so

$$q_{B,r}\left[\mu\left(\bigcup_{i \geq n_0} E_i\right)\right] < \varepsilon$$

and therefore

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} \mu(E_i).$$

iii) μ *is $(v_{B,r})$ -continuous.* Let $r \in \mathcal{R}$, $B \in \mathcal{B}$ and $\varepsilon > 0$. Since \mathcal{G} is $(v_{B,r})$ -continuous there exists $\delta > 0$ such that

$$r[\mathcal{G}_B(f)] < \varepsilon$$

for all $f \in \mathcal{S}_B$ with $f|_{T-E} \equiv 0$ for some $E \in \Sigma$ of measure $v_{B,r}(E) < \delta$. Therefore, if $E \in \Sigma$ and $v_{B,r}(E) < \delta$ then for every finite family $\{x_1, \dots, x_n\} \subset B$ and every finite partition $\{E_1, \dots, E_n\} \subset \Sigma$ of E , we have

$$r\left[\sum_{i=1}^n x_i \mu(E_i)\right] = r\left[\mathcal{G}\left(\sum_{i=1}^n x_i \chi_{E_i}\right)\right] \leq \varepsilon$$

and consequently,

$$\|\mu\|_{B,r}(E) \leq \varepsilon.$$

iv) μ *represents \mathcal{F} .* Let $B \in \mathcal{B}$ and $f \in \mathcal{C}_B$. As in Proposition 3 we can find a sequence $(f_n) \subset \mathcal{S}_B$ which is uniformly convergent to f , and therefore,

$$\mathcal{F}(f) = \lim_n \mathcal{G}_B(f_n) = \lim_n \int_T f_n d\mu = \int_T f d\mu.$$

Theorem 7. *If $\{v_{B,r}: B \in \mathcal{B}, r \in \mathcal{R}\}$ are Radon measures (i.e. regular Borel measures) and \mathcal{F} is as in Theorem 6 verifying 6.1, then the measure μ of 6.1 is unique.*

Proof. Suppose that there exist two $(v_{B,r})$ -continuous measures $\mu, \mu': \Sigma \rightarrow Y$ such that

$$\mathcal{F}(f) = \int_T f d\mu = \int_T f d\mu'$$

holds for all $f \in \mathcal{C}$. Then Proposition 5 implies the existence of two $(v_{B,r})$ -continuous extensions \mathcal{F}^s and $\mathcal{F}^{s'}$ of \mathcal{F} to $\mathcal{C} + \mathcal{S}$. If $x \in X$ and $E \in \Sigma$, let us consider $B \in \mathcal{B}$ with $x \in X_B$ and $r \in \mathcal{R}$ arbitrary. Then two sequences (K_n) and (G_n) of compact and open subsets of T , respectively, can be found such that $K_n \subset E \subset G_n$ and

$$v_{B,r}(G_n - K_n) \leq 1/n$$

for all $n \in \mathbb{N}$.

For every $n \in \mathbb{N}$, let $f_n: T \rightarrow [0, 1]$ be a continuous function with $f_n|_{K_n} \equiv 1$ and $\text{supp}(f_n) \subset G_n$, then

$$\begin{aligned} r[\mu(E)(x) - \mu'(E)(x)] &= r[\mathcal{F}^s(x\chi_E) - \mathcal{F}^{s'}(x\chi_E)] \leq \\ &\leq r[\mathcal{F}^s(x\chi_E - xf_n)] + r[\mathcal{F}^s(xf_n) - \mathcal{F}^{s'}(xf_n)] + r[\mathcal{F}^{s'}(xf_n - x\chi_E)]. \end{aligned}$$

Hence it results that $r[\mu(E)(x) - \mu'(E)(x)] = 0$ (and therefore $\mu(E) = \mu'(E)$) because

$$r[\mathcal{F}^{s'}(xf_n) - \mathcal{F}^{s'}(xf_n)] = r[\mathcal{F}(xf_n) - \mathcal{F}(xf_n)] = 0$$

and

$$\lim_n r[\mathcal{F}^s(x\chi_E - xf_n)] = \lim_n r[\mathcal{F}^{s'}(xf_n - x\chi_E)] = 0$$

since $x\chi_E - xf_n$ takes non zero values in $G_n - K_n$,

$$\lim_n v_{B,r}(G_n - K_n) = 0,$$

and \mathcal{F}^s and $\mathcal{F}^{s'}$ are $(v_{B,r})$ -continuous.

Theorem 8. Let us suppose that the family $\{v_{B,r}: B \in \mathcal{B}, r \in \mathcal{R}\}$ is uniformly tight (i.e., given $E \in \Sigma$ and $\varepsilon > 0$ there exists a compact $K \subset T$ such that $K \subset E$ and $v_{B,r}(E - K) < \varepsilon$ for all $B \in \mathcal{B}$ and $r \in \mathcal{R}$), and let $\mathcal{F}: \mathcal{C} \rightarrow Z$ be a linear operator with continuous restrictions \mathcal{F}_B ($B \in \mathcal{B}$). Then there exists a $(v_{B,r})$ -continuous measure $\mu: \Sigma \rightarrow Y$ such that

$$\mathcal{F}(f) = \int_T f d\mu$$

for all $f \in \mathcal{C}$, if and only if the operator \mathcal{F} is $(v_{B,r})$ -continuous. In this case the measure μ is unique.

Proof. If such a measure exists, then the $(v_{B,r})$ -continuity of \mathcal{F} follows from Proposition 3, and the uniqueness of μ is deduced from Theorem 7.

Let us suppose that \mathcal{F} is $(v_{B,r})$ -continuous, then we will prove that 6.3 holds. If $E \in \Sigma$ we can find an increasing sequence of compact subsets $(K_n) \subset T$ and a decreasing sequence of open subsets $(G_n) \subset T$ such that $K_n \subset E \subset G_n$ and

$$v_{B,r}(G_n - K_n) \leq 1/n$$

for all $r \in \mathcal{R}$ and $B \in \mathcal{B}$. Let $f_n: T \rightarrow [0, 1]$ be a continuous function such that $f_n|_{K_n} \equiv 1$ and $f_n|_{T - G_n} \equiv 0$, for all $n \in \mathbb{N}$.

Define

$$(7) \quad \mathcal{G}_0(x\chi_E) = \lim_n \mathcal{F}(xf_n)$$

for all $x \in X$. Let us prove that this limit exists and that it is independent of the sequence (f_n) . If $n, m \in \mathbb{N}$ are such that $n \leq m$, then for every $B \in \mathcal{B}$ and $x \in B$, the function $xf_m - xf_n$ belongs to \mathcal{C}_B and vanishes outside of $G_m - K_n$. Moreover, for every $r \in \mathcal{R}$ we have

$$\lim_{\substack{m, n \rightarrow \infty \\ m \geq n}} v_{B,r}(G_m - K_n) = 0,$$

and the $(v_{B,r})$ -continuity of \mathcal{F} yields

$$\lim_{\substack{m, n \rightarrow \infty \\ m \geq n}} r[\mathcal{F}(xf_m - xf_n)] = 0.$$

So $\{\mathcal{F}(xf_n)\}_{n \in \mathbb{N}}$ is a Cauchy sequence and therefore the limit (7) exists.

Let us now consider other sequences (K'_n) , (G'_n) and (f'_n) satisfying the above conditions. If $B \in \mathcal{B}$ and $x \in B$, then the function $xf_n - xf'_n \in \mathcal{C}_B$ vanishes outside $(G_n \cup G'_n) - (K_n \cap K'_n)$ and

$$\lim_n v_{B,r}[(G_n \cup G'_n) - (K_n \cap K'_n)] = 0$$

holds for all $r \in \mathcal{R}$, and therefore the $(v_{B,r})$ -continuity of \mathcal{F} implies

$$\lim_n \mathcal{F}(xf_n) = \lim_n \mathcal{F}(xf'_n).$$

For a simple function $f = \sum_{i=1}^n x_i \chi_{E_i}$ let us define an operator

$$\mathcal{G}(f) = \sum_{i=1}^n \mathcal{G}_0(x_i \chi_{E_i}),$$

which is clearly linear and with continuous restrictions \mathcal{G}_B (since \mathcal{F} has these properties), so the proof will be complete if we prove that 6.3 holds.

Since \mathcal{F} is $(v_{B,r})$ -continuous, then for every $B \in \mathcal{B}$, $r \in \mathcal{R}$ and $\varepsilon > 0$ there exist $\delta > 0$ and $1 \geq \delta' > 0$ such that $r(\mathcal{F}(f)) < \varepsilon$ holds for all $f \in \mathcal{C}_B$ which verify $q_B \circ f \mid T - E \leq \delta'$ for some $E \in \Sigma$ with $v_{B,r}(E) < \delta$. Then, if $g \in \mathcal{S}_B$ is such that $q_B \circ g \mid T - E \leq \delta'$ for some $E \in \Sigma$ with $v_{B,r}(E) < \delta/2$, there exist $E' \in \Sigma$ and $f \in \mathcal{C}_B$ such that $v_{B,r}(E') < \delta/2$, $g \mid T - E' \equiv f \mid T - E'$ and

$$r(\mathcal{G}(g)) \leq r(\mathcal{F}(f)) + \varepsilon.$$

Therefore, $q_B \circ f \mid T - (E \cup E') \leq \delta'$, $v_{B,r}(E \cup E') < \delta$, $r(\mathcal{G}(g)) \leq 2\varepsilon$ and \mathcal{G} is $(v_{B,r})$ -continuous.

Moreover, if $B \in \mathcal{B}$ and the sequence $(h_n) \subset \mathcal{S}_B$ is uniformly convergent to $f \in \mathcal{G}_B$, then for every $\varepsilon > 0$ and $r \in \mathcal{R}$ there exist $\delta > 0$ and $1 \geq \delta' > 0$ such that $r(\mathcal{F}(g)) < \varepsilon$ for every $g \in \mathcal{C}_B$ which verifies $q_B \circ g \mid T - E \leq \delta'$ for some $E \in \Sigma$ with $v_{B,r}(E) < \delta/2$. Moreover, we can find $n_0 \in \mathbb{N}$, $f_{n_0} \in \mathcal{C}_B$ and $E \in \Sigma$ with $v_{B,r}(E) < \delta$ such that

$q_B(h_{n_0} - f) < \delta', f_{n_0} \mid T - E \equiv h_{n_0} \mid T - E$ and

$$r(\mathcal{G}(h_{n_0}) - \mathcal{F}(f_{n_0})) < \varepsilon.$$

Since $q_B(f_{n_0} - f) \mid T - E \leq \delta'$ and \mathcal{F} is $(v_{B,r})$ -continuous, it results that $r(\mathcal{F}(f_{n_0} - f)) < \varepsilon$. Therefore, $r(\mathcal{G}(h_{n_0}) - \mathcal{F}(f)) < 2\varepsilon$ holds and 6.3 is verified.

References

- [1] *Balbás, A. and P. Jiménez Guerra*: Un teorema de Radon-Nikodym para integrales bilineales. *Rev. R. Acad. Ci. Madrid*, 78 (1984), 217—220.
- [2] *Balbás, A. and P. Jiménez Guerra*: A Radon-Nikodym theorem for a bilinear integral in locally convex spaces. *Math. Japonica*, 32 (1987).
- [3] *Ballvé, M. E.*: Integración vectorial bilineal. U.N.E.D., 1984.
- [4] *Bartle, R. G.*: A general bilinear vector integral. *Studia Math.*, 15 (1956), 337—352.
- [5] *Bombal, F.*: Medida e integración en espacios bornológicos. *Rev. R. Acad. Ci. Madrid*, 65 (1981), 115—137.
- [6] *Bombal, F.*: El theorema de Radon-Nikodym en espacios bornológicos. *Rev. R. Acad. Ci. Madrid*, 65 (1981), 140—154.
- [7] *Devieve, C.*: Integration of vector valued functions with respect to vector valued measures. *Rev. Roum. Math. P. et Appl.*, 26 (1981), 943—957.
- [8] *Dobrákov, I.*: On integration in Banach spaces I. *Czech. Math. J.*, 20 (95) (1970), 511—536.
- [9] *Dobrákov, I.*: On integration in Banach spaces II. *Czech. Math. J.*, 20 (95) (1970), 680—695.
- [10] *Dobrákov, I.*: On representation of linear operators on $C_0(T, X)$. *Czech. Math. J.*, 21 (96) (1971), 13—30.
- [11] *Dunford, N. and J. Schwartz*: Linear operators part I. Interscience Pub., New York, 1958.
- [12] *Maynard, H. B.*: A Radon-Nikodym theorem for operator-valued measures. *Trans. Amer. Math. Soc.*, 173 (1972), 449—463.
- [13] *Rao Chivukula, R. and A. S. Sastry*: Product vector measures via Bartle integrals. *J. Math. Anal. and App.*, 96 (1983), 180—195.
- [14] *Sivasankara, S. A.*: Vector integrals and product of vector measures. Univ. Microfilm. Inter., Michigan, 1983.
- [15] *Smith, W. V. and D. H. Tucker*: Weak integral convergence theorems and operator measures. *Pacific J. Math.*, 111 (1984), 243—256.

Authors' address: Dpto. de Matemáticas Fundamentales, Facultad de Ciencias, U.N.E.D., Ciudad Universitaria, Madrid-28040 (Spain).