

Ján Duplák

Identities and deleting maps on quasigroups

Czechoslovak Mathematical Journal, Vol. 38 (1988), No. 1, 1–7

Persistent URL: <http://dml.cz/dmlcz/102198>

Terms of use:

© Institute of Mathematics AS CR, 1988

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

IDENTITIES AND DELETING MAPS ON QUASIGROUPS

JÁN DUPLÁK, Prešov

(Received March 22, 1985)

V. D. Belousov in [2] and M. A. Taylor in [6] have proved a theorem that is a generalization of a theorem of Belousov (see Theorem 2.1.1 in [3]). In this paper we give a generalization of these results and its applications.

1. PRELIMINARIES

An algebra $(Q, f_1, \dots, f_n) = (Q, F)$ is called an *algebra of quasigroups* if (Q, f_i) is a quasigroup for all $i \in \{1, 2, \dots, n\}$. An algebra (Q, f_1, f_2, f_3) is called a *primitive quasigroup* if there exists a permutation (ijk) of the set $\{1, 2, 3\}$ such that (Q, f_j) and (Q, f_k) are respectively the left and the right division groupoids of a quasigroup (Q, f_i) ; if f_i is denoted by \cdot , then we put $f_j = /$, $f_k = \backslash$, thus $(Q, \cdot, /, \backslash)$ means a primitive quasigroup. An algebra (Q, F) is called an *algebra of primitive quasigroups* if for each $f \in F$ there exist $g, h \in F$ such that (Q, f, g, h) is a primitive quasigroup. For every quasigroup (Q, f) there exists a primitive quasigroup (Q, f, g, h) and for every algebra of quasigroups (Q, F) there exists an algebra of primitive quasigroups (Q, G) with $F \subset G$.

Let (Q, A) be a quasigroup; we define $A[x, y] = z$ iff $A^{-1}[x, z] = y$ iff $^{-1}A[z, y] = x$ iff $^{-1}(A^{-1})[y, z] = x$ iff $(^{-1}A)^{-1}[z, x] = y$ iff $A^*[y, x] = z$. The set $\{A, ^{-1}A, A^{-1}, ^{-1}(A^{-1}), (^{-1}A)^{-1}, A^*\} = \Sigma A$ is called the *system of division operations of A*. An algebra (Q, F) is said to be an *algebra of parastrophic quasigroups* if $\Sigma f \subset F$ for each $f \in F$. For every algebra of quasigroups (Q, F) there exists exactly one algebra of parastrophic quasigroups $(Q, \Sigma F)$, where $\Sigma F = \bigcup \{\Sigma f; f \in F\}$.

Throughout the paper, for a quasigroup (Q, \cdot) we put $L_a x = a \cdot x$, $R_a x = x \cdot a$, $T_a x = x \backslash a$, $L_a^{-1} x = a \backslash x$, $R_a^{-1} x = x / a$, $T_a^{-1} x = a / x$,

$$T_0 = \{L, R, T, L^{-1}, R^{-1}, T^{-1}\}.$$

If a quasigroup operation is denoted by another symbol, say \circ , then we put $a \circ x =$

$= L_a^\circ x, x \circ a = R_a^\circ x, \dots, T_\circ^\circ = \{L^\circ, R^\circ, \dots\}$. If (Q, F) is an algebra of quasigroups we denote $T_\circ^F = \bigcup \{T_\circ^\circ; \circ \in F\}$.

A word on algebra $(Q, f_1, f_2, \dots, f_n)$ is a formal expression consisting of variables, brackets and operations f_1, \dots, f_n . The length $l(w)$ of a word w is the number of occurrences of variables in w . In the following, the set of all variables that occur in a word w will be denoted by $V(w)$.

Let $x, y, \dots, z, x_1, \dots, x_n, \dots$ be variables and let a_1, \dots, a_n, \dots be elements of a set Q . A *retraction map* with invariant variables x, y, \dots, z (or a retraction map, if there is no danger of confusion) is a map $w \mapsto w_1$, where $V(w) \subset \{x, y, \dots, z, x_1, \dots, x_n, \dots\}$ and w_1 is a formal expression obtained from a word w on an algebra (Q, F) by replacing each variable x_i in w by a_i , for all $x_i \in V(w)$; if a set of invariant variables of a retraction map is empty then the image of a word w is an element of Q .

An *identity* on an algebra (Q, F) is a pair (w, w') of words on (Q, F) that is written $w \simeq w'$. We say that an identity $w \simeq w'$ is valid on (Q, F) (or (Q, F) satisfies $w \simeq w'$) and write $w = w'$ if for every retraction map ϱ with no invariant variables $\varrho w = \varrho w'$, i.e. $\varrho w, \varrho w'$ are equal elements in Q . Words w, w' on (Q, F) are said to be *equivalent* if (Q, F) satisfies the identity $w \simeq w'$. A word w_1 is said to be a *subword of an identity* $w \simeq w'$ if w_1 is a subword of w or w' . Identities $w \simeq w', w_1 \simeq w'_1$ are called *equivalent* if the validity of one of them implies the validity of the other. Let ϱ be a retraction map and let w, w' be words on an algebra (Q, F) . We say that $\varrho w, \varrho w'$ are equivalent and write $\varrho w \simeq \varrho w'$ or $\varrho w = \varrho w'$ if for each retraction map σ with no invariant variables $\sigma w = \sigma w'$.

An identity $w \simeq w'$ on an algebra (Q, F) is called *balanced* if each variable occurs exactly twice in $w \simeq w'$, once on each side. The length $l(w \simeq w')$ of an identity $w \simeq w'$ is the sum of the lengths of w and w' .

Let w, w_1 be non-empty words on an algebra of quasigroups (Q, F) , and let w_1 be a subword of w . We define $Z(w, w_1)$ as the set of non-negative integers as follows:

- (i) $Z(w, w) = \{0\}$ for any word w ,
- (ii) $n \in Z(w, w_1)$ for $w \neq w_1$ iff there exists a word w_2 of length $n + 1$ such that either $w_1 \cdot w_2$ or $w_2 \cdot w_1$ is a subword of w .

Let f be an n -ary operation on a set Q and let $\alpha_1, \alpha_2, \dots, \alpha_{n+1}$ be permutations of Q . Then the n -ary operation $\alpha_{n+1} f(\alpha_1 x_1, \dots, \alpha_n x_n)$ is called an isotope of f ; the isotope will be denoted by $f^{(\alpha_1, \dots, \alpha_{n+1})}$. An algebra (Q, F) is called an *isotope of an algebra* (Q, F') if every $f \in F$ is an isotope of some $f' \in F'$ and conversely. It is known that every isotope of a quasigroup is a quasigroup. An isotope of a group is called a transitive quasigroup.

A property P of a quasigroup (Q, \cdot) is said to be *universal* if every isotope of (Q, \cdot) has property P . It is known that the transitivity of quasigroups is a universal property of quasigroups.

Let (Q, F) be an algebra of quasigroups, W the set of all words on the algebra (Q, F) and let (W, F) be the algebra of words on (Q, F) . Let x, y, \dots, z be variables. A map $\delta: W \rightarrow W, w \mapsto \delta w$, where δw is the word that we get from w by deleting all

variables except x, y, \dots, z , all superfluous operations and all superfluous brackets (in this order) is said to be a deleting map with invariant variables x, y, \dots, z (briefly a deleting map). Obviously, each deleting map is an endomorphism.

Let $(W, F), (W', F')$ be the algebras of words on algebras of quasigroups. A map $\omega: W \rightarrow W', w \mapsto \omega w$, where ωw is the word that we get from w if each operation symbol in w is replaced by an operation symbol in F' (equal operation symbols in w are not necessarily replaced by the same symbol) is called a *change operation map*; if each operation in F' is an isotope of an operation in F , we say that ω is a *change isotopy operation map*.

Let $w \simeq w'$ be an identity on an algebra of quasigroups (Q, F) and let P be a property of a quasigroup (Q, \cdot) , where the operation (\cdot) occurs in $w \simeq w'$; we say that P is an invariant (isotopy invariant) of $w \simeq w'$ if for every change operation (resp. change isotopy operation) map ω there exists an operation \circ occurring in $\omega w \simeq \omega w'$ such that (Q, \circ) has the property P .

1.1. Lemma. *Let $w \simeq w'$ be a balanced identity of length ≤ 6 on an algebra of quasigroups (Q, F) . Then $w \simeq w'$ is equivalent to at least one of the following identities*

$$(I) \quad x \cdot (y \square z) = x \circ (y \nabla z)$$

$$(II) \quad x \cdot (y \square z) = (x \circ y) \nabla z$$

given on the algebra $(Q, \Sigma F)$.

Proof. Let $V(w) = V(w') = \{x, y, z\}$. First, let $Z(w, x) = Z(w', x) = 1$; then $w = x \cdot (y \square z)$ and $w' = x \circ (y \nabla z)$ for convenient operations $\cdot, \square, \circ, \nabla$ in ΣF (if, for example, $w = (z \blacksquare y) \otimes x$ we put (\cdot) to be the dual of \otimes , \square the dual of \blacksquare). Now, let $Z(w, x) = Z(w', z) = 1$ i.e. there exists no $t \in V(w)$ such that $Z(w, t) = Z(w', t) = 1$; then obviously $w = x \cdot (y \square z)$, $w' = (x \circ y) \nabla z$ for convenient operations in ΣF . This completes the proof.

Identity (II) is called the *general associative law* (see [3, p. 76]).

1.2. Lemma. *A balanced identity $w \simeq w'$ of length ≤ 6 on an algebra of quasigroups (Q, F) is equivalent to the general associative law iff there exists $t \in V(w \simeq w')$ such that $Z(w, t) + Z(w', t) = 1$.*

Proof. Evident.

1.3. Theorem (about four quasigroups, see [1], [3]). *An algebra of quasigroups $(Q, \cdot, \circ, \square, \nabla)$ that satisfies the identity (II) is an algebra of transitive quasigroups all isotopic to the same group.*

Proof. From (II) it follows that

$$b \circ y = a \circ x \Leftrightarrow L_b L_y^\square = L_a L_x^\square \Leftrightarrow L_a^{-1} L_b = L_x^\square (L_y^\square)^{-1},$$

$$d \circ y = c \circ x \Leftrightarrow L_d L_y^\square = L_c L_x^\square \Leftrightarrow L_c^{-1} L_d = L_x^\square (L_y^\square)^{-1},$$

therefore

$$b \circ y = a \circ x \quad \text{and} \quad c \circ x = d \circ y \Leftrightarrow L_a^{-1} L_b = L_c^{-1} L_d.$$

The simultaneous equations $b \circ y = a \circ x$, $c \circ x = d \circ y$ have a solution for arbitrary three elements of the set $\{a, b, c, d\}$, so according to Theorem 2.1 in [5], (Q, \cdot) is a transitive quasigroup. For the rest of the proof see [3, p. 77].

1.4. Corollary. *The transitivity of a quasigroup is an invariant of the general associative law.*

2. GENERALIZATIONS OF A THEOREM OF BELOUSOV

2.1. Lemma. *Let w be a non-empty word on an algebra of quasigroups (Q, F) , let δ be a deleting and ϱ a retraction map, both with invariant variables x, y, \dots, z . Then there exists a change isotopy operation map ω such that $\varrho w = \omega \delta w$.*

Proof. We shall proceed by induction on the length of w . Let $l(w) = 1$; if δw is empty then $\varrho w = a \in Q$ and we put $\omega \delta w = a$ (an isotope of a 0-ary operation is a 0-ary operation); if δw is non-empty, say $w = x$, then $\delta w = x$ and we put $\omega = 1$. Further, assume that the theorem is valid for all words of length $< n$ and let $1 < l(w) = n$. Then $w = w_1 \cdot w_2$, where $l(w_1) < n$, $l(w_2) < n$, therefore there exists a change isotopy operation map ω such that $\varrho w_1 = \omega \delta w_1$ and $\varrho w_2 = \omega \delta w_2$. Let $\delta w_1, \delta w_2$ be non-empty words; since $\varrho(w_1 \cdot w_2) = \varrho w_1 \cdot \varrho w_2$ we define $\omega(\delta w_1 \cdot \delta w_2) = \omega \delta w_1 \cdot \omega \delta w_2$ (i.e. the operation (\cdot) is not changed). Since δ is an endomorphism, $\varrho w = \varrho w_1 \cdot \varrho w_2 = \omega \delta w_1 \cdot \omega \delta w_2 = \omega(\delta w_1 \cdot \delta w_2) = \omega \delta(w_1 \cdot w_2) = \omega \delta w$. Now suppose that δw_1 is empty and δw_2 is non-empty word. Then there exists a permutation α of Q such that $\varrho w = \alpha \varrho w_2$. Since $l(w_2) < n$, there exists a change isotopy operation map ω such that $\varrho w_2 = \omega \delta w_2$, where $V(\omega \delta w_2) \subset \{x, y, \dots, z\}$ (therefore $\sigma \omega \delta w_2 = \omega \delta w_2$ for every retraction map σ with invariant variables x, y, \dots, z). If $\omega \delta w_2 = w_3 \circ w_4$ for some non-empty words w_3, w_4 then we put $\alpha(w_3 \circ w_4) = w_3 \nabla w_4$, thus $\varrho w = \alpha \varrho w_2 = \alpha \omega \delta w_2 = \alpha(w_3 \circ w_4) = w_3 \nabla w_4$ and that is the above case. If $\delta w_2 = x$, then by the induction hypothesis $\varrho w_2 = \omega' \delta w_2 = \beta x$, where ω' is a change isotopy operation map and β is a permutation of Q ; now we put $\omega(\varepsilon) = \alpha \beta \varepsilon$, where ε is the identity map on Q . We have $\varrho w = \alpha \varrho w_2 = \alpha \beta x = \alpha \beta \varepsilon x = \omega(\varepsilon x) = \omega \delta w_2 = \omega \delta w$.

2.2. Theorem. *Let an algebra of quasigroups (Q, F) satisfy an identity $w \simeq w'$ and let δ be a deleting map. Let a universal property P of a quasigroup be an isotopy invariant of $\delta w \simeq \delta w'$. Then there exists (\cdot) in F such that (Q, \cdot) has the property P .*

Proof. Let ϱ be a retraction map with the same invariant variables as δ . By Lemma 2.1, there exists a change isotopy operation map ω such that $\varrho w = \omega \delta w$ and $\varrho w' = \omega \delta w'$. Since $w = w'$, $\varrho w = \varrho w'$ and hence $\omega \delta w = \omega \delta w'$. Therefore by the assumption of the theorem there exists an operation \circ in $\omega \delta w \simeq \omega \delta w'$ such that (Q, \circ) has the universal property P . Because any operation occurring in $\delta w \simeq \delta w'$ is an isotope of an operation in $\omega \delta w \simeq \omega \delta w'$ and any operation occurring in $\delta w \simeq \delta w'$ is in F , there exists (\cdot) in F that is an isotope of \circ . Since P is a universal property, (Q, \cdot) has the property P .

2.3. Corollary. *Let an algebra of quasigroups (Q, F) satisfy an identity $w \simeq w'$ and let δ be a deleting map. If $\delta w \simeq \delta w'$ is equivalent to the general associative law on $(Q, \Sigma F)$ then there exists \circ in F such that (Q, \circ) is a transitive quasigroup.*

Proof. Follows from the statement that the transitivity of a quasigroup is a universal property, from Corollary 1.4 and Theorem 2.2.

The identities $w \simeq w'$ that satisfy the conditions of Corollary 2.3 include, for example, the general medial law (see [3, p. 76]).

2.4. Theorem (of Belousov). *Let $w \simeq w'$ be a balanced identity on a quasigroup (Q, \cdot) , let $x \cdot y$ be a subword of w' and let neither $x \cdot y$ nor $y \cdot x$ be a subword of w . Then (Q, \cdot) is a transitive quasigroup.*

Proof. Since $x \cdot y, y \cdot x$ are not subwords of w , there exist at least three variables, say x, y, z , in w . Let δ be the deleting map with invariant variables x, y, z . Then obviously $Z(\delta w, z) = 0$ and $Z(\delta w', z) = 1$ so that $Z(\delta w, z) + Z(\delta w', z) = 1$. Evidently $\delta w \simeq \delta w'$ is a balanced identity, therefore by Lemma 1.2, the identity is equivalent to the general associative law.

3. SOME CLASSES OF TRANSITIVE QUASIGROUPS

3.1. Theorem. *Let $w = w'$ be an identity on an algebra of quasigroups (Q, F) such that*

- (1) $V(w \simeq w') = \{x, y, z\}$ and each variable occurs exactly twice in $w \simeq w'$,
- (2) $w \simeq w'$ is not of type $x \circ w_1 = x$ or $w_1 \circ x = x$,
- (3) if a word w_1 of length 2 is a subword of $w \simeq w'$ then $V(w_1)$ consists of two distinct variables,
- (4) if a word w_1 of length 3 is a subword of $w \simeq w'$ then $V(w_1) = \{x, y, z\}$,
- (5) for each $t \in \{x, y, z\}$, $\{Z(w, t), Z(w', t)\} \cap \{\{1, 2\}, \{2, 3\}\} = \emptyset$,
- (6) if words w_1, w_2 of length 2 occur in $w \simeq w'$ as subwords then $V(w_1) \neq V(w_2)$.

Then

- (i) for every change operation map ω the identity $\omega w \simeq \omega w'$ satisfies all conditions (1)–(6),
- (ii) there exists a balanced identity of length 6 on $(Q, \Sigma F)$ equivalent to the general associative law,
- (iii) there exists a group (Q, \circ) such that (Q, \cdot) is an isotope of (Q, \circ) for every operation (\cdot) in $w \simeq w'$.

Proof. (i) is easy. (ii) Without loss of generality assume $l(w) \leq 3$. Obviously there exists a variable, say x , that occurs exactly once on each side of $w \simeq w'$. Therefore, there exist $A, B, C, D \in T^F$ and subwords r, s, t, v of w with variables y, z such that if $w \simeq w'$ is rewritten with translations of (Q, F) we get

$$(a) A_r B_s x = x \quad \text{or} \quad (b) A_r B_s C_t x = x \quad \text{or} \quad (c) A_r B_s C_t D_v x = x$$

($A_x x = x$ contradicts (2)). In the case (a), we have $l(r) = l(s) = 2$, hence $r = y \cdot z$, $s = y \circ z$, but this contradicts (6). If (b) holds then $\{r, s, t\} = \{y, z, y \cdot z\}$; first we put $r = y$, $s = z$, $t = y \cdot z$ then $C_{y \cdot z} x = B_z^{-1} A_y^{-1} x$ is equivalent to (ii). If $r = y \cdot z$, $s = z$, $t = y$ then $B_z C_y x = A_{y \cdot z}^{-1} x$ is equivalent to (ii). Finally, if $r = y$, $s = y \cdot z$, $t = z$ then put $y \cdot z = u$ i.e. $y = u/z$ so that $B_u C_z x = A_{u/z}^{-1} x$ is equivalent to (ii). (iii) follows from Theorem 1.3.

3.2. Corollary. *A balanced identity $w \simeq w'$ of length ≤ 6 on an algebra of quasigroups satisfies all conditions (1)–(6) of Theorem 3.1 iff there exists $t \in V(w \simeq w')$ such that $Z(w, t) + Z(w', t) = 1$.*

Proof. Easy.

The identities that satisfy the conditions (1)–(6) of Theorem 3.1 and that are not balanced include, for example, the following identities (see [3, p. 59]): $yx \cdot xz = yz$, $yx \cdot zx = yz$, $xz \cdot xy = yz$, $x \cdot z(yx) = yz$, $x(yz \cdot yx) = z$, $(y \square x) \cdot (z \circ x) = y \triangle z$, $(x \square y) \cdot (y \circ z) = x \nabla z$.

3.3. Theorem. *Let $w \simeq w'$ be an identity on an algebra of quasigroups (Q, F) such that the conditions (1), (2) and (3) of Theorem 3.1 hold. Then there exists a balanced identity on $(Q, \Sigma F)$ equivalent to $w \simeq w'$.*

Proof. (i) Let a word w_1 of length 3 be a subword of w and let $V(w_1) = \{x, y\}$, i.e. (4) of Theorem 3.1 be not valid. Then $w_1 = x \circ (x \square y)$ for some convenient operations \circ, \square in F . If w_1 is expressed from $w \simeq w'$ then we get $w_1 \simeq w_2$, where $w_2 = z \cdot (z \nabla y)$ for some $\cdot, \nabla \in \Sigma F$. Further $w_1 \simeq w_2$ is rewritten with translations, so $L_x^{\circ} L_x^{\square} y = L_z \cdot L_z^{\nabla} y$ whence $L_z^{-1} L_x^{\circ} y = L_z^{\nabla} (L_x^{\square})^{-1} y$ is equivalent to (I). (ii) Let $Z(w, x) \in \{\{1, 2\}, \{2, 3\}\}$ i.e. (5) of Theorem 3.1 is not valid. If $Z(w, x) = \{1, 2\}$ then $w \simeq w'$ is equivalent to $x \cdot (x \circ (y \square z)) = y \nabla z$ and $x \circ (y \square z) = x \setminus (y \triangle z)$. If $Z(w, x) = \{2, 3\}$ then $w \simeq w'$ is equivalent to $x \cdot (x \circ (y \square (y \nabla z))) = z$ whence $Z(w, y) = \{0, 1\}$, that is (i). (iii) Let $x \circ y$, $x \square y$ be subwords of $w \simeq w'$. Then $w \simeq w'$ is equivalent to $(z \nabla (x \circ y)) \cdot (x \square y) = z$ as well as $z \nabla (x \circ y) = z / (x \square y)$. Finally, let (i)–(iii) be not valid. Then the statements (1)–(6) of Theorem 3.1 hold and therefore we can use the theorem.

3.4. Theorem. *Let $W \simeq W'$ be an identity on an algebra of quasigroups (Q, F) and let there exist variables $x, y, z \in V(W \simeq W')$ such that if δ is the deleting map with invariant variables x, y, z and $\delta W = w$, $\delta W' = w'$ then $w \simeq w'$ is an identity which satisfies conditions (1)–(6) of Theorem 3.1. Then there exists a group (Q, \circ) such that for every operation (\cdot) in $w \simeq w'$ (Q, \cdot) is an isotope of (Q, \circ) .*

Proof. Follows from Corollary 2.3 and Theorem 3.1.

3.5. Theorem. *Let w be a word on an algebra of quasigroups (Q, F) , $V(w) = \{x, y\}$, $A, B \in \mathbf{T}^F$ and let δ be the deleting map with invariant variables x, y, z . If $W = W'$ is an identity on (Q, F) such that $\delta W \simeq \delta W'$ is at least one of the*

identities

$$(7) \quad A_z(B_z x \cdot y) \simeq w$$

$$(8) \quad A_z(x \cdot B_z y) \simeq w$$

then there exists a group $(Q, +)$ such that for every operation (\cdot) in $\delta W \simeq \delta W', (Q, \cdot)$ is an isotope of $(Q, +)$.

Proof. Let us denote $w = x \circ y$, $z \nabla x = B_z x$, $A_z^{-1} t = z \square t$ for all x, y, t . Then from (7) we have

$$(i) \quad (z \nabla x) \cdot y \simeq z \square (x \circ y).$$

Since $(\circ) = (\cdot)^{(B_z, 1, A_z)}, (Q, \circ)$ is a quasigroup. Thus (i) is the general associative law. The rest of the proof is similar.

Among identities of type (7) belongs the identity $z(xz \cdot y) = x \cdot xy$ (see [4]). From theorem 3.5 directly follows that a quasigroup which satisfies this identity is a transitive quasigroup.

References

- [1] *B. Д. Белоусов*: Уравновешенные тождества в квазигруппах, Матем. сб. 70 (112) (1966), 55–97.
- [2] *B. Д. Белоусов*: Одна теорема об уравновешенных тождествах, Квазигруппы и латинские квадраты, Мат. Исслед. 71 (1983), 22–24.
- [3] *J. Dénes, A. D. Keedwell*: Latin squares and their applications, Akadémiai Kiadó, Budapest 1974.
- [4] *J. Duplák*: Rot — quasigroups, Mat. čas. 23 (1973), 223–230.
- [5] *J. Duplák*: On quasiidentities of transitive quasigroups, Math. Slovaca 34 (1984), 281–294.
- [6] *M. A. Taylor*: A generalization of a theorem of Belousov, Bull. London Math. Soc. 10 (1978), No 3, 285–286.

Author's address: 081 16 Prešov, Gottwaldova 1, Czechoslovakia (Katedra matematiky PdF UPJŠ).