

Josef Niederle

Completely meet-irreducible tolerances in distributive Noetherian lattices

Czechoslovak Mathematical Journal, Vol. 39 (1989), No. 2, 348–349

Persistent URL: <http://dml.cz/dmlcz/102306>

Terms of use:

© Institute of Mathematics AS CR, 1989

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

COMPLETELY MEET-IRREDUCIBLE TOLERANCES
IN DISTRIBUTIVE NOETHERIAN LATTICES

JOSEF NIEDERLE, Brno

(Received January 23, 1987)

In [3], it has been shown that the poset of all meet-irreducible tolerances in any finite distributive lattice L is order-isomorphic to the set of all intervals in the poset of all join-irreducible elements of L ordered by inclusion. Note that in the finite case the poset of all join-irreducible elements is order-isomorphic to the poset of all meet-irreducible elements. In this paper, the above result will be generalized to distributive Noetherian lattices.

Recall that a lattice L is said to be *Noetherian* if it satisfies the Ascending Chain Condition (ACC): each non-empty subset (or, equivalently, subchain) in L contains a maximal element (see [1] for details). A lattice is said to be *upper bounded* if it contains a greatest element, *conditionally complete* if any subset of L with an upper (lower) bound in L has also a least upper (greatest lower) bound in L , and *compact* if each element of L is compact, i.e. each subset X of L includes a finite subset Y such that $\bigvee X = \bigvee Y$ whenever $\bigvee X$ exists.

Proposition 1. *A lattice is Noetherian if and only if it is upper bounded, conditionally complete and compact.*

Proof. Let L be a Noetherian lattice. As L contains a maximal element, this is the greatest element of L . Let X be any nonempty subset of L . Define $X^\vee = \{x \in L \mid \exists_{n \in \mathbb{N}} \exists_{x_0, \dots, x_n \in X} (x = x_0 \vee \dots \vee x_n)\}$. This subset X^\vee contains a maximal element; denote it by m . Since $m \vee x \in X^\vee$ for any $x \in X$, it follows that m is an upper bound of X in L . If z is also an upper bound of X in L , then we have $m = x_0 \vee \dots \vee x_n \leq z$ for suitable elements $x_0, \dots, x_n \in X$, hence $m = \bigvee X$ in L . Consequently, L is conditionally complete. We have just shown that $\bigvee X \in X^\vee$, and so there exists a finite subset $Y \subseteq X$ such that $\bigvee X = \bigvee Y$. Thus the lattice L is compact. Conversely, let L be an upper bounded, conditionally complete and compact lattice. Let X be an arbitrary non-empty subchain in L . Then $\bigvee X \in X^\vee = X$, therefore X contains a maximal element. Q.E.D.

Completely (=strictly) meet-irreducible tolerances in distributive lattices were recognized in [2] and [3] as tolerances formed by a proper prime ideal that is maximal among all ideals not containing a given element b , and by a proper dual prime ideal

that is maximal among all dual ideals not containing a given element a , where $a < b$. See [2] and [3] for definitions and basic properties.

Proposition 2. *For a distributive Noetherian lattice L , the set of all completely meet-irreducible tolerances in L ordered by inclusion is order-isomorphic to the set of all intervals in the poset of all meet-irreducible elements of L with completely meet-irreducible greatest elements, which is ordered by inclusion, formally*

$$\mathbf{CM}(\mathbf{TL}(L)) \cong (\mathbf{M}(L) \times \mathbf{CM}(L)) \cap \leq .$$

Proof. Let $T \in \mathbf{CM}(\mathbf{TL}(L))$, i.e. $T = (I \times I) \cup (F \times F)$, $a, b \in L$, $a < b$, where I is a proper prime ideal which is maximal among all ideals not containing b , and F is a proper dual prime ideal which is maximal among all dual ideals not containing a . Then $J = L \setminus F$ is a proper prime ideal and $J \subseteq I$ holds. Ideals in Noetherian lattices are principal, and greatest elements of prime ideals are meet-irreducible (cf. [1]). If $\bigvee I = \bigwedge X$ for some $X \subseteq L$, then $\bigvee I \notin X$ would imply, in view of maximality of I with respect to b , that $b \leq x$ for any $x \in X$. This would yield $b \leq \bigwedge X = \bigvee I$, which contradicts the assumption. Therefore $\bigvee I$ is completely meet-irreducible. Define $h(T) = [\bigvee J, \bigvee I] \in (\mathbf{M}(L) \times \mathbf{CM}(L)) \cap \leq$. It is obvious that h is injective and order-preserving. It remains to show that it has an inverse.

For $[v, w] \in (\mathbf{M}(L) \times \mathbf{CM}(L)) \cap \leq$ define $g([v, w]) = ((L \setminus \langle v \rangle) \times (L \setminus \langle w \rangle)) \cup \langle w \rangle \times \langle w \rangle$. It is clear that $L \setminus \langle v \rangle$ is a dual prime ideal that is maximal among all dual ideals not containing v . Further, the element w is a lower bound of the set $X = \{x \in L \mid w < x\}$, which is not empty as it contains $\bigvee L$ by Proposition 1. Again by Proposition 1, the lattice L is conditionally complete, and therefore $\bigwedge X$ exists. As w is completely meet-irreducible, $w < \bigwedge X$. Denote $b = \bigwedge X$. Let K be an ideal that includes $\langle w \rangle$ and does not contain b . Then $w \leq \bigvee K$ but $b \not\leq \bigvee K$, and this implies $\bigvee K = w$. Hence $\langle w \rangle$ is maximal among all ideals not containing b . Consequently, $g([v, w])$ is a completely meet-irreducible tolerance. The mapping g is obviously injective and order-preserving. It is clear that $g = h^{-1}$. Q.E.D.

References

- [1] *G. Birkhoff*: Lattice theory. 3rd ed. Amer. Math. Soc., Providence 1979.
- [2] *J. Niederle*: A note on tolerance lattices. Časop. pěst. matem. 107 (1982), 221–224.
- [3] *J. Niederle*: On skeletal and irreducible elements in tolerance lattices of finite distributive lattices. Časop. pěst. matem. 107 (1982), 23–29.

Author's address: Viniční 60, 615 00 Brno 15, Czechoslovakia.