

Vladimír Müller; Wiesław Żelazko

$B(X)$  is generated in strong operator topology by two of its elements

*Czechoslovak Mathematical Journal*, Vol. 39 (1989), No. 3, 486–489

Persistent URL: <http://dml.cz/dmlcz/102320>

## Terms of use:

© Institute of Mathematics AS CR, 1989

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

$B(X)$  IS GENERATED IN STRONG OPERATOR TOPOLOGY  
BY TWO OF ITS ELEMENTS

VLADIMÍR MÜLLER, Praha and WIESŁAW ŻELAZKO, Warszawa

(Received November 19, 1987)

Let  $X$  be a real or complex Banach space and  $\tau$  a topology on  $B(X)$ . We say that  $B(X)$  is  $\tau$ -generated by its subset  $G$  if it coincides with the smallest  $\tau$ -closed subalgebra of  $B(X)$  containing  $G$ . In particular we say that  $B(X)$  is *strongly generated* by  $G$  if  $\tau$  is the strong operator topology. Similarly we say that  $B(X)$  is generated by  $G$  if  $\tau$  is the norm topology and  $B(X)$  is algebraically generated by  $G$  if  $\tau$  is the discrete topology.

As a motivation for the result presented here we mention a similar theorem known for  $X$  being a separable Hilbert space (see [1], [2], cf also [3] and the references therein) and the results given in [6]–[8]. In [6] it is shown that for any Hilbert space  $H$  there are two subalgebras  $A_1, A_2 \subset B(H)$  with square zero and therefore commutative such that the set  $G = A_1 \cup A_2$  algebraically generates  $B(H)$ . In [7] this result is extended to those Banach spaces  $X$  which are „ $n$ -th powers”,  $n > 1$ , i.e. which can be decomposed into direct sums  $X = X_1 \oplus \dots \oplus X_n$  of closed subspaces  $X_i$  which are mutually isomorphic. In [8] it is shown that for every Banach space  $X$  the algebra  $B(X)$  is strongly generated by two commutative subalgebras (in fact by two subalgebras of square zero) if  $\dim X > 1$ . Our present result improves the last one for separable Banach space so that instead of commutativity we have a single generation. Our proof is based upon the following result due to Ovsepian and Pełczyński ([4], Theorem 1) on the existence of total bounded biorthogonal systems in separable Banach spaces:

**Theorem [O–P].** *Let  $X$  be a separable Banach space. Then there is a sequence  $(x_i)$  of elements in  $X$  and a sequence  $(f_i)$  of functionals in  $X$  such that*

- (1)  $f_m(x_n) = \delta_{m,n}$  (the Kronecker symbol) for  $m, n = 1, 2, \dots$ .
- (2) The linear span of  $(x_i)$  is dense in  $X$  in the norm topology.
- (3) If  $f_n(x) = 0$  for all  $n$  then  $x = 0$ .
- (4)  $\sup_n \|x_n\| \|f_n\| = M < \infty$ .

By [5] we can assume  $M < 1 + \varepsilon$  for a given positive  $\varepsilon$ . In what follows we shall assume that the sequences  $(x_i)$  and  $(f_i)$  satisfying (1)–(4) are normalized so that they

satisfy

$$(5) \quad \|x_i\| = 1 \quad \text{and} \quad \|f_i\| \leq M \quad \text{for all } i.$$

We can now formulate our result.

**Theorem.** *Let  $X$  be a real or complex Banach space. There exist operators  $R$  and  $S$  in  $B(X)$  such that  $B(X)$  is strongly generated by  $\{R, S\}$ .*

*Proof.* For operators  $R$  and  $S$  in  $B(X)$  denote by  $\text{Alg}(R, S)$  the subalgebra of  $B(X)$  algebraically generated by  $R$  and  $S$ . It consists of linear combinations of the products of powers of  $R$  and  $S$ . We have to construct two operators  $R$  and  $S$  so that  $\text{Alg}(R, S)$  is strongly dense in  $B(X)$ . This means that for a given  $T$  in  $B(X)$ ,  $y_1, \dots, y_n$  linearly independent elements in  $X$  and a positive  $\varepsilon$ , there is an operator  $B$  in  $\text{Alg}(R, S)$  such that

$$(6) \quad \|By_i - Ty_i\| < \varepsilon \quad \text{for } 1 \leq i \leq n.$$

In order to have (6) it is sufficient to find operators  $B_i$  in  $\text{Alg}(R, S)$ ,  $1 \leq i \leq n$ , such that  $\|B_i y_i - T y_i\| < \varepsilon$  and  $B_i y_j = 0$  for  $j \neq i$ , because then (6) is satisfied by  $B = \sum_{i=1}^n B_i$ . Thus for proving strong density of  $\text{Alg}(R, S)$  in  $B(X)$  we have to show that for given linearly independent  $y_0, y_1, \dots, y_n$  in  $X$ , positive  $\varepsilon$  and an element  $z$  in  $X$  there is an operator  $B$  in  $\text{Alg}(R, S)$  such that

$$(7) \quad \|By_0 - z\| < \varepsilon \quad \text{and} \quad B y_i = 0 \quad \text{for } i = 1, 2, \dots, n.$$

Let  $(x_i)$  and  $(f_i)$  be a sequence of elements and functionals satisfying conditions (1)–(5) and put

$$(8) \quad R = \sum_{i=1}^{\infty} 2^{-i} (f_i \otimes x_{i+1}) \quad \text{and} \quad S = \sum_{i=1}^{\infty} 2^{-i} (f_{i+1} \otimes x_i),$$

where  $f \otimes x$  is the one-dimensional operator on  $X$  given by  $u \mapsto f(u)x$ . By (5) we have  $\|R\|, \|S\| \leq M$  and so  $R, S \in B(X)$ .

First we show that all operators  $f_m \otimes x_n$ ,  $1 \leq m, n < \infty$ , are in  $\text{Alg}(R, S)$ . To this end we prove the formula

$$(9) \quad f_m \otimes x_n = 2^p R^{n-1} (4SR - RS) S^{m-1}, \quad p = \binom{m}{2} + \binom{n}{2},$$

$$1 \leq m, n < \infty,$$

where  $\binom{k}{2}$  is 0 if  $k = 1$ . Using (1) we obtain immediately from (8)

$$(10) \quad R x_n = 2^{-n} x_{n+1} \quad \text{and} \quad S x_n = 2^{-n+1} x_{n-1}, \quad n = 1, 2, \dots,$$

where  $x_0 = 0$ . This implies

$$(11) \quad (4SR - RS) x_1 = x_1 \quad \text{and} \quad (4SR - RS) x_m = 0 \quad \text{for } m \neq 1.$$

Denote by  $A_{m,n}$  the right-hand side operator in formula (9). We see by (10) and (11)

that  $A_{m,n}x_k = 0$  for  $k \neq m$  and

$$A_{m,n}x_m = 2^p R^{n-1} (4SR - RS) 2^{-m+1} \dots 2^{-1} x_1 = 2^q R^{n-1} x_1 = x_n,$$

$$p = \binom{m}{2} + \binom{n}{2}, \quad q = \binom{n}{2}.$$

We infer that  $A_{m,n} = f_m \otimes x_n$  since both operators agree on linearly dense sequence  $(x_i)$  and formula (9) holds true.

Let  $y_0, y_1, \dots, y_n$  be linearly independent elements in  $X$ , let  $z$  be in  $X$  and let  $\varepsilon$  be a positive number. We shall be done if we find an operator  $B$  in  $\text{Alg}(R, S)$  so that relations (7) are satisfied. To this end observe that the sequences  $\{f_i(y_0)\}_{i=1}^\infty, \{f_i(y_1)\}_{i=1}^\infty, \dots, \{f_i(y_n)\}_{i=1}^\infty$  are linearly independent elements in  $l^\infty$ , otherwise there would exist non-zero coefficients  $\lambda_0, \dots, \lambda_n$  such that  $0 = \sum_{j=0}^n \lambda_j f_i(y_j) = f_i(\sum_{j=0}^n \lambda_j y_j)$  for all  $i$  and by (3) we would have  $\sum_{j=0}^n \lambda_j y_j = 0$  which is a contradiction. The linear independence of these sequences implies that there are indices  $i_0, i_1, \dots, i_n$  such that the finite sequences

$$\{f_{i_k}(y_0)\}_{k=0}^n, \{f_{i_k}(y_1)\}_{k=0}^n, \dots, \{f_{i_k}(y_n)\}_{k=0}^n$$

are linearly independent. This means that there are coefficients  $c_0, c_1, \dots, c_n$  such that

$$(12) \quad \sum_{k=0}^n c_k f_{i_k}(y_0) = 1 \quad \text{and} \quad \sum_{k=0}^n c_k f_{i_k}(y_j) = 0, \quad 1 \leq j \leq n.$$

Define now  $A = \sum_{k=0}^n c_k (f_{i_k} \otimes x_1)$ .

By (9) we have  $A \in \text{Alg}(R, S)$  and by (12) we conclude

$$(13) \quad Ay_0 = \sum_{k=0}^n c_k f_{i_k}(y_0) x_1 = x_1 \quad \text{and} \quad Ay_i = \sum_{k=0}^n c_k f_{i_k}(y_i) x_1 = 0, \\ 1 \leq i \leq n.$$

For a given positive  $\varepsilon$  and  $z$  in  $X$  we find by (2) a finite linear combination

$$(14) \quad z_0 = \sum_{j=1}^m b_j x_j$$

such that

$$(15) \quad \|z - z_0\| < \varepsilon.$$

Define  $B = \sum_{j=1}^m b_j (f_1 \otimes x_j) A$ . This is again an operator in  $\text{Alg}(R, S)$ . By (13) and (14) we have

$$By_0 = \sum_{j=1}^m b_j f_1(Ay_0) x_j = \sum_{j=1}^m b_j x_j = z_0$$

and

$$By_i = \sum_{j=1}^m b_j f_1(Ay_i) x_j = 0 \quad \text{for } i = 1, 2, \dots, n.$$

Thus by (15) we obtain (7) and the conclusion follows.

Remarks. 1° Without assuming that  $X$  is a separable space our conclusion fails to be true. This follows from the fact that for any operators  $R, S$  in  $B(X)$  and for any element  $x$  in  $X$  the orbit  $0(x) = \{Tx: T \in \text{Alg}(R, S)\}$  is separable. Thus for non-separable  $X$  there is an element  $z$  in  $X$  such that  $\text{dist}(z, 0(x)) = 1$ , and there is no operator  $T$  in  $\text{Alg}(R, S)$  satisfying  $\|Tx - (f \otimes z)x\| < 1$  if  $f(x) = 1$ .

2° It is still an open problem whether  $B(X)$  can be separable for an infinite-dimensional Banach space (for this information the authors are indebted to Tadeusz Figiel), and for most familiar Banach spaces the algebra  $B(X)$  is known to be non-separable. Thus for many cases (perhaps for all cases) our theorem cannot be improved by replacing strong generation by generation. In particular the algebra  $B(H)$  cannot be generated by two operators if  $H$  is an infinite-dimensional Hilbert space.

3° In our theorem we can replace strong generation by weak generation (generation in the weak operator topology). This follows from the fact that for linear subspaces of  $B(X)$  their closures in strong and in weak operator topologies coincide.

#### References

- [1] *Ch. Davis*: Generators of the ring of bounded operators, Proc. Amer. Math. Soc. 6 (1955), 970–972.
- [2] *E. A. Nordgren, M. Rajabali-pour, H. Radjavi, P. Rosenthal*: Quadratic operators and invariant subspaces, Studia Math. 88, in press.
- [3] *C. L. Olsen, W. R. Zame*: Some  $C^*$ -algebras with a single generator, Trans. Amer. Math. Soc. 215 (1976), 205–217.
- [4] *R. I. Ovspeian, A. Pelczyński*: Existence of a fundamental total and bounded biorthogonal sequence, Studia Math. 54 (1975), 149–159.
- [5] *A. Pelczyński*: All separable Banach spaces admit for every  $\varepsilon > 0$  fundamental total and bounded by  $1 + \varepsilon$  biorthogonal sequences, Studia Math. 55 (1976), 295–304.
- [6] *W. Żelazko*:  $B(H)$  is generated by two of its abelian subalgebras, Proc. Int. Conf. On invariant subspaces and allied topics (Delhi 15–19 Dec. 1986), in press.
- [7] *W. Żelazko*: Algebraic generation of  $B(X)$  by two subalgebras with square zero, Studia Math. 90, in press.
- [8] *W. Żelazko*:  $B(X)$  is generated in strong operator topology by two subalgebras with square zero, submitted.

*Authors' addresses*: V. Müller, Žitná 25, 115 67 Praha 1, Czechoslovakia (Matematický ústav ČSAV); W. Żelazko, Śniadeckich 8, 00-950 Warszawa, Poland (Instytut Matematyczny PAN).