

Surjit Singh Khurana; Jorge E. Vielma
Strict topology and perfect measures

Czechoslovak Mathematical Journal, Vol. 40 (1990), No. 1, 1–7

Persistent URL: <http://dml.cz/dmlcz/102354>

Terms of use:

© Institute of Mathematics AS CR, 1990

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

STRICT TOPOLOGY AND PERFECT MEASURES

SURJIT SINGH KHURANA, JORGE E. VIELMA, Iowa City

(Received September 10, 1986 in revised form February 27, 1987)

Let X be a completely regular Hausdorff space, E a normed space over K , the field of real or complex numbers, and $C_b(X, E)$ the space of all continuous E -valued functions on X . When $E = K = \mathbb{R}$, the set of real numbers, a topology β_p is defined, in [6], on $C_b(X, E)$ which gives its dual $M_p(X)$ the set of all Baire perfect measures on X . In this case we consider the general case when E is a normed space.

Notations of [3] will be used. X will always denote a completely regular Hausdorff space and E a normed space over K (scalars), the field of real or complex numbers. All linear spaces are taken over K . $C_b(X, E)$ will be denoted by $C_b(X)$ when $E = K$. \tilde{X} and νX will be respectively the Stone-Ćech compactification and real compactification of X . $M(X)$, $M_\sigma(X)$, $M(X, E')$ have the meanings as in ([3], p. 196). For a continuous function f from X into a topological space Y , \tilde{f} , \bar{f} will respectively denote its unique extensions to \tilde{X} and νX if extensions are possible. Notations of [7] for locally convex spaces will be used. A locally convex space F is strongly Mackey if every relatively countably compact subset of $(F', \sigma(F', F))$ is equicontinuous. A subset Z of X will be called a *zero set* if $Z = \bar{f}^{-1}\{0\}$ for some $f \in C_b(X)$. The norm topology on $C_b(X, E)$ is defined by $\|f\| = \sup_{x \in X} |f(x)|$. For a $\mu \in (C_b(X), \|\cdot\|)'$, $\tilde{\mu}: C(\tilde{X}) \rightarrow K$, is defined by $\tilde{\mu}(f) = \mu(f \setminus X)$.

As in [6], $M_p(X)$ denotes the space of all scalar-valued Baire perfect measures on X . A subset G in a completely regular space Y will be called *distinguished* if there exists a continuous mapping φ from Y onto a separable metric space such that $G = \varphi^{-1}(\varphi(G))$. The class of all distinguished subsets of \tilde{X} disjoint from X will be denoted by $\mathcal{D}(\tilde{X}) = D$. For a $D \in \mathcal{D}$, the topology γ_D on $C_b(X, E)$ is defined to be the one generated by the seminorms $\|\cdot\|_g$, as g varies over $B_D(X)$, all bounded scalar-valued functions on \tilde{X} , vanishing at infinity and zero on D , $\|f\|_g = \sup_{x \in \tilde{X}} (\|f\|(x) |g(x)|)$.

As in ([9], Theorem 2.4) it can be verified that γ_D is the finest locally convex topology agreeing with the topology of uniform convergence on compact subsets of $\tilde{X} \setminus D$, on norm-bounded subsets of $C_b(X, E)$ (note for a compact $C \subset \tilde{X} \setminus D$, $f \in C_b(X, E)$

norm sup of f over C is in the sense $\sup \|f\| \sim (C)$. We define $\beta_p = \Lambda\{\gamma_D: D \in \mathcal{D}\}$. As observed in [6] for every $D \in \mathcal{D}$, $D \cap \nu X = \emptyset$. For a function $f \in C_b(X, E)$, $\|f\|: X \rightarrow R$ is defined by $\|f\|(x) = \|f(x)\|$. A locally convex topology on $C_b(X, E)$ will be called *locally solid* if it has a 0-nbd. base consisting of absolutely convex sets V , such that $f \in V$, $\|g\| \leq f$ implies $g \in V$ (f, g in $C_b(X, E)$). For a duality $\langle F, G \rangle$, $A \subset F$. The polar of $A = A^0 = \{g \in G, |\langle a, g \rangle| \leq 1, \text{ for each } a \in A\}$. For any collection $\{A_\alpha\}_{\alpha \in I}$ of subsets of a locally convex space F , $\Gamma_\alpha A_\alpha$ will denote the absolutely convex hull of $\bigcup_{\alpha \in I} A_\alpha$. We define $M_p(X, E') = \{\mu \in M_\sigma(X, E'), \mu_x \in M_p(X) \text{ for every } x \in E\}$. It is known that $\mu \in M_p(X)$ implies $|\mu| \in M_p(X)$. First we prove that $\mu \in M_p(X, E')$ implies $|\mu| \in M_p(X)$.

Theorem 1. For a $\mu \in M_p(X, E')$, $|\mu| \in M_p(X)$.

Proof. From ([6], Theorem 2.1) it is sufficient to prove that $|\mu| \sim^*(D) = 0$ for every $D \in \mathcal{D}$. Take a Baire subset V_0 of \tilde{X} such that $|\mu| \sim(B) = 0$, for any Baire $B \subset V_0 \setminus D$. For any $x \in E$, $\|x\| \leq 1$, $|\mu_x| \leq |\mu| \sim$, and so $|\mu_x| \sim(B) = 0$. Since $|\mu_x|$ is perfect, this implies $|\mu_x| \sim(V_0) = 0$. So we get $|\mu_x|(V_0 \cap X) = 0$. Take any finite partition $\{V_i: 1 \leq i \leq n\}$ of $V_0 \cap X$, and any collection $\{x_i: 1 \leq i \leq n\}$ in E with $\|x_i\| \leq 1$, for every i . From what is proved above it follows that $|\sum_{i=1}^n \mu_{x_i}(V_i)| = 0$ and so $|\mu|(V_0 \cap X) = 0$. This means $|\mu| \sim(V_0) = 0$ and so $|\mu| \sim^*(D) = 0$. Thus $|\mu|$ is perfect.

The following theorem is a simple consequence of the definition of β_p .

Theorem 2.

- (i) $\beta_0 \leq \beta_p \leq \beta_1$.
- (ii) β_p is the finest locally convex topology agreeing with itself on norm-bounded subsets of $C_b(X, E)$.
- (iii) $C_b(X) \otimes E$ is dense in $(C_b(X, E), \beta_p)$ if it is dense in $(C_b(X, E), \beta_1)$ ([3], p. 206).

Proof follows easily from the definition of β_p .

Theorem 3. Let Y be a completely regular Hausdorff space and $\varphi: X \rightarrow Y$ a continuous mapping. Then the canonical mapping $(C_b(Y, E), \beta_p) \rightarrow (C_b(X, E), \beta_p)$. $(f \rightarrow f \circ \varphi)$ is continuous.

Proof. For a distinguished set D_0 in \tilde{Y} , $D_0 \subset \tilde{Y} \setminus Y$, $D = \tilde{\varphi}^{-1}(D_0) \in \mathcal{D}$, where $\tilde{\varphi}: \tilde{X} \rightarrow \tilde{Y}$ is the unique continuous extension. This implies that the mapping

$$(C_b(Y, E), \gamma_{D_0}) \rightarrow (C_b(X, E), \beta_p)$$

is continuous. The result follows now.

For a $\mu \in (C_b(X, E), \|\cdot\|)$, define $\lambda_\mu: C_b(X)^+ \rightarrow [0, \infty)$, $\lambda_\mu(f) = \sup \{|\mu(g)|: g \in C_b(X, E), \|g\| \leq f\}$.

Lemma 4. λ_μ is additive and positively homogeneous.

Proof. Take $f \in C_b(X)^+$, $g \in C_b(X)^+$, such that $f + g \geq \eta > 0$, on X , for some η .

Taking any $h \in C_b(X, E)$, with $\|h\| \leq f + g$, we get that

$$\left\| \frac{hf}{f+g} \right\| \leq f, \quad \left\| \frac{hg}{f+g} \right\| \leq g.$$

From

$$\mu(h) = \mu\left(\frac{hf}{f+g}\right) + \mu\left(\frac{hg}{f+g}\right)$$

it easily follows that

$$\lambda_\mu(f+g) \leq \lambda_\mu(f) + \lambda_\mu(g).$$

On the other hand, take h_1, h_2 in $C_b(X, E)$, $\|h_1\| \leq f$, $\|h_2\| \leq g$. This gives that $\|h_1 + h_2\| \leq f + g$, which implies $\lambda_\mu(f+g) \geq \lambda_\mu(f) + \lambda_\mu(g)$. Thus $\lambda_\mu(f) + \lambda_\mu(g) = \lambda_\mu(f+g)$. In particular, $\lambda_\mu(\frac{1}{2}) + \lambda_\mu(\frac{1}{2}) = \lambda_\mu(1)$. Take any f_1, g_1 in $C_b(X)^+$. From above it follows that $\lambda_\mu(f_1) + \lambda_\mu(g_1) + \lambda_\mu(1)$, $\lambda_\mu(f_1) + \lambda_\mu(g_1 + 1) = \lambda_\mu(f_1 + g_1 + 1) = \lambda_\mu(f_1 + g_1) + \lambda_\mu(1)$ and so $\lambda_\mu(f_1 + g_1) = \lambda_\mu(f_1) + \lambda_\mu(g_1)$. Also it is easily verified that $\lambda_\mu(pf) = p \lambda_\mu(f)$, for any $p \geq 0$ and $f \in C_b(X)^+$.

Theorem 5. *The space $(C_b(X, E), \beta_p)$ is locally solid, and has a 0-nbd. base consisting of solid absolutely convex sets.*

Proof. For any $D \in \mathcal{D}$, take $g_D: \tilde{X} \rightarrow R$, $g_D \equiv 0$ on D , g_D bounded and vanishing at infinity and put $V_D = \{f \in C_b(X, E), \sup_{x \in \tilde{X}} \|f\| \sim(x) g_D(x) \leq 1\}$ and $V = \bigcup_{D \in \mathcal{D}} V_D$.

This gives $V^0 = \bigcap V_D^0$, polar being taken in $(C_b(X, E), \beta_p)'$. Take $\mu \in V^0$. Since $\beta_p \leq \|\cdot\|$, $\mu \in (C_b(X, E), \|\cdot\|)'$. From Lemma 4 and the fact that each V_D is locally solid it follows that $\mu \in V^0$ if and only if $\lambda_\mu(\|g\|) \leq 1$, for every $g \in V_D$, for each D . Using Lemma 4, it is easily verified that $W = \{f \in C_b(X, E), \lambda_\mu(\|f\|) \leq 1, \text{ for every } \mu \in V^0\}$ is convex, contains V , is contained in V^{00} , and is locally solid. This proves the result.

Corollary 6. *A net $f_\alpha \rightarrow 0$, in $(C_b(X, E), \beta_p)$, if and only if $\|f_\alpha\| \rightarrow 0$ in $(C_b(X), \beta_p)$.*

Proof. Assuming $f_\alpha \rightarrow 0$ in $(C_b(X, E), \beta_p)$ take an absolutely convex, solid β_p 0-nbd W in $(C_b(X), \beta_p)$. This means for every $D \in \mathcal{D}$, there exists a $g_D \in B_D(\tilde{X})$, such that $W \supset P = \Gamma_D\{g \in C_b(X): \sup_{x \in \tilde{X}} |g| \sim(x) g_D(x) \leq 1\}$. Take $W_0 =$

$$= \Gamma_D\{f \in C_b(X, E): \sup_{x \in \tilde{X}} \|f\| \sim(x) g_D(x) \leq 1\}.$$

Since W is locally solid, $f_\alpha \in W_0$ implies $\|f_\alpha\| \in W$. This proves $\|f_\alpha\| \rightarrow 0$. Conversely suppose $\|f_\alpha\| \rightarrow 0$ in $(C_b(X), \beta_p)$. Fix a $y \in E$ with $\|y\| = 1$. We first prove that $\|f_\alpha\| y \rightarrow 0$ in $(C_b(X, E), \beta_p)$. Take an absolutely convex solid 0-nbd. V_0 in $(C_b(X, E), \beta_p)$. With above notations $V_0 \supset W_0$, for some g_D 's in $B_D(\tilde{X})$. Form P , as defined above, with these g_D 's. It is easy to see

$$\text{now that } \|f_\alpha\| \in P \text{ implies } \|f_\alpha\| = \sum_{i=1}^n \lambda_i g_{D_i}, \text{ with } \sum_{i=1}^n |\lambda_i| \leq 1, \sup_{x \in \tilde{X}} |g_i| \sim(x) g_{D(i)}(x) \leq 1, \text{ for } D(i) \in \mathcal{D}, g_i \in C_b(X), 1 \leq i \leq n, \text{ for some } n. \text{ Thus, } \|f_\alpha\| y = \sum_{i=1}^n \lambda_i g_i y$$

so $\|f_\alpha\| y \in W_0$. Since $\|f_\alpha\| \leq \| \|f_\alpha\| y \|$ and V_0 is locally solid $f_\alpha \in V_0$. The result follows now.

Theorem 7. *If X is realcompact $\beta_p \leq \beta_\infty$. In this case $C_b(X) \otimes E$ is dense in $(C_b(X, E), \beta_p)$.*

Proof. If $E = K = \mathbb{R}$, this result is proved in ([6], Prop. 4.6). First we prove that $f_\alpha \rightarrow 0$ in $(C_b(X, E), \beta_\infty)$ if and only if $\|f_\alpha\| \rightarrow 0$ in $(C_b(X), \beta_\infty)$. Suppose $\|f_\alpha\| \rightarrow 0$ in $(C_b(X), \beta_\infty)$. Take $x \in E$ with $\|x\| = 1$. We claim that $\|f_\alpha\| \otimes x \rightarrow 0$ in $(C_b(X, E), \beta_\infty)$. Let A be an equicontinuous subset of $M_\infty(X, E')$. This means $|A| = \{|\mu| : \mu \in A\}$ is an equicontinuous subset of $M_\infty(X)$. (This is proved in [3], Theorem 3.7, p. 202). Now $|\mu(\|f_\alpha\| \otimes x)| \leq |\mu|(\| \|f_\alpha\| \otimes x \|) = |\mu|(\|f_\alpha\|) \rightarrow 0$, uniformly for $\mu \in A$. This proves the claim. Now $\|f_\alpha\| \leq \| \|f_\alpha\| \otimes x \|$, and $\|f_\alpha\| \otimes x \rightarrow 0$. Since $(C_b(X, E), \beta_\infty)$ is locally solid ([3], Theorem 8.1), we get $f_\alpha \rightarrow 0$.

Now suppose $f_\alpha \rightarrow 0$ in β_∞ . This means $\|f_\alpha\| \rightarrow 0$ in $(C_b(X), \beta_\infty)$ ([3], Theorem 8.1). Since $\beta_p \leq \beta_\infty$, considered as topologies on $C_b(X)$ ([6], Theorem 4.1; this is proved for $K = \mathbb{R}$, but easily extends to the case when $K = \mathbb{C}$), we get $\|f_\alpha\| \rightarrow 0$ in β_p . By Corollary 6, this means $f_\alpha \rightarrow 0$ in $(C_b(X, E), \beta_p)$. Thus $\beta_p \leq \beta_\infty$, as topologies on $C_b(X, E)$. Since $C_b(X) \otimes E$ is dense in $(C_b(X, E), \beta_\infty)$, it follows that $C_b(X) \otimes E$ is dense in $(C_b(X, E), \beta_p)$.

Theorem 8. *If $C_b(X) \otimes E$ is dense in $(C_b(X, E), \beta_p)$, then*

- (i) *for any $\mu \in M_p(X, E')$, $L_1(\mu, X, E) \supset C_b(X, E)$;*
- (ii) *$(C_b(X, E), \beta_p)' = M_p(X, E)$, $L \in (C_b(X, E), \beta_p)'$ being related to corresponding $\mu \in M_p(X, E')$ by $L(f) = \mu(f)$, $\# f \in C_b(X, E)$.*

Proof is very similar to ([3], Theorem 5.3) and is omitted.

In the next theorem we give a new characterization of the topology β_p , which avoids the use of distinguished sets.

Let

$$\mathcal{F} = \{(Y, \varphi): Y \text{ a separable metric space,} \\ \varphi: X \rightarrow Y \text{ a continuous onto mapping}\}.$$

Every element $F = (Y, \varphi) \in \mathcal{F}$ gives rise to a linear mapping $T_F: C_b(Y, E) \rightarrow C_b(X, E)$, $h \rightarrow h \circ \varphi$.

Theorem 9. *β_p is the finest locally solid, locally convex topology V_p on $C_b(X, E)$ such that the mappings*

$$T_F: (C_b(Y, E), \beta_0) \rightarrow (C_b(X, E), V_p)$$

are continuous for every $F = (Y, \varphi) \in \mathcal{F}$.

Proof. When X is a separable metric space, $\beta_p = \beta_0$ (simple verification). Thus T_F is continuous when $V_p = \beta_p$. This means V_p exists and $V_p \geq \beta_p$. To prove $V_p \leq \| \cdot \|$ we take a sequence $\{f_n\} \subset C_b(X, E)$, $f_n \rightarrow 0$ in $\| \cdot \|$. Fix $e \in E$, $\|e\| = 1$. The continuous mapping $\varphi: X \rightarrow \mathbb{R}^N$, $\varphi(x) = \{\|f_n\|(x)\}$ maps X onto the separable metric space $Y = \varphi(X)$. The sequence $\{g_i\} \subset C_b(Y, E)$, $g_i(\{\|f_n\|(x)\}) = \|f_i\|(x) \otimes e$ uniformly

converges to 0 and so converges to 0 in $(C_b(Y, E), \beta_0)$. Thus $\|f_i\| \otimes e$ converges to 0 in $(C_b(X, E), V_p)$. Since $(C_b(X, E), V_p)$ is locally solid, this means $f_i \rightarrow 0$ in $(C_b(X, E), V_p)$.

To prove $V_p = \beta_p$ we first consider the case when $E = K$. Take a sequence $\{f_n\} \subset C_b(X)$, $f_n \downarrow 0$. The mapping

$$\varphi: X \rightarrow R^N, \quad x \rightarrow \{f_n(x)\}$$

is a continuous mapping from X onto a separable metric space $\varphi(X)$. Fix a $\mu \in (C_b(X), V_p)$. Then $\varphi * \mu \in M_t(\varphi(X))$ (note $\varphi * \mu(g) = \mu(g \circ \varphi)$). Since the sequence $\{g_n\} \subset C_b(\varphi(X))$, $g_n(\{f_i\}) = f_n$, monotonically decreases to 0, we get $\varphi * \mu(g_n) \rightarrow 0$, from which it follows that $\mu(f_n) \rightarrow 0$. Thus $(C_b(X), V_p) \subset M_\sigma(X)$. Next we will prove that $(C_b(X), V_p)' = M_p(X)$. Take a $\mu \in (C_b(X), V_p)'$. By the locally solid property of V_p , $|\mu| \in (C_b(X), V_p) \subset M_\sigma$. Thus for any metric space Y and every $\varphi: X \rightarrow Y$, a continuous onto mapping, $\varphi * |\mu| \in M_t(Y)$. Thus $|\mu| \in M_p(X)$ ([6], Lemma 2.2, p. 469). By ([6], Theorem 2.1) $\mu \in M_p(X)$. (Though results proved in [6] are for $k = R$, they easily extend to when $K = C$). Since $\beta_p \leq V_p$, we get $(C_b(X), V_p)' = M_p(X)$. Now we are ready to prove that when $E = K$, $\beta_p = V_p$. Take $H \subset M_p(X)$, H V_p -equicontinuous. There exists an absolutely convex solid V_p 0-nbd W in $C_b(X)$, such that $W \subset H^0 = \{g \in C_b(X): |\mu(g)| \leq 1, \forall \mu \in H\}$. If $g \in W$ and $\mu \in H$, then $|\mu|(|g|) = \sup \{|\mu(h)|: |h| \leq |g|, h \in C_b(X)\}$. Since W is solid, we get $|\mu|(|g|) \leq 1$. Thus $|H|$ is V_p -equicontinuous, and so for any $(Y, \varphi) \in \mathcal{F}$, $\varphi * |H|$ is β_0 -equicontinuous in $M_t(Y)$. By ([6], Prop. 2.6, p. 471), H is β_p -equicontinuous. This proves V_p and β_p on $C_b(X)$.

Now we come to the general case $(C_b(X, E), V_p)$. Fix $e \in E$, $\|e\| = 1$. We shall prove that the mapping $\psi: (C_b(X), \beta_p) \rightarrow (C_b(X, E), V_p)$, $g \rightarrow g \otimes e$ is continuous. Taking any $\mathcal{F} = (Y, \varphi) \in F$, the mappings $T_F: (C_b(Y, E), \beta_0) \rightarrow (C_b(X, E), V_p)$, $g \rightarrow g \circ \varphi$, and $\psi_0: (C_b(Y), \beta_0) \rightarrow (C_b(Y, E), \beta_0)$, $g \rightarrow g \otimes e$ are continuous. Let $\psi_1: (C_b(Y), \beta_0) \rightarrow (C_b(X), \beta_p)$, $g \rightarrow g \circ \varphi$. For any locally solid, absolutely convex 0-nbd U in $(C_b(X, E), V_p)$, $\psi_0^{-1}(T_F^{-1}(U))$ is a 0-nbd in $(C_b(Y), \beta_0)$. Since $\psi_1^{-1}(\psi^{-1}(U)) = \psi_0^{-1}(T_F^{-1}(U))$ (simple verification), and $V_p = \beta_p$ on $C_b(X)$, we get ψ is continuous. Now take a net $f_\alpha \rightarrow 0$ in $(C_b(X, E), \beta_p)$. This gives $\|f_\alpha\| \rightarrow 0$ in $(C_b(X), \beta_p)$. Since ψ is continuous, $\|f_\alpha\| \otimes e \rightarrow 0$ in $(C_b(X, E), V_p)$. Since $(C_b(X, E), V_p)$ is locally solid and $\|f_\alpha\| \leq \|(\|f_\alpha\| \otimes e)\|$, we get $f_\alpha \rightarrow 0$ in $(C_b(X, E), V_p)$. This proves the theorem.

Theorem 10. *Let X be a P -space ([2], p. 62). If $C_b(X) \otimes E$ is dense in $(C_b(X, E), \beta_p)$, then $(C_b(X, E), \beta_p)$ is Mackey. If E is a Banach space $(C_b(X, E), \beta_p)$ is strongly Mackey.*

Proof. Putting $F = C_b(X, E)$ and $F' = M_p(X, E')$, let A be a norm-bounded relatively countably compact subset of $(F', \sigma(F', F))$. Since νX is topologically complete and P -space, $M_\sigma(\nu X) = M_\tau(\nu X)$ ([10], p. 469). Further, since νX is also

a P -space ([2], p. 125), $M_t(vX) = M_t(vX)$. Now using the fact that vX is realcompact, we get $M_p(vX) \subset M_\infty(vX)$. Combining these facts we get $M_t(vX, E') = M_p(vX, E')$. From this it easily follows that $(C_b(vX, E), \beta_p)' = (C_b(vX, E), \beta_c)'$. Since $(C_b(vX, E), \beta_0)$ is Mackey ([4]) and $\beta_0 \leq \beta_p$, we get $\beta_0 = \beta_p$. By Theorem 3, $(C_b(vX, E), \beta_p) \rightarrow (C_b(X, E), \beta_p)$ ($f \rightarrow f|_X$) is continuous. This means a $\mu \in M_p(X, E')$ gives a

$$\hat{\mu} \in M_p(vX, E'),$$

$$\hat{\mu}(f) = \mu(f|_X), \quad f \in C_b(vX, E).$$

Also it is a simple verification that $|\hat{\mu}|^\sim = |\mu|^\sim$. Thus $\hat{A} = \{\hat{\mu}: \mu \in A\}$ is norm-bounded and $\sigma(M_p(vX, E'), C(vX, E))$ relatively countable subset of $M_p(vX, E')$. Since $M_p(vX, E') = M_t(vX, E')$, \hat{A} is a β_r -equicontinuous subset of $(C_b(vX, E), \beta_r)'$. There exists an increasing sequence of compact subsets K_n of vX , such that

$$|\mu|^\sim(\tilde{X} \setminus K_n) \leq \frac{p}{(p+1)(n+1)2^{n+1}},$$

for each $\mu \in H$, where $p = \sup\{|\mu|(X): \mu \in H\}$ [4]. Take any $D \in \mathcal{D}$, $D \subset \tilde{X} \setminus X$. This means $D \subset \tilde{X} \setminus vX$. Define $g_D: \tilde{X} \rightarrow R$,

$$g_D = \sum_{i=1}^{\infty} \frac{4(p+1)}{n} \chi_{(K_n \setminus K_{n-1})} \quad (K_0 = \emptyset).$$

g_D vanishes at infinity.

Take $f \in C_b(X, E)$, $\|f\|^\sim(x) g_D(x) \leq 1$, for every $x \in \tilde{X}$. This gives $\|f\|^\sim \leq n/4(p+1)$ on $K_n \setminus K_{n-1}$. For a $\mu \in H$,

$$|\mu(f)| \leq |\mu|(\|f\|) = |\mu|^\sim(\|f\|^\sim) = \sum_{i=1}^{\infty} \int_{K_n \setminus K_{n-1}} \|f\|^\sim d|\mu|^\sim \leq$$

$$\leq \sum \frac{n}{4(p+1)} \frac{1}{n} p \leq 1.$$

This proves A^0 is a 0-nbd. in $(C_b(X, E), \beta_p)$. This proves the theorem.

Theorem 11. *Let X be a paracompact locally compact D_0 -space and E is a normed space. Then $(C_b(X, E), \beta_p)$ is Mackey. If E is a Banach space, then $(C_b(X, E), \beta_p)$ is strongly Mackey.*

Proof. A paracompact locally compact space is topologically complete ([1], Theorem 11.2, p. 92). Since X is a D_0 -space, this implies X is realcompact ([6], Theorem 4.1). Thus $M_p(X) = M_t(X)$, which implies $M_p(X, E') = M_t(X, E')$. Since the result is known to be true in the case of topology β_0 ([5]), and $\beta_p \geq \beta_0$, the result now follows.

References

- [1] *W.W. Comfort, S. Negrepointis*: Continuous pseudometrics, Marcel Dekker, New York 1975.
- [2] *L. Gilman, M. Jerison*: Rings of continuous functions, D. van Nostrand Co., Inc., New York 1960.
- [3] *S. S. Khurana*: Topologies on spaces of vector-valued continuous functions, Trans. Amer. Math. Soc. *24* (1978) 195—211.
- [4] *S. Choo, S. S. Khurana*: Strict topology and P -spaces, Proc. Amer. Math. Soc. *61* (1976) 280—284.
- [5] *S. Khurana*: Strict topology on paracompact locally compact spaces, Canad. J. Math. *29* (1977) 216, 219; Corrigendum *30* (1978) 671.
- [6] *G. Koumoullis*: Perfect, u -additive measures and strict topologies, Illinois J. Math. *26* (1982) 466—478.
- [7] *H. H. Schaefer*: Topological vector spaces, Macmillan, New York, 1966.
- [8] *A. H. Schuchat*: Integral representation theorems in topological vector spaces, Trans. Amer. Math. Soc. *172* (1972) 373—397.
- [9] *F. D. Sentilles*: Bounded continuous functions on completely regular spaces, Trans. Amer. Math. Soc. *168* (1972) 311—336.
- [10] *Robert Wheeler*: The strict topology for P -spaces, Proc. Amer. Math. Soc. *41* (1973) 466—472.

Authors' address: The University of Iowa, Iowa City, Iowa 52242, U.S.A.