

Ján Jakubík

Cyclically ordered groups with unique addition

Czechoslovak Mathematical Journal, Vol. 40 (1990), No. 3, 534–538

Persistent URL: <http://dml.cz/dmlcz/102406>

Terms of use:

© Institute of Mathematics AS CR, 1990

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

CYCLICALLY ORDERED GROUPS WITH UNIQUE ADDITION

JÁN JAKUBÍK, Košice

(Received December 28, 1988)

Linearly ordered groups with unique addition (for definitions, cf. below) were investigated by T. Ohkuma [8]. The case of lattice ordered groups having this property was dealt with by P. Conrad and M. Darnel [2], and by the author [4].

Cyclically ordered groups were studied in [1], [5], [6], [7], [9]–[13]. The notion of cyclically ordered group is a generalization of the notion of linearly ordered group.

An example of cyclically ordered group is the additive group K of all complex numbers z with $|z| = 1$, where the cyclic order is defined in a natural way.

In this note the notion of cyclically ordered group with unique addition will be introduced. Let \mathcal{C}_u be the class of all cyclically ordered groups G such that (i) G fails to be linearly ordered, and (ii) G has a unique addition.

It will be proved that each element of \mathcal{C}_u is isomorphic to a subgroup of K with the inherited cyclic order. Next it will be shown that there are exactly $2^{2^{\aleph_0}}$ nonisomorphic types of cyclically ordered groups belonging to \mathcal{C}_u .

1. PRELIMINARIES

A linearly ordered group $G_1 = (G; \leq, +_1)$ is said to have a unique addition, if whenever $G_2 = (G; \leq, +_2)$ is linearly ordered group such that the neutral element of the group $(G; +_1)$ is the same as the neutral element of the group $(G; +_2)$, then the operation $+_1$ coincides with the operation $+_2$.

For the notion of cyclically ordered group cf., e.g., Fuchs [3], Chap. IV. Section 6. In the present paper we shall apply the same terminology and denotations concerning cyclically ordered groups as in [7]: in particular, the group operation in a cyclically ordered group will be written additively and the relation of cyclic order will be denoted by the symbol $[x, y, z]$ (or, shortly, by $[]$).

The cyclically ordered group K mentioned above can be described, up to isomorphism, as follows: K is the set of all reals x with $0 \leq x < 1$, the operation $+$ is the addition mod 1, and for $a, b, c \in K$ we have $[a, b, c]$ if and only if

$$(1) \quad a < b < c \text{ or } b < c < a \text{ or } c < a < b$$

is valid.

The relations between linearly ordered groups and cyclically ordered groups are well-known; cf., e.g., [6], Section 3.

For cyclically ordered groups we can apply a definition analogous to that applied above for linearly ordered groups, namely:

A cyclically ordered group $G_1 = (G; [\], +_1)$ will be said to have a unique addition if, whenever $G_2 = (G; [\], +_2)$ is a cyclically ordered group with $0_1 = 0_2$ (where 0_i is the neutral element of G_i ($i = 1, 2$)), then the operation $+_1$ coincides with the operation $+_2$.

2. ELEMENTS OF \mathcal{C}_u AS SUBGROUPS OF K

Let us denote by \mathcal{O}_u the class of all linearly ordered groups with unique addition. Next let \mathcal{C}_u^0 the class of all cyclically ordered groups with unique addition.

Let $G_1 = (G; [\], +)$ be a cyclically ordered group. In view of the Representation Theorem (which is due to Swierczkowski [10] (cf. also [7], Theorem 1.1)) there is a linearly ordered group L such that G_1 is isomorphic to a subgroup of the cyclically ordered group $K \otimes L$ (for denotations, cf. [7], Section 1). Hence without loss of generality we can suppose that G_1 is a subgroup of $K \otimes L$.

We denote by $G_1(K)$ and $G_1(L)$ the natural projection of G_1 into K or into L , respectively. Also, without loss of generality we can assume that $G_1(L) = L$.

2.1. Lemma. *Let $G_1(K) = \{0\}$. Then $G_1 \in \mathcal{C}_u^0$ if and only if $G_1 = L \in \mathcal{O}_u$.*

Proof. From $G_1(K) = \{0\}$ we obtain that $G_1(L) = G_1$, and thus in view of the above assumption, $G_1 = L$. Hence G_1 is linearly ordered. It is easy to verify that G_1 as cyclically ordered group has a unique addition if and only if G_1 as linearly ordered group has a unique addition.

Put $G_0 = \{g \in G; g(K) = 0\}$. Then in view of [6], Cor. 3.6, G_0 is the largest linearly ordered subgroup of G_1 ; moreover, G_0 is a normal subgroup of G_1 and it is c -convex in G_1 ([6], Section 4).

If $G_0 = \{0\}$, then for $g = (a, x) \in G$ we put $\varphi(g) = a$; it is easy to verify that φ is an isomorphism of G onto $G_1(K)$. Hence we have

2.2. Lemma. *Let $G_0 = \{0\}$. Then G_1 is isomorphic to a cyclically ordered subgroup of K .*

2.3. Lemma. *Let $G_1(K) \neq \{0\}$. Assume that G_1 belongs to \mathcal{C}_u . Then $G_0 = \{0\}$.*

Proof. By way of contradiction, suppose that $G_0 \neq \{0\}$. Since $G_1(K) \neq \{0\}$ there is $g \in G_1$ with $g = (a, y)$ such that $0 \neq a \in K$ and $y \in L$.

We define a mapping $\psi: G \rightarrow G$ as follows. We choose a fixed $z \in G_0$, $z \neq 0$ and for each $g_1 \in G$ we put

$$\begin{aligned} \psi(g_1) &= g_1 + z \quad \text{if } g_1 \in g + G_0, \quad \text{and} \\ \psi(g_1) &= g_1 \quad \text{otherwise.} \end{aligned}$$

Then ψ is an automorphism of the cyclically ordered set $(G; [\])$ such that $\psi(0) = 0$. Now we define a binary operation $+_1$ on G by putting

$$g_1 +_1 g_2 = \psi(\psi^{-1}(g_1) + \psi^{-1}(g_2))$$

for each $g_1, g_2 \in G$. The operation $+_1$ does not coincide with $+$ (since $g + g \neq g +_1 g$) and $(G; [\], +_1)$ is a cyclically ordered group whose neutral element is 0; in this way we arrived at a contradiction.

From 2.2 and 2.3 we obtain as a corollary

2.4. Theorem. *Let G_1 be a cyclically ordered group belonging to \mathcal{C}_u . Then G_1 is isomorphic to a cyclically ordered subgroup of K .*

3. THE POWER OF THE CLASS \mathcal{C}_u

Let us denote by R the additive group of all reals with the natural linear order.

3.1. Lemma. *Let $(G'; \leq, +) \in \mathcal{O}_u$. Assume that $(G'; \leq, +)$ is an l -subgroup of R such that $1 \in G'$. Put $G = \{x \in G' : 0 \leq x < 1\}$ and let $+_1$ be the binary operation on G defined as to be the addition mod 1. For $x, y, z \in G$ put $[x, y, z]$ if the relation (1) is valid. Then*

- (i) $G_1 = (G; [\], +_1)$ is a cyclically ordered group;
- (ii) if $G \neq \{0\}$; then $G_1 \in \mathcal{C}_u$.

Proof. (i) can be verified by a routine calculation; it will be omitted. Assume that $G \neq \{0\}$. Then G_1 fails to be a linearly ordered group. It remains to show that G_1 has a unique addition.

Let $+_2$ be a binary operation on G such that $G_2 = (G; [\], +_2)$ is a cyclically ordered group such that its neutral element is 0. Assume that $+$ is the original group operation on R (i.e., the addition of reals); hence $+$ is also the group operation of G' . Let us define a new binary operation $+^2$ on G' in the following manner:

Let $x, y \in G'$. There are uniquely determined integers x_0, y_0 and uniquely determined elements x^1, y^1 of G such that $x = x^0 + x^1$ and $y = y^0 + y^1$. We put

$$(2) \quad x +^2 y = (x^0 + y^0) + z^1,$$

where $z^1 = x^1 +_2 y^1$ if $x^1 +_2 y^1 < 1$, and $z^1 = (x^1 +_2 y^1) - 1$ otherwise. Then the set G' with the natural linear order and with the operation $+^2$ is a linearly ordered group with the neutral element 0. Since G' has a unique addition, the operation $+^2$ coincides with the operation $+$ on G' . Thus from (2) we infer that the operation $+_2$ on G must coincide with the operation $+_1$. Therefore G_1 has a unique addition.

3.2. Lemma. *Let $(G'', \leq, +)$ be an l -subgroup of R with $1 \in G''$. Let $G^* = \{x \in G'' : 0 \leq x < 1\}$ and let us define the cyclically ordered group $G_3 = (G^*, [\], +_2)$ analogously as we did for G_1 in 3.1 with the distinction that we now have G^* instead of G' . Suppose that the cyclically ordered groups G_1 and G_3*

are isomorphic. Then the linearly ordered groups $(G'; \leq, +)$ and $(G^*; \leq, +)$ are isomorphic as well.

Proof. Let φ be an isomorphism of G_1 onto G_3 . For $x \in G'$ let x^0 and x^1 be as in the proof of 3.2. Put

$$\psi(x) = x^0 + \varphi(x^1).$$

Then ψ is an isomorphism of $(G'; \leq, +)$ onto $(G^*; \leq, +)$.

The following theorem is the main result of [8].

3.3. Theorem. (Ohkuma) *There exists a subset $\{G^i: i \in I\}$ of \mathcal{C}_u such that*

- (i) $\text{card } I = 2^{2^{\aleph_0}}$;
- (ii) if $i(1)$ and $i(2)$ are distinct elements of I , then $G^{i(1)}$ fails to be isomorphic to $G^{i(2)}$;
- (iii) for each $i \in I$, G^i is an l -subgroup of R and contains all rational numbers.

(The assertions (i) and (ii) are expressed in Theorem 3 of [8] (cf. also [2]); (iii) follows from the constructions established in Section 2 and Section 3 of [8].)

If $(G'; \leq, +) = G^i$ for some $i \in I$, then let $\chi(G^i) = G_1$, where G_1 is as in 3.1. In view of 3.2, if $i(1)$ and $i(2)$ are distinct elements of I , then $\chi(G^{i(1)})$ is not isomorphic to $\chi(G^{i(2)})$.

Now from 2.4, 3.1 and 3.3 we obtain:

3.4. Theorem. *There are exactly $2^{2^{\aleph_0}}$ nonisomorphic types of cyclically ordered groups belonging to \mathcal{C}_u .*

Next, from 3.3, 3.4 and 2.1 we infer that in 3.4 the class \mathcal{C}_u can be replaced by \mathcal{C}_u^0 .

References

- [1] Černák, Š., Jakubík, J.: Completion of a cyclically ordered group. Czech. Math. J. 37, 1987, 157–174.
- [2] Conrad, P., Darnel, M.: 1-groups with unique addition. Algebra and Order. Proc. First Int. Symp. Ordered Algebraic Structures, Luminy-Marseille 1984, Helderman Verlag, Berlin 1986, 15–27.
- [3] Fuks, L.: Častično uporjadočennye algebraičeskie sistemy, Moskva 1965.
- [4] Jakubík, J.: On lattice ordered groups having a unique addition. Czech. Math. J. 40, 1990, 311–314.
- [5] Jakubík, J.: Retracts of abelian cyclically ordered groups. Archivum Math. 25, 1989, 13–18.
- [6] Jakubík, J., Pringerová, G.: Representations of cyclically ordered groups. Čas. pěst. matem. 113, 1988, 197–208.
- [7] Jakubík, J., Pringerová, G.: Radical classes of cyclically ordered groups. Math. Slovaca 38, 1988, 255–268.
- [8] Ohkuma, T.: Sur quelques ensembles ordonnés linéairement. Fund. Math. 43, 1955, 326–337.
- [9] Rieger, L.: On ordered and cyclically ordered groups, I, II, III. Věstník král. české spol. nauk 1946, 1–31; 1947, 1–33; 1948, 1–26 (In Czech.)
- [10] Swierczkowski, S.: On cyclically ordered groups. Fundam. Math. 47, 1959, 161–166.

- [11] *Zabarina, A. J.*: K teorii cikličeski uporjadočennych grupp. *Matem. zametki* 31, 1982, 3–12.
- [12] *Zabarina, A. J.*: O linejnom i cikličeskom porjadkach v gruppe. *Sibir. matem. žurn.* 26, 1985, 204–207.
- [13] *Zabarina, A. J., Pestov, G. G.*: K teoreme Sverčkovskogo. *Sibir. matem. ž.* 25, 1984, 46–53.

Author's address: 040 01 Košice, Grešákova 6, Czechoslovakia (Matematický ústav SAV, dislokované pracovisko v Košiciach).