

Igor Edmundovich Zverovich; Vadim E. Zverovich
A note on domatically critical and cocritical graphs

Czechoslovak Mathematical Journal, Vol. 41 (1991), No. 2, 278–281

Persistent URL: <http://dml.cz/dmlcz/102460>

Terms of use:

© Institute of Mathematics AS CR, 1991

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON DOMATICALLY CRITICAL AND COCRITICAL GRAPHS

I. E. ZVEROVICH, V. E. ZVEROVICH, Minsk

(Received February 28, 1989)

This paper deals with domatically critical and cocritical graphs. Two problems concerning such graphs are settled.

With minor adaptations, we adopt the terminology of Harary [3].

Let $G = (V(G), E(G))$ be an undirected graph with no loops and multiple edges. A set D of vertices in G is said to be a *dominating set* if every vertex not in D is adjacent to some vertex in D . A set of vertices S is independent if no two vertices in S are adjacent. A domatic partition (D -partition) of G is a partition of $V(G)$ into dominating sets. The maximum order of a D -partition of G is called the *domatic number* of G and is denoted by $d(G)$.

The join of two graphs G, H is the graph $G + H = (V(G) \cup V(H), E)$ where $E = E(G) \cup E(H) \cup \{(u, v) \mid u \in V(G), v \in V(H)\}$. We denote by $p(G)$ and $q(G)$ the number of vertices and edges of G , respectively. Finally, $\delta(G)$ will denote the minimum degree among the vertices of G .

A graph G is called *domatically critical*, if $d(G \setminus e) < d(G)$ for each edge e of G [1].

We shall say that the partition V_1, V_2, \dots, V_d of $V(G)$ possesses property (P), if it satisfies the following conditions:

- (i) V_i is an independent set for any $i \in \{1, 2, \dots, d\}$,
- (ii) the subgraph $G_{i,j}$ of G , induced by $V_i \cup V_j$, is a disjoint union of stars (K_1 is not a star) for any $i, j \in \{1, 2, \dots, d\}$, $i \neq j$.

Conjecture [4]. *Let G be a graph, $d(G) = d$ and let there exist a partition V_1, V_2, \dots, V_d of $V(G)$ satisfying (P). Then G is domatically critical.*

The conjecture is certainly true for all graphs with $d(G) = 1$ or 2. Indeed, if $d(G) = 1$ then (from (P)) G is \bar{K}_n ; if $d(G) = 2$ then (from (P)) G is a disjoint union of stars without isolated vertices. Both cases give domatically critical graphs. However this does not hold in case $d(G) \geq 3$.

Theorem 1. *For every integer $d \geq 3$ there exists a graph G with $d(G) = d$ which has the following properties:*

- (i) *there is a partition of $V(G)$ satisfying (P);*
- (ii) *G is not domatically critical.*

We shall need the following propositions.

Proposition 1 [2]. For any graph G , $d(G) \leq \delta(G) + 1$.

Proposition 2 [2]. For any graph G , $d(G + K_n) = d(G) + n$.

Proposition 3. A graph G is domatically critical with the domatic number $d(G) = d$, if and only if any maximum D -partition of G satisfies (P).

Proof. The „only if” part of the proposition follows from definitions.

To prove the sufficiency, consider any maximum D -partition R of G . Since R satisfies (P) the partition R of $G \setminus e$ is not domatic for any e of G .

Obviously $d(G \setminus e) \leq d(G)$. Assume $d(G \setminus e) = d(G)$ for some edge e of G . Then there exists a D -partition R' of $G \setminus e$ of order $d(G)$. This partition R' is a maximum D -partition of $G - a$ contradiction. Hence $d(G \setminus e) < d(G)$ for any edge e of G and the result follows.

Proposition 4. Let $G = H + K_n$. Then G is a domatically critical graph, if and only if H is one.

Proof. Obviously it is sufficient to prove the proposition in case $n = 1$: $G = H + \{v\}$.

Necessity. Assume H is not domatically critical: there exists e of H such that $d(H \setminus e) = d(H) = d(G) - 1$ (using Proposition 2). Consider a D -partition R of $H \setminus e$ of order $d(G) - 1$. Then $R^* = R \cup \{v\}$ is a D -partition of $G \setminus e$ of order $d(G) - 1$ – this contradicts the domatic criticality of G .

Sufficiency. By contradiction. Let G be not domatically critical: there exists e of G such that $d(G \setminus e) = d(G) = d(H) + 1$ (using Proposition 2).

There are two possibilities.

(a) The edge e is non-incident to v . Consider a maximum D -partition $R = \{V_1, V_2, \dots, V_{d+1}\}$ of $G \setminus e$, where $d = d(H) \geq 1$, and assume (without loss of generality) that $v \in V_1$. Then $R^* = \{V_2 \cup (V_1 \setminus \{v\}), V_3, \dots, V_{d+1}\} \neq \emptyset$ is a D -partition of $H \setminus e$ of order $d(H)$ – this is impossible, as H is domatically critical.

(b) The edge e is incident to v . Consider the partitions R, R^* constructed above. Clearly R^* is a D -partition of H of order $d(H)$. Since $V_1 \setminus \{v\} \neq \emptyset$ (v is not dominating in $G \setminus e$) and V_2 is dominating in $G \setminus e$ (V_2 exists, as $d(H) + 1 \geq 2$) the set $V_2 \cup (V_1 \setminus \{v\})$ is dependent. Hence the maximum D -partition R^* of H does not satisfy (P). By Proposition 3, H is not domatically critical. Again, we arrive at a contradiction. Thus all cases have been considered and the proof is complete.

Proof of Theorem 1. Let H be the graph in Figure 1. On the one hand, $d(H) \leq \delta(H) + 1 = 3$ (using Proposition 1). On the other hand, the sets $X = \{x_1, x_2, x_3\}$, $Y = \{y_1, y_2, y_3\}$, $Z = \{z_1, z_2, z_3\}$ form a D -partition of H . Hence $d(H) = 3$. The D -partition $\{\{x_1, y_3, z_2\}, \{x_2, y_1, y_2\}, \{x_3, z_1, z_3\}\}$ of H does not satisfy (P), as the set $\{x_2, y_1, y_2\}$ is dependent. By Proposition 3, the graph H is not domatically critical.

Now we shall prove that the graph $G = H + K_n$, $n \geq 0$ has the properties (i), (ii) of Theorem 1. Let $V(K_n) = \{v_1, v_2, \dots, v_n\}$. By Propositions 2, 4, $d(G) = d(H) + n \geq 3$ and G is not domatically critical (as H is not so). Obviously the D -partition




Figure 1

$\{X, Y, Z\}$ of H satisfies (P), therefore the D -partition $\{X, Y, Z, \{v_i\}, i = \overline{1..n}\}$ of G satisfies (P), too. This completes the proof.

A graph G is called *domatically cocritical*, if for every pair of its non-adjacent vertices u, v the inequality $d(G \cup (u, v)) > d(G)$ holds.

Problem [5]. Does there exist a domatically cocritical graph G whose complement \bar{G} has more than $p(G) - d(G)$ edges?

The answer is affirmative.

Theorem 2. For every positive integer k there exists a domatically cocritical graph G for which

$$q(\bar{G}) = k + p(G) - d(G).$$

Proof. Consider the graph G_k whose complement \bar{G}_k is shown in Figure 2.




Figure 2

Clearly $p(G_k) = 6k + 3$ and $q(\bar{G}_k) = 4k + 2$. Each dominating set of G_k contains at least two vertices. Hence $d(G_k) \leq \lceil p(G_k)/2 \rceil = 3k + 1$. Let $I = \{r + 6t \mid r = \overline{1..3}, t = \overline{0..k-1}\}$. The sets $\{i, i + 3\}$, $\{6k + 1, 6k + 2, 6k + 3\}$ form a D -partition of G_k with $3k + 1$ classes, therefore $d(G_k) = 3k + 1$.

It is not difficult to see that $d(G_k \cup e) > d(G_k)$ for each edge e of \bar{G}_k . Thus the graph G_k is domatically cocritical and $q(\bar{G}_k) - p(G_k) + d(G_k) = (4k + 2) - (6k + 3) + (3k + 1) = k$, as required.

References

- [1] *Cockayne E. J.*: Domination of undirected graphs — a survey. *Lecture Notes Math.*, 1978, Vol. 642, 141–147.
- [2] *Cockayne E. J., Hedetniemi, S. T.*: Toward a theory of domination in graphs. *Networks*, 1977, Vol. 7, 247–261.
- [3] *Harary F.*: *Graph Theory*, Addison-Wesley, Reading, Massachusetts, 1969.
- [4] *Zelinka B.*: Domatically critical graphs. *Czechoslovak Math. Journal*, 1980, Vol. 30, No 3, 486–489.
- [5] *Zelinka B.*: Domatically cocritical graphs. *Časopis pro pěst. mat.*, 1983, Vol. 108, No 1, 82–88.

Authors' address: Mayakovskogo, 152, kv. 56, 220028 Minsk, USSR.