

Ján Jakubík

Lattice ordered groups with unique addition must be archimedean

Czechoslovak Mathematical Journal, Vol. 41 (1991), No. 3, 559–563

Persistent URL: <http://dml.cz/dmlcz/102489>

Terms of use:

© Institute of Mathematics AS CR, 1991

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

LATTICE ORDERED GROUPS WITH UNIQUE ADDITION
MUST BE ARCHIMEDEAN

JÁN JAKUBÍK, Košice

(Received November 28, 1990)

In the present paper we will give an affirmative solution to a question proposed in a recent paper by Conrad and Darnel.

The linearly ordered groups with unique addition were identified by Ohkuma in [4]. Lattice ordered groups with unique addition were studied by Conrad and Darnel [1] and by the author [2]. The analogous notion for cyclically ordered groups was investigated in [3].

In [1], p. 19 it is remarked that every example thus far of a lattice ordered group with unique addition is an archimedean lattice ordered group, and that it is an open question whether or not a lattice ordered group with unique addition must be archimedean. The same question was proposed in [5], p. 266 (Problem 16).

It will be shown below that the answer to this question is 'YES'.

1. PRELIMINARIES

We recall the following definition (cf. [1], [2]):

A lattice ordered group $G_1 = (G; \leq, +_1)$ is said to have a *unique addition* if, whenever $G_2 = (G; \leq, +_2)$ is a lattice ordered group such that the neutral element of the group $(G; +_1)$ is the same as the neutral element of the group $(G; +_2)$, then the operation $+_1$ coincides with the operation $+_2$.

1.1. Lemma. (Cf. [1], Corollary 2.) *If $G_1 = (G; \leq, +_1)$ is a lattice ordered group with unique addition, then each automorphism τ of the lattice $(G; \leq)$ with $0\tau = 0$ is a group automorphism.*

The following assertion is easy to verify (cf. also [1]).

1.2. Lemma. *Let $G_1 = (G; \leq, +_1)$ be a lattice ordered group with unique addition. Then the group $(G; +_1)$ is abelian.*

2. AUTOMORPHISMS OF THE LATTICE $(G; \leq)$

Let $G_1 = (G; \leq, +)$ be a lattice ordered group. As usual, we put

$$G^+ = \{g \in G: g \geq 0\}.$$

In this section some auxiliary results on the automorphisms of the lattices $(G; \leq)$ and $(G^+; \leq)$ will be established.

If $x, y \in G$ and $x \wedge y = 0$, then the elements x and y are called disjoint.

The following lemma can be verified by a routine calculation.

2.1. Lemma. *Let x and y be disjoint elements of G . Then the element $x - y$ is the (uniquely determined) relative complement of the element 0 in the interval $[-y, x]$.*

2.2. Lemma. *Let φ be an automorphism of the lattice $(G^+; \leq)$. For each $x \in G$ put $\psi(x) = \varphi(x \vee 0) - \varphi(-(x \wedge 0))$. Then ψ is an automorphism of the lattice $(G; \leq)$ with $\psi(0) = 0$.*

Proof. Since $\varphi(0) = 0$, the relation $\psi(0) = 0$ obviously holds.

Let $x, y \in G, x \leq y$. Then we have $x \vee 0 \leq y \vee 0$, whence $\varphi(x \vee 0) \leq \varphi(y \vee 0)$. Analogously we obtain that $-\varphi(-(x \wedge 0)) \leq -\varphi(-(y \wedge 0))$. Thus $\psi(x) \leq \psi(y)$. Hence ψ is isotone.

Let $t_1, t_2 \in G, \psi(t_1) \leq \psi(t_2)$. It is well-known that the elements $t_1 \vee 0$ and $-(t_1 \wedge 0)$ are disjoint. Since φ is an automorphism of (G^+, \leq) , the elements $\varphi(t_1 \vee 0)$ and $\varphi(-(t_1 \wedge 0))$ are disjoint as well. Hence in view of 2.1, the element $\psi(t_1)$ is a relative complement of the element 0 in the interval $[-\varphi(-(t_1 \wedge 0)), \varphi(t_1 \vee 0)]$. An analogous assertion holds for $\psi(t_2)$. Hence

$$\begin{aligned} \psi(t_1) \vee 0 &= \varphi(t_1 \vee 0), & \psi(t_1) \wedge 0 &= -\varphi(-(t_1 \wedge 0)), \\ \psi(t_2) \vee 0 &= \varphi(t_2 \vee 0), & \psi(t_2) \wedge 0 &= -\varphi(-(t_2 \wedge 0)). \end{aligned}$$

Since $\psi(t_1) \leq \psi(t_2)$ we obtain that the relations

$$\varphi(t_1 \vee 0) \leq \varphi(t_2 \vee 0), \quad -\varphi(-(t_1 \wedge 0)) \leq -\varphi(-(t_2 \wedge 0))$$

are valid. Thus $t_1 \vee 0 \leq t_2 \vee 0$ and $t_1 \wedge 0 \leq t_2 \wedge 0$. Therefore $t_1 \leq t_2$. In particular, if $\psi(t_1) = \psi(t_2)$, then $t_1 = t_2$ and hence ψ is a monomorphism.

Let $x \in G$. Denote $\varphi^{-1}(x \vee 0) = x_1, \varphi^{-1}(-(x \wedge 0)) = x_2$. The elements x_1 and x_2 are disjoint. Put $t = x_1 - x_2$. Then in view of 2.1 we have $t \vee 0 = x_1$ and $t \wedge 0 = -x_2$, whence

$$\psi(t) = \varphi(x_1) - \varphi(x_2) = (x \vee 0) + (x \wedge 0) = x.$$

Thus ψ is an epimorphism.

In the remaining part of this section we assume that $G_1 = (G; \leq, +_1)$ is an abelian lattice ordered group which fails to be archimedean. Thus there are elements a and b_1 in G such that

$$0 < na < b_1$$

is valid for each positive integer n ; this situation will be denoted by writing $a \ll b_1$. Let an element a with the described property be fixed. Denote

$$B_0 = \{b \in G: a \ll b\}.$$

Next, for each positive integer n we define by induction a subset B_n of G^+ as follows:

B_n is the set of all elements $y \in G^+$ having the property that there exists $z \in B_{n-1}$ with $z < y \vee z \leq z + a$.

Put $B = \bigcup B_n$ ($n = 0, 1, 2, \dots$). From the definition of B we immediately obtain:

2.3. Lemma. $B \neq \emptyset$, and the element 0 does not belong to B .

Let us consider the mapping $\varphi: G^+ \rightarrow G^+$ which is defined by

$$\varphi(x) = x + a \quad \text{if } x \in B, \quad \text{and}$$

$$\varphi(x) = x \quad \text{otherwise}.$$

2.4. Lemma. Let $n \in \{0, 1, 2, \dots\}$, $y \in B_n$, $y_1 \in G$, $y < y_1$. Then $y_1 \in B_n$.

Proof. We proceed by induction on n . The assertion obviously holds for $n = 0$. Let $n > 0$ and suppose that the assertion is valid for B_{n-1} . Let z be as in the definition of B_n . Put

$$z' = z + (y_1 - y).$$

Then in view of the induction assumption we have $z' \in B_{n-1}$. Next,

$$z' < y_1 \vee z' \leq z' + a,$$

whence $y_1 \in B_n$.

2.5. Corollary. Let $y \in B$, $y_1 \in G$, $y < y_1$. Then $y_1 \in B$.

2.6. Lemma. The mapping φ is isotone.

Proof. Let $y_1, y_2 \in G^+$, $y_1 \leq y_2$. If either both y_1 and y_2 belong to B or both y_1 and y_2 belong to $G^+ \setminus B$, then clearly $\varphi(y_1) \leq \varphi(y_2)$. If $y_1 \in B$, then in view of 2.5 the relation $y_2 \in B$ is valid. Thus it suffices to consider the case $y_1 \notin B$ and $y_2 \in B$; hence

$$\varphi(y_1) = y_1 < y_2 < y_2 + a = \varphi(y_2).$$

2.7. Lemma. Let $n \in \{0, 1, 2, \dots\}$, $y \in B_n$. Then $y - a \in B_n$.

Proof. The assertion obviously holds for $n = 0$. Let $n > 0$ and assume that the assertion is valid for B_{n-1} . Let z be as in the definition of B_n . Put $z' = z - a$ and $y' = y - a$. Then $z' \in B_{n-1}$ and

$$z' < y' \vee z' \leq z' + a,$$

whence $y' \in B_n$.

2.8. Corollary. Let $y \in B$. Then $y - a \in B$.

2.9. Lemma. The mapping φ is a surjective monomorphism.

Proof. Let $y \in G^+$. If $y \notin B$, then $\varphi(y) = y$. Next, suppose that y belongs to B . Then in view of 2.8 the relation $y - a \in B$ is valid. Thus $\varphi(y - a) = y$. Hence φ is surjective.

Let $y_1, y_2 \in G^+, y_1 \neq y_2$. If either $y_1, y_2 \in B$ or $y_1, y_2 \in G^+ \setminus B$, then we obviously have $\varphi(y_1) \neq \varphi(y_2)$. Assume that $y_1 \in B$ and $y_2 \notin B$. Then in view of 2.5, $\varphi(y_1) = y_1 + a \in B$. Next, $\varphi(y_2) = y_2 \notin B$. Thus $\varphi(y_1) \neq \varphi(y_2)$.

2.10. Lemma. *Let $y_1, y_2 \in G^+, \varphi(y_1) < \varphi(y_2)$. Then $y_1 < y_2$.*

Proof. If either $y_1, y_2 \in B$ or $y_1, y_2 \in G^+ \setminus B$, then $y_1 < y_2$. Next, 2.5 and 2.7 yield that for each $t \in G^+$ the relation

$$t \in B \Leftrightarrow \varphi(t) \in B$$

is valid. Hence in view of 2.5 it suffices to consider the case $y_1 \notin B$ and $y_2 \in B$. Thus there exists $n \in \{0, 1, 2, \dots\}$ such that $y_2 \in B_n$ and

$$\varphi(y_1) = y_1, \quad \varphi(y_2) = y_2 + a.$$

Therefore $y_1 < y_2 + a$. Clearly $y_1 \neq y_2$.

Assume that $y_1 \not\leq y_2$. Then

$$y_2 < y_1 \vee y_2 \leq y_2 + a.$$

Therefore $y_1 \in B_{n+1} \subseteq B$, which is a contradiction.

2.11. Lemma. *The mapping φ is an automorphism of the lattice $(G^+; \leq)$.*

Proof. This is a consequence of 2.6, 2.9 and 2.10.

3. UNIQUE ADDITION

3.1. Proposition. *Let $G_1 = (G; \leq, +)$ be an abelian lattice ordered group. Assume that G_1 fails to be archimedean. Then there is an automorphism ψ of the lattice $(G; \leq)$ with $\psi(0) = 0$ such that ψ is not a group automorphism.*

Proof. Let φ be as in Section 2. By means of φ we construct the automorphism ψ as in 2.2. In view of this construction,

$$\psi(x) = \varphi(x) \quad \text{for each } x \in G^+.$$

Next, let the element $a \in G$ be as in Section 2. In view of 2.3, there exists $b \in B$. Then according to 2.5 we have $2b \in B$ and thus

$$\psi(2b) = 2b + a \neq 2b + 2a = 2\psi(b);$$

hence ψ is not a group automorphism.

3.2. Theorem. *Let $G_1 = (G; \leq, +)$ be a lattice ordered group with unique addition. Then G_1 is archimedean.*

Proof. According to 1.2, G_1 is abelian. By way of contradiction, assume that G_1 fails to be archimedean. Let ψ be as in 3.1. By applying 1.1 we arrive at a contradiction.

References

- [1] *r'. Conrad, M. Darnel*: l-groups with unique addition. Algebra and Order. Proc. First. Int. Symp. Ordered Algebraic Structures, Luminy-Marseille 1984, Helderman Verlag, Berlin 1986, 15–27.
- [2] *J. Jakubik*: On lattice ordered groups having a unique addition. Czechoslov. Math. J. 40, 1990, 311–314.
- [3] *J. Jakubik*: Cyclically ordered groups with unique addition. Czechoslov. Math. J. 40, 1990, 534–538.
- [4] *T. Ohkuma*: Sur quelques ensembles ordonnés linéairement. Fund. Math. 43, 1955, 326–337.
- [5] Ordered algebraic structures. Proc. Conf. Curaçao, 1989. Edited by J. Martinez. Kluwer Acad. Publ., Dordrecht–Boston–London 1989.

Author's address: 040 01 Košice, Grešákova 6, Czechoslovakia (Matematický ústav SAV, dislokované pracovisko).