

Jindřich Nečas; Miloš Štípl

A paradox in the theory of linear elasticity

Aplikace matematiky, Vol. 21 (1976), No. 6, 431--433

Persistent URL: <http://dml.cz/dmlcz/103667>

Terms of use:

© Institute of Mathematics AS CR, 1976

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A PARADOX IN THE THEORY OF LINEAR ELASTICITY

JINDŘICH NEČAS, MILOŠ ŠTÍPL

(Received November 26, 1975)

Let $\Omega = \{x \in E_3; \|x\| < 1\}$. Let $\mathcal{D}(\Omega)$ be the class of real functions, each of which is infinitely differentiable and has its support in Ω . Let $W^{1,2}(\Omega)$, $W_0^{1,2}(\Omega)$ be the usual Sobolev spaces. Let us define C_{ijkl} , $i, j, k, l = 1, 2, 3$ (the tensor of the elastic coefficients) in Ω as

$$C_{ijkl}(x) = \frac{1}{2}(\delta_{ik}\delta_{lj} + \delta_{il}\delta_{jk}) + \delta_{ij}\delta_{kl} + \frac{3}{\|x\|^2}(\delta_{ij}x_kx_l + \delta_{kl}x_ix_j) + \frac{9}{\|x\|^4}x_ix_jx_kx_l, \quad \|x\| \neq 0,$$

where δ_{ij} is the Kronecker symbol delta. Let us denote the strain tensor by $e_{kl} = \frac{1}{2}(\partial u_k/\partial x_l + \partial u_l/\partial x_k)$ (where u is the displacement vector), $k, l = 1, 2, 3$.

Let $u_0 \in [W^{1,2}(\Omega)]^3$. We say that the vector function $u \in [W^{1,2}(\Omega)]^3$ is a generalized solution of the second problem of the mathematical theory of elasticity in Ω with the boundary condition $u = u_0$ on $\partial\Omega$, if the following conditions are fulfilled:

(i)
$$\int_{\Omega} C_{ijkl} \frac{\partial v_i}{\partial x_j} e_{kl} dx = 0 \quad \text{for every } v \in [W_0^{1,2}(\Omega)]^3$$

(we neglect body forces),

(ii)
$$u - u_0 \in [W_0^{1,2}(\Omega)]^3.$$

Put

$$\alpha = \frac{3(1 - \sqrt{17})}{2\sqrt{17}}.$$

Theorem. *The displacement vector $u(x) = x\|x\|^\alpha = (x_1\|x\|^\alpha, x_2\|x\|^\alpha, x_3\|x\|^\alpha)$ is the generalized solution of the second problem of the mathematical theory of elasticity in Ω with the boundary condition $u(x) = x$ on $\partial\Omega$.*

Proof. We shall prove the relation (i). The other one is obvious. If $\|x\| \neq 0$, then

(1)
$$\frac{\partial}{\partial x_j}(C_{ijkl}e_{kl}) = 0, \quad i = 1, 2, 3.$$

Let φ be an arbitrary function from $[\mathcal{D}(\Omega)]^3$. Let $\psi \in \mathcal{D}(\Omega)$ be such a function that $\psi(x) = 1$ when $\|x\| < \frac{1}{2}$. We write

$$\psi_\varepsilon(x) = \psi(x/\varepsilon), \quad \varphi_\varepsilon(x) = \varphi(x)(1 - \psi_\varepsilon(x)), \quad \varepsilon \in (0, 1).$$

Then

$$\int_{\Omega} C_{ijkl} \frac{\partial \varphi_i}{\partial x_j} e_{kl} \, dx = \int_{\Omega} C_{ijkl} \frac{\partial \varphi_{\varepsilon i}}{\partial x_j} e_{kl} \, dx + \int_{\|x\| < \varepsilon} C_{ijkl} \frac{\partial (\varphi_i \psi_\varepsilon)}{\partial x_j} e_{kl} \, dx.$$

The first integral on the right hand side is, according to Green's theorem and to (1), equal to zero. Because C_{ijkl} and $\partial(\varphi_i \psi_\varepsilon)/\partial x_j$ are bounded and $|e_{kl}| \leq \|x\|^\alpha$, it is

$$\left| \int_{\Omega} C_{ijkl} \frac{\partial \varphi_i}{\partial x_j} e_{kl} \, dx \right| \leq C\varepsilon^{\alpha+3}, \quad C > 0.$$

Because $(\alpha + 3) > 0$, the relation (i) holds for every function $v \in [\mathcal{D}(\Omega)]^3$. The set $[\mathcal{D}(\Omega)]^3$ is dense in $[W_0^{1,2}(\Omega)]^3$, hence (i) holds.

The uniqueness of the solution follows from the relation

$$C_{ijkl}(x) \zeta_{ij} \zeta_{kl} \geq \zeta_{ij} \zeta_{ij} \quad \text{for every } \zeta \in E_6, \quad \zeta_{ij} = \zeta_{ji}, \quad \|x\| \neq 0.$$

From the physical point of view we may compare this deformation to an explosion. When the radius of the sphere Ω increases by an arbitrary $\varepsilon > 0$, then the points from a neighbourhood of the origin "cross the boundary of Ω " (i.e., for the boundary condition $u_0(x) = \varepsilon x$ it is $\|x + u(x)\| > 1 + \varepsilon$ in a neighbourhood of the origin). The displacement vector and the stress tensor are unbounded.

The tensor C_{ijkl} is constant on the radial lines (except for the origin) and invariant with respect to the rotation about the origin. The behaviour of the derived material is paradoxical. Let us have a constant tensor $\bar{C}_{ijkl} = C_{ijkl}(\frac{1}{2}, 0, 0)$. Consider the cube $\langle 0, 1 \rangle^3$ of derived homogeneous material. In the case of a constant hydrostatic pressure the body extends in the direction of the axis x_1 . In the case of a pure tension in the direction of the axis x_1 the body contracts.

Nonetheless, all the assumptions of the mathematical theory of the linear elasticity are satisfied (i.e., the coefficients C_{ijkl} are measurable, bounded, the form $C_{ijkl} \zeta_{ij} \zeta_{kl}$ is elliptic).

References

- [1] *E. De Giorgi*: Un esempio di estremali discontinue per un problema variazionale di tipo ellittico, *Boll. U. M. I.*, Vol. I., 1968, 135–137.
- [2] *J. Nečas*: Les méthodes directes en théorie des équations elliptiques, Praha 1967.
- [3] *L. F. Nye*: Physical properties of crystals, Oxford 1957.

Souhrn

PARADOX V TEORII LINEÁRNÍ PRUŽNOSTI

JINDŘICH NEČAS, MILOŠ ŠTÍPL

Uvažujme systém parciálních diferenciálních rovnic lineární pružnosti. Ukážeme, že řešení tohoto systému s omezenou okrajovou podmínkou není (obecně) omezené (tj. nejsou omezené složky vektoru posunutí). Tento příklad je modifikací příkladu z článku E. De Giorgiho [1].

Author's addresses: Doc. Dr. Jindřich Nečas, DrSc., Matematický ústav ČSAV, Žitná 25, 115 67 Praha 1; Miloš Štípl, Za Hládkovem 7, 169 00 Praha 6.