

Aplikace matematiky

Jiří Rohn

Nonsingularity and P -matrices.

Aplikace matematiky, Vol. 35 (1990), No. 3, 215--219

Persistent URL: <http://dml.cz/dmlcz/104405>

Terms of use:

© Institute of Mathematics AS CR, 1990

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

NONSINGULARITY AND P -MATRICES

Jiří ROHN

(Received February 17, 1989)

Summary. New proofs of two previously published theorems relating nonsingularity of interval matrices to P -matrices are given.

Keywords: Nonsingular matrices, P -matrices.

AMS Classification: 15A03, 15A57, 65G10.

In [5] we proved, in a broader frame of the problem of solving linear interval systems, two theorems relating nonsingularity of interval matrices to P -matrices (Theorems 1 and 2 below). It is the purpose of this paper to give alternative proofs of them, from which it can be perhaps better seen how nonsingularity is intertwined with P -property. We also include some consequences implied by the properties of P -matrices.

We begin with this simple auxiliary result:

Lemma. *Let A be a nonsingular $n \times n$ matrix and let B be an $n \times n$ matrix whose rows, except the j -th, are zero. Let $1 + (BA^{-1})_{jj} \leq 0$. Then there exists a $t \in (0, 1]$ such that $A + tB$ is singular.*

Proof. Consider the function φ of one real variable defined by $\varphi(\tau) = 1 + \tau(BA^{-1})_{jj}$. Since $\varphi(0) > 0$ and $\varphi(1) \leq 0$, there exists a $t \in (0, 1]$ such that $\varphi(t) = 0$. Then the matrix $A + tB = (E + tBA^{-1})A$ is singular since $\det(E + tBA^{-1}) = 1 + t(BA^{-1})_{jj} = 0$. ■

Let A^-, A^+ be two $n \times n$ matrices, $A^- \leq A^+$ (the inequality to be understood componentwise). The set of matrices

$$A^I = \{A; A^- \leq A \leq A^+\}$$

is called an interval matrix; we say that A^I is nonsingular (in [5]: regular) if each $A \in A^I$ is nonsingular. A square matrix A is said to be a P -matrix [1] if all its principal minors are positive.

First, we have this result:

Theorem 1. *Let A^I be nonsingular. Then for each $A_1, A_2 \in A^I$, both $A_1A_2^{-1}$ and $A_1^{-1}A_2$ are P -matrices.*

Proof. The proof consists of several steps. Let $A_1, A_2 \in A^I$.

(1) We shall prove that all *leading* principal minors m_1, \dots, m_n of $A_1A_2^{-1}$ are positive. Put $D = A_1 - A_2$ so that $A_1A_2^{-1} = E + DA_2^{-1}$, and denote by D_j ($j = 1, \dots, n$) the matrix whose first j rows are identical with those of D and the remaining ones are zero. Then

$$m_j = \det(E + D_jA_2^{-1})$$

holds for $j = 1, \dots, n$. We shall prove by induction that $m_j > 0$ for each j :

(1.1) Case $j = 1$. Since $m_1 = \det(E + D_1A_2^{-1}) = 1 + (D_1A_2^{-1})_{11}$, the above lemma implies $m_1 > 0$, for otherwise the matrix $A_2 + tD_1$ would be singular for some $t \in (0, 1]$ but $A_2 + tD_1 \in A^I$, which is a contradiction.

(1.2) Case $j > 1$. Assume that $m_{j-1} > 0$ and consider the matrix

$$(E + D_jA_2^{-1})(E + D_{j-1}A_2^{-1})^{-1} = E + (D_j - D_{j-1})(A_2 + D_{j-1})^{-1}.$$

Taking determinants on both sides we obtain

$$\frac{m_j}{m_{j-1}} = 1 + [(D_j - D_{j-1})(A_2 + D_{j-1})^{-1}]_{jj}.$$

If the right-hand side were nonpositive, then, according to the lemma, $A_2 + D_{j-1} + t(D_j - D_{j-1})$ would be singular for some $t \in (0, 1]$, which is a contradiction since it is a matrix from A^I . Hence

$$\frac{m_j}{m_{j-1}} > 0$$

holds, which in conjunction with the induction hypothesis gives that $m_j > 0$, which concludes the inductive proof.

(2) Second we shall prove that each principal minor of $A_1A_2^{-1}$ is positive. Consider a principal minor formed from the rows and columns with indices k_1, \dots, k_r , $1 \leq r \leq n$. Let R be any permutation matrix with $R_{k_jj} = 1$ ($j = 1, \dots, r$). Then the above minor is equal to the r -th leading principal minor of $R^T A_1 A_2^{-1} R = (R^T A_1 R) \cdot (R^T A_2 R)^{-1}$. Since the interval matrix $\{R^T A R; A \in A^I\}$ is nonsingular, all leading principal minors of $(R^T A_1 R)(R^T A_2 R)^{-1}$ are positive due to (1).

(3) To prove that $A_1^{-1}A_2$ is also a P -matrix, consider the transpose interval matrix $(A^I)^T = \{A^T; A \in A^I\}$. According to part (2), its nonsingularity implies that $(A_2^T)(A_1^T)^{-1} = (A_1^{-1}A_2)^T$ is a P -matrix, hence so is $A_1^{-1}A_2$. This completes the proof. ■

We shall now show that the result can be in a certain sense reversed, so that the P -property of a finite number of matrices of the form $A_1^{-1}A_2$ will imply nonsingularity of A^I . To this end, let us denote

$$A_c = \frac{1}{2}(A^- + A^+),$$

$$\Delta = \frac{1}{2}(A^+ - A^-),$$

then $A^- = A_c - \Delta$, $A^+ = A_c + \Delta$, $\Delta \geq 0$. A diagonal matrix S satisfying $|S_{ii}| = 1$ for each i is called a signature matrix, so that there are 2^n signature matrices of size n .

Theorem 2. *An interval matrix A^I is nonsingular if and only if for each signature matrix S , $A_c - S\Delta$ is nonsingular and $(A_c - S\Delta)^{-1}(A_c + S\Delta)$ is a P -matrix.*

Proof. The “only if” part being an obvious consequence of Theorem 1, we must prove the “if” part only. This will be done if we prove that for each $A \in A^I$ and each $b \in R^n$, the system of linear equations

$$Ax = b$$

has a solution, which, according to a theorem proved in [6], is equivalent to the fact that for each signature matrix S , the system of linear inequalities

$$(*) \quad SAx \geq Sb$$

has a solution. To show this, consider the linear complementarity problem

$$x_1 = (A_c - S\Delta)^{-1}(A_c + S\Delta)x_2 + (A_c - S\Delta)^{-1}b,$$

$$x_1^T x_2 = 0,$$

$$x_1 \geq 0, \quad x_2 \geq 0.$$

Since $(A_c - S\Delta)^{-1}(A_c + S\Delta)$ is a P -matrix by the assumption, this problem has a solution x_1, x_2 , as proved in [7]. Then

$$A_c(x_1 - x_2) - S\Delta(x_1 + x_2) = b$$

and for each $A \in A^I$ we have

$$SA(x_1 - x_2) = SA_c(x_1 - x_2) + S(A - A_c)(x_1 - x_2) \geq$$

$$\geq SA_c(x_1 - x_2) - \Delta(x_1 + x_2) = Sb,$$

hence $(*)$ has a solution, which by virtue of the above-quoted theorem proves that A^I is nonsingular. ■

It is worth noting that the matrices $(A_c - S\Delta)^{-1}(A_c + S\Delta)$ cannot be replaced by matrices of the type $(A_c - S\Delta)(A_c + S\Delta)^{-1}$ in the formulation of Theorem 2:

Example 1 (communicated to the author by M. Baumann). Let

$$A^- = \begin{pmatrix} 3 & -1 \\ 1 & 3 \end{pmatrix}, \quad A^+ = \begin{pmatrix} 7 & 3 \\ 5 & 7 \end{pmatrix}.$$

Then $(A_c - S\Delta)(A_c + S\Delta)^{-1}$ is a P -matrix for each signature matrix S , but A^I contains the singular matrix

$$\begin{pmatrix} 3 & 3 \\ 3 & 3 \end{pmatrix}.$$

Since each positive definite (not necessarily symmetric) matrix is a P -matrix [1], we obtain a consequence:

Corollary 1. *For each signature matrix S , let $A_c - S\Delta$ be nonsingular and $(A_c - S\Delta)^{-1}(A_c + S\Delta)$ positive definite. Then A^I is nonsingular.*

The converse implication is, however, not true:

Example 2. Let

$$A^- = \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix}, \quad A^+ = \begin{pmatrix} 3 & 1 \\ 1 & 0 \end{pmatrix}.$$

Then A^I is obviously nonsingular, but none of the matrices $(A_c - S\Delta)^{-1}(A_c + S\Delta)$ is positive definite.

Finally, using the well-known properties of P -matrices, we may draw some consequences regarding nonsingular interval matrices:

Corollary 2. *Let A^I be nonsingular. Then for each $A_1, A_2 \in A^I$ we have:*

- (i) *each diagonal element of both $A_1^{-1}A_2$ and $A_1A_2^{-1}$ is positive,*
- (ii) *for each signature matrix S there exist x_1, x_2 such that $A_1x_1 = A_2x_2$, $Sx_1 > 0$, $Sx_2 > 0$,*
- (iii) *for each signature matrix S there exist x_1, x_2 such that $A_1^{-1}x_1 = A_2^{-1}x_2$, $Sx_1 > 0$, $Sx_2 > 0$,*
- (iv) *if $A_1x_1 = A_2x_2$ for some $x_1 \neq 0$, $x_2 \neq 0$, then $(x_1)_i(x_2)_i > 0$ for some $i \in \{1, \dots, n\}$,*
- (v) *if $A_1^{-1}x_1 = A_2^{-1}x_2$ for some $x_1 \neq 0$, $x_2 \neq 0$, then $(x_1)_i(x_2)_i > 0$ for some $i \in \{1, \dots, n\}$.*

Proof. (i) follows from the fact that each diagonal element (i.e., first order minor) of a P -matrix is positive. (ii) Let S be a signature matrix. Then the interval matrix $\{AS; A \in A^I\}$ is nonsingular, hence $(A_1S)^{-1}(A_2S) = SA_1^{-1}A_2S$ is a P -matrix; then, as proved by Gale and Nikaido [3], there exists a $y_2 > 0$ such that $y_1 = SA_1^{-1}A_2Sy_2 > 0$. Setting $x_1 = Sy_1$, $x_2 = Sy_2$, we obtain vectors with the properties stated. (iii) is proved in a similar manner as (ii). (iv) If $A_1x_1 = A_2x_2$, then $x_1 = A_1^{-1}A_2x_2$ and since $A_1^{-1}A_2$ is a P -matrix, the result follows from the characterization by Fiedler and Pták [2]. (v) follows in a similar way from the fact that $A_1A_2^{-1}$ is a P -matrix. ■

The necessary and sufficient nonsingularity conditions given in Theorem 2 are generally very difficult to verify. This fact becomes more understandable in the light of the recent result by Poljak and Rohn [4] stating that testing nonsingularity of an interval matrix is an NP-complete problem.

References

- [1] *M. Fiedler*: Special Matrices and Their Use in Numerical Mathematics (Czech). SNTL, Prague 1981.
- [2] *M. Fiedler, V. Pták*: On Matrices with Non-Positive Off-Diagonal Elements and Positive Principal Minors. Czechoslovak Math. J. 12 (1962), 382–400.
- [3] *D. Gale, H. Nikaido*: The Jacobian Matrix and Global Univalence of Mappings. Math. Annalen 159 (1965), 81–93.
- [4] *S. Poljak, J. Rohn*: Radius of Nonsingularity. KAM Series 88–117, Charles University, Prague 1988.
- [5] *J. Rohn*: Systems of Linear Interval Equations. Lin. Alg. and Its Appls. 126 (1989), 39–78.
- [6] *J. Rohn*: Characterization of a Linear Program in Standard Form by a Family of Linear Programs with Inequality Constraints. To appear in Ekon.-Mat. Obzor.
- [7] *H. Samelson, R. Thrall, O. Wesler*: A Partition Theorem for Euclidean n -Space. Proc. of the Amer. Math. Society 9 (1958), 805–807.

Souhrn

REGULARITA A P -MATICE

Jiří ROHN

Jsou uvedeny nové důkazy dvou dříve publikovaných vět o vztahu regularity intervalových matic k reálným P -maticím.

Резюме

РЕГУЛЯРНОСТЬ И P -МАТРИЦЫ

ЖИ́РИ РО́Н

В статье приведены новые доказательства двух ранее опубликованных теорем о взаимном отношении регулярных интервальных матриц и P -матриц.

Author's address: RNDr. Jiří Rohn, Matematicko-fyzikální fakulta UK, Malostranské nám. 25, 118 00 Praha 1.