

Ladislav Procházka

Note on quasi-isomorphism of torsion free abelian groups of finite rank
(Preliminary communication)

Commentationes Mathematicae Universitatis Carolinae, Vol. 3 (1962), No. 1, 18--19

Persistent URL: <http://dml.cz/dmlcz/104904>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1962

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

3, 1 (1962)

NOTE ON QUASI-ISOMORPHISM OF TORSION FREE ABELIAN GROUPS OF

FINITE RANK
(Preliminary communication)
Ladislav PROCHÁZKA, Praha

By a group we understand always an abelian group.

According to [1] two torsion free groups G and H of finite rank are called quasi-isomorphic (such two groups are said in [2] to be almost isomorphic) if there exists a subgroup $G' \subseteq G$ such that $G' \cong H$ and G' has finite index in G . Using this concept of quasi-isomorphism the following conjecture is formulated in [1] (see [1], p.41): If G is a torsion free group of finite rank r containing a free subgroup F of the same rank r such that for each prime p , the divisible part of the p -primary component of torsion group G/F has rank at least $r-1$, then any group H which is quasi-isomorphic to G is already isomorphic to G .

It can be proved that this hypothesis really holds; the complete proof of corresponding assertion will be presented in Czechoslovak Mathematical Journal. Let us remark only that this proof is based on some results of the paper [3] concerning invariants theory of torsion free groups of finite rank and the author's paper [4].

References

- [1] R.A. Beaumont and R.S. Pierce: Torsion free groups of rank two, *Memoirs Amer. Math. Soc.*, 1961, 38, pp.1-41
- [2] B. Jónsson: On direct decomposition of torsion free abelian groups, *Math. Scand.*, 7, 1959, pp.361-371

- [3] А. Мальцев: Абелевы группы конечного ранга без кручения, Мат. сб. 4 (46), 1938, стр. 45-68
- [4] Л. Прохазка: О r -ранге абелевых групп без кручения конечного ранга, Чех.мат.ж., 12 (87), 1962, стр. 3-48