

Zdeněk Hedrlín

On a number of commuting transformations

Commentationes Mathematicae Universitatis Carolinae, Vol. 4 (1963), No. 3, 132--136

Persistent URL: <http://dml.cz/dmlcz/104943>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1963

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

4, 3 (1963)

ON A NUMBER OF COMMUTING TRANSFORMATIONS

Z. HEDRLÍN, Praha

The aim of this remark is to prove the following:

Theorem: Let f be a transformation of an n -point set X . Then there exist at least n different transformations of X commuting with f , that is, there exist transformations g_1, g_2, \dots, g_s , $s \geq n$, $g_i \neq g_j$ for $i \neq j$, such that

$$g_i[f(x)] = f[g_i(x)] \quad \text{for all } x \in X \text{ and all } i = 1, 2, \dots, s.$$

We use the following notation:

i denotes the identity transformation of X and we write $i = f^0$, $f^i = f \circ f^{i-1}$, and $F = \bigcup_{i=0}^{n-1} \{f^i\}$. We write $F(x) =$

$$= \bigcup_{i=0}^{n-1} \{f^i(x)\}.$$

$y \in X$ is said to be maximal according to f , or simply maximal, if $f(x) \neq y$ for every $x \in X$. If Y is a set, $|Y|$ denotes the cardinal of Y . We assume that $|X| = n$.

Lemma 1. Let g commute with f , $g(x_1) = x_2$. Then

$$|F(x_1)| \geq |F(x_2)|.$$

Proof. We have $f^i(x_2) = f^i[g(x_1)] = g[f^i(x_1)]$. Hence, if

$$f^i(x_2) \neq f^j(x_2), \text{ then also } f^i(x_1) \neq f^j(x_1).$$

Lemma 2. Let $F(x_1) \cap F(x_2) \neq \emptyset$, $F(x_2) \cap F(x_3) \neq \emptyset$. Then $F(x_1) \cap F(x_3) \neq \emptyset$.

Proof. We have $f(x_1) = f(x_2)$, $f(x_2) = f(x_3)$, for some i, j, k, l . Hence, $f(x_1) = f(x_3)$, and the lemma is proved.

$x_1, x_2 \in X$ are said to belong to the same component according to f , or simply to the same component, if $F(x_1) \cap F(x_2) \neq \emptyset$. By Lemma 2; two components are either equal or disjoint.

Lemma 3. Let there exist only one component and only one maximal element according to f . Then $F(y) = X$, where y is the maximal element.

Proof. Let $x_0 \notin F(y)$. As $x_0 \neq y$, x_0 is not maximal, there exists x_1 such that $f(x_1) = x_0$. Evidently, $x_1 \notin F(y)$. Continuing this process we get a sequence x_m . As $|X| = n$, there exists only finite number of different x_i . Therefore there must exist x_j and natural k such that $f(x_j) = x_j$, $x_j \notin F(y)$. Hence, $F(x_j) \cap F(y) = \emptyset$, and x_j and y belong to different components.

Lemma 4. Let there exist only one component, and no maximal element according to f . Then $F(y) = X$ for all $y \in X$. The proof is evident.

Now, we are going to prove the theorem by induction. If $n = 1$, then the theorem is evidently true. Let $n > 1$. We assume that the theorem is proved for all $m \leq n - 1$. We divide the proof in three sections according to the properties of f .

- (a) There exist more than one component.
- (b) There exists only one component containing at most one

maximal element.

(c) There exists only one component containing more than one maximal element.

(a) We denote the components Y_1, Y_2, \dots, Y_k . We may assume that $|Y_i| \geq |Y_{i+1}|$, $i = 1, 2, \dots, k-1$. Evidently, every Y_i is fixed under f , that is $f(Y_i) = \bigcup_{y_i \in Y_i} \{y_i\} \subset Y_i$. If we denote by $f|_{Y_i}$, as usual, the restriction of f onto Y_i , then $f|_{Y_i}$ is a transformation of Y_i . By assumption, there exist at least $|Y_i|$ different transformations of Y_i commuting with $f|_{Y_i}$. We denote this set by F_i . Let G be a system of transformations of X such that

$$g \in G \text{ if and only if } g|_{Y_i} \in F_i \text{ for each } i = 1, 2, \dots, k.$$

We have

$$|G| = \prod_{i=1}^k |F_i| \geq \prod_{i=1}^k |Y_i|.$$

If no $|Y_i| = 1$, then the theorem is true, as every transformation in G commutes with f .

Let $|Y_i| = 1$ for $i = 1, 2, \dots, r$, that is $Y_i = \{y_i\}$. For each $i = 1, 2, \dots, r$, we define a constant transformation $h_i(x) = y_i$ for every $x \in X$. Evidently, all h_i , $i = 1, 2, \dots, r$, commute with f . Let us denote $G' = G \cup \left(\bigcup_{i=1}^r \{h_i\} \right)$. We get

$$|G'| \geq \prod_{i=r+1}^k |Y_i| + r.$$

Evidently, $\sum_{i=r+1}^k |Y_i| = n - r$, and every $|Y_i| \geq 2$, $i = r+1, r+2, \dots, k$. Hence, $|G'| \geq n$, and the case (a) is proved.

(b) If there exists only one component with at most one

element y , then, by lemma 3 and 4, the points $y, f(y), f^2(y), \dots, f^{n-1}(y)$ are different. That proves the case (b).

(c) Let y, y_1, y_2, \dots, y_t be maximal elements, $|F(y)| \leq |F(y_i)|$ for $i = 1, 2, \dots, t$. We denote $X' = X \setminus \{y\}$, $f' = f|X'$. Evidently, f' is a transformation of X' , $|X'| = n - 1$. Hence, there exist different transformations g'_1, g'_2, \dots, g'_s , $s \geq n - 1$, such that every g'_i commutes with f' . We may assume that $g'_i, i = 1, 2, \dots, s$, are all transformations of X' commuting with f' . We are going to prove that every g'_i can be commutatively extended to X , that is, for each $g'_i, i = 1, 2, \dots, s$, there exists a transformation g_i of X such that $g_i|X' = g'_i$, and g_i commutes with f . By assumption, y is a maximal element and therefore

$$|F[f(y)]| = |F(y)| - 1 = |F'[f(y)]|,$$

where F' denotes the set of transformations of X' belonging to f' . As $|F(y)| \leq |F(y_i)|$, by lemma 1, $g'_i[f(y)]$ is not maximal element according to f .

Thus, there exists at least one element x'_i such that

$$f(x'_i) = g'_i[f(y)].$$

If we define $g_i|X' = g'_i$, $g_i(y) = x'_i$, then g_i is the required commutative extension of g'_i .

It remains only to prove the existence of g'_i which can be extended in two different ways. To prove it we show that under assumptions of (c), there exists a natural k such that

$$f^k(y) = f(z), \quad z \neq f^k(y).$$

Let $y_1 \neq y$, y_1 be maximal. We have $F(y) \cap F(y_1) \neq \emptyset$, $y_1 \notin F(y)$, $y \notin F(y_1)$. Hence, there exists natural

k such that $f^k(y) \in F(y_1)$, $f^{k-1}(y) \text{ non } \in F(y_1)$. There exists an integer m such that $f^m(y_1) = f^k(y)$. Put $z = f^{m-1}(y_1)$. Evidently, $z \neq f^{k-1}(y)$, as $f^{k-1}(y) \text{ non } \in F(y_1)$.

Now, $f^{k-1}|X'$ commutes with f' and can be extended in two different ways. The proof is finished.