

Roman Frič

A note on Fréchet spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 13 (1972), No. 3, 411--418

Persistent URL: <http://dml.cz/dmlcz/105429>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1972

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A NOTE ON FRÉCHET SPACES ¹⁾

R. FRÍČ, Žilina

Recall that a Fréchet space (L, λ) is a T_1 topological space such that for every subset A we have $\lambda A = \{x \mid x = \lim x_m, x_m \in A\}$, i.e. λA is the set of all limit points of sequences of points of A ; the space (L, λ) is said to be sequentially regular if for every sequence $\langle x_m \rangle$ of points of L and every point x such that $x \in L - \lambda U(x_m)$ there is a continuous function f on (L, λ) , $0 \leq f(x) \leq 1$, and a subsequence $\langle m_i \rangle$ of $\langle m \rangle$ such that $f(x) = 0$, $f[U(x_{m_i})] = 1$ (cf. [3]).

Following [5] a T_1 topological space (L, λ) is called κ_0 -regular if for every countable subset A and every point x such that $x \in L - \lambda A$ there is a continuous function f on (L, λ) , $0 \leq f(x) \leq 1$ such that $f(x) = 0$, $f[A] = 1$. It can be readily seen that every κ_0 -regular Fréchet space is sequentially regular. J. Novák asked in [5] whether every sequentially regular Fréchet space is κ_0 -regular.

1) The article is a part of [1].

The main purpose of the present paper is to show that the answer is no. The space \bar{A}_∞ constructed by F.B. Jones in [2] ²⁾ (as a Moore space which is not completely regular) is a counter-example. We also give a necessary and sufficient condition for a Fréchet sequentially regular space to be \mathcal{K}_0 -regular and two sufficient conditions for an \mathcal{K}_0 -regular Fréchet space to be completely regular.

Example. Let L be the subset of all points (x, y) of the Euclidean plane $\mathbb{R} \times \mathbb{R}$ such that $y \geq 0$ provided with the following refinement of the product topology: for $\kappa > 0$, the sets

$$V^\kappa(x, 0) = \{(x, 0)\} \cup \{(u, v) \mid (u, v) \in L, (u-x)^2 + (v-\kappa)^2 < \kappa^2\}$$

are also neighbourhoods of the point $(x, 0)$ (Niemytzky space).

Denote by \mathcal{A} the just described topology. Clearly, (L, \mathcal{A}) satisfies the first axiom of countability and hence it is Fréchet. The subspace $(D, \mathcal{A}/D)$ of (L, \mathcal{A}) where $D = \{(x, 0) \mid x \in \mathbb{R}\}$, is discrete. The space (L, \mathcal{A}) is completely regular and hence sequentially regular. The set D is the union of two disjoint uncountable sets, denote them by A and by B , such that if U is an open set containing uncountably many points of one of them, then $\mathcal{A}U$ contains uncountably many points of the other (for the proof see [2]).

Let $\langle (L_m, \mathcal{A}_m) \rangle_{m=1}^\infty$ be a simple sequence of disjoint copies of the space (L, \mathcal{A}) . For convenience we may

2) It is Professor J. Novák who called my attention to that article.

imagine these spaces as lying in different planes of the three-dimensional Euclidean space parallel to the plane of L . For each point set H in L and to every natural m there corresponds in a natural way the set H_m in L_m (the set H is the projection of every H_m). The symbol q denotes always a point of D .

Let $\sum_{m=1}^{\infty} (L_m, \lambda_m)$ be the topological sum of the above sequence. We modify it in the following manner:

1. If m is odd ($m = 1, 3, 5, \dots$) and q is a point of B , then we identify points q_m and q_{m+1} to $(q_m; q_{m+1})$; if m is even ($m = 2, 4, 6, \dots$) and q is a point of A , then we identify points q_m and q_{m+1} to $(q_m; q_{m+1})$ (the projection of $(q_m; q_{m+1})$ is q in this case). Let for $\kappa > 0$ the sets

$$W^{\kappa}((q_m; q_{m+1})) = \\ = \{(q_m; q_{m+1})\} \cup \{V_m^{\kappa}(q) - (q)\} \cup \{V_{m+1}^{\kappa}(q) - (q_{m+1})\}$$

be fundamental systems of neighbourhoods of these points, i.e. we take a quotient space of $\sum_{m=1}^{\infty} (L_m, \lambda_m)$.

2. We add one "ideal" point μ (distinct from all) to the modified $\sum_{m=1}^{\infty} (L_m, \lambda)$.

Let for $k = 1, 2, 3, \dots$, the sets

$$O_k(\mu) = \{\mu\} \cup \left\{ \bigcup_{m > k} \bigcup_{y > 0} (x_m, y_m) \right\} \cup \left\{ \bigcup_{m > k} (q_m; q_{m+1}) \right\}$$

form a fundamental system of neighbourhoods of μ .

Denote by $(L_{\infty}, \lambda_{\infty})$ this modified space (cf. [2], where $\bar{L}_{\infty} = (L_{\infty}, \lambda_{\infty})$). The space $(L_{\infty}, \lambda_{\infty})$ satisfies the first axiom of countability and hence it is

Fréchet, it is "completely regular at every point" except μ but it is not completely regular (at μ) since $\mu \in \epsilon L_\infty - \lambda_\infty A_1$, but for each continuous function f on $(L_\infty, \lambda_\infty)$ we have $f(\mu) \in \overline{f[A_1]}$ (cf. [2]).

Proposition. The Fréchet space $(L_\infty, \lambda_\infty)$ is sequentially regular but fails to be κ_0 -regular.

Proof. First prove that $(L_\infty, \lambda_\infty)$ is sequentially regular. Since $(L_\infty, \lambda_\infty)$ is "completely regular and hence sequentially regular at every point" except μ , we have to prove that if $\langle x_m \rangle$ is a sequence of points of L_∞ such that $\mu \in L_\infty - \lambda_\infty \bigcup_{m=1}^{\infty} (x_m)$, then there is a continuous function f on $(L_\infty, \lambda_\infty)$ and a subsequence $\langle x_{m_i} \rangle$ of $\langle x_m \rangle$ such that

$$f(\mu) = 0, \quad f(x_{m_i}) = 1, \quad i = 1, 2, 3, \dots$$

Since there is a natural κ_0 such that $x_m \in L_\infty - \theta_{\kappa_0}(\mu)$ for all m , we always can and do select a subsequence $\langle x'_{m_i} \rangle$ of $\langle x_m \rangle$ such that

a) $\langle x'_{m_i} \rangle$ is a constant sequence or the projection of no x'_{m_i} lies in $D \subset L$. In this case the construction of f and the subsequence $\langle x_{m_i} \rangle$ of $\langle x'_{m_i} \rangle$ and hence of $\langle x_m \rangle$ is easy and is omitted.

b) If $(x'_i, 0) \in D \subset L$ is the projection of x'_{m_i} , i.e. x'_{m_i} is either of the form of $(q_m^{(i)}; q_{m+1}^{(i)})$, $m \leq \kappa_0$, or $x'_{m_i} \in A_1$, then there is a strictly monotone, say increasing, subsequence $\langle x_i \rangle$ of the sequence $\langle x'_i \rangle$ of real numbers x'_i . Let $\langle \kappa_i \rangle$ be a sequence of positive real numbers such that

$$x_{i-1} + \kappa_{i-1} < x_i - \kappa_i < x_i + \kappa_i < x_{i+1} - \kappa_{i+1}, i = 1, 2, 3, \dots$$

Denote by $U(x_{m_i}) = (V^{n_i}(x_i, 0))_1$ if $x_{m_i} \in A_1$ and

$$U(x_{m_i}) = W^{n_i}((q_m^{(i)}; q_{m+1}^{(i)}))$$

otherwise. Now, let f be a function on $(L_\infty, \mathcal{A}_\infty)$ defined in the following manner:

$$f(x) = 1 \text{ for } x = x_{m_i};$$

$f(x) = 0$ for each x on the boundary of the neighbourhood $U(x_{m_i})$ of x_{m_i} and linear on the segment from x_{m_i} to x , $i = 1, 2, 3, \dots$;

$$f(x) = 0 \text{ for } x \in L_\infty - \bigcup_{i=1}^{\infty} U(x_{m_i}).$$

It is easy to verify that f has the desired properties. If the sequence $\langle x_i \rangle$ is decreasing, then the procedure is similar.

Secondly, denote by

$$C = \{(x, y) \mid (x, y) \in L - D; x, y \text{ rational}\}.$$

The set C_1 is countable and can be arranged into a sequence $\langle x_m \rangle$ and $\rho \in L_\infty - \bigcup_{m=1}^{\infty} (x_m)$. As

$$A_1 \subset \bigcup_{m=1}^{\infty} (x_m), \text{ we have } f(\rho) \in \overline{\bigcup_{m=1}^{\infty} (f(x_m))}$$

for each continuous function f on $(L_\infty, \mathcal{A}_\infty)$. Therefore $(L_\infty, \mathcal{A}_\infty)$ fails to be κ_0 -regular. This completes the proof.

Let (L, \mathcal{A}) be a Fréchet sequentially regular space. Recall that the completely regular modification $\tilde{\mathcal{A}}$ of \mathcal{A} is the finest of all completely regular topologies for L coarser than \mathcal{A} , the systems of continuous functions on

(L, λ) and on $(L, \tilde{\lambda})$ coincide and $\lim x_n = x$ if and only if the sequence $\langle x_n \rangle$ is eventually in every $\tilde{\lambda}$ -neighbourhood of x (see [3]). A point x_0 is called a side-point of a sequence $\langle x_n \rangle$ in $(L, \tilde{\lambda})$ if any subsequence $\langle x_{n_i} \rangle$ of $\langle x_n \rangle$ does not converge to x_0 and the sequence $\langle x_n \rangle$ is frequently in every $\tilde{\lambda}$ -neighbourhood of x_0 .

Theorem 1. A Fréchet sequentially regular space (L, λ) is x_0 -regular if and only if there is no sequence in $(L, \tilde{\lambda})$ having a side-point, where $\tilde{\lambda}$ is the completely regular modification of λ .

Proof. I. If there is a sequence $\langle x_n \rangle$ in $(L, \tilde{\lambda})$ having a side-point x_0 , then

$$x_0 \in L - \lambda U(x_n), \quad x_0 \in \tilde{\lambda} U(x_n).$$

Thus for each continuous function f on $(L, \tilde{\lambda})$ and hence, as mentioned above, on (L, λ) we have

$$f(x_0) \in \overline{U(f(x_n))}.$$

But this implies that (L, λ) cannot be x_0 -regular.

II. If (L, λ) is not x_0 -regular, then there is a sequence $\langle x_n \rangle$ of points $x_n \in L$ and a point $x_0 \in L$ such that

$$x_0 \in L - \lambda U(x_n)$$

and for each continuous function f on (L, λ) there is a subsequence $\langle n_i \rangle$ of $\langle n \rangle$ such that

$$\lim f(x_{n_i}) = f(x_0).$$

From the definition of $\tilde{\lambda}$ it follows that

$$x_0 \in \tilde{\lambda} U(x_n) ,$$

i.e. x_0 is a side-point of the sequence $\langle x_n \rangle$ in $(L, \tilde{\lambda})$.

Theorem 2. A regular separable λ_0 -regular Fréchet space (L, λ) is completely regular.

Proof. Denote by $S \subset L$ a countable set such that $G \cap S \neq \emptyset$ for each non-empty open set $G \subset L$. Let $F \subset L$ be a non-empty closed set and $x_0 \in L - F$. Then there is a neighbourhood $W(x_0)$ such that $\lambda W(x_0) \subset L - F$ and $(L - \lambda W(x_0)) \cap S \neq \emptyset$. Hence $(L - W(x_0)) \cap S \neq \emptyset$. Now, arrange the countable set $(L - W(x_0)) \cap S$, either finite or infinite, into a sequence $\langle x_n \rangle$. Evidently

$$x_0 \in (L - \lambda U(x_n)) \subset L - F .$$

Since (L, λ) is λ_0 -regular, there is a continuous function f on (L, λ) such that

$$f(x_0) = 0 , f[U(x_n)] = 1 = f[F] .$$

Corollary. A first-countable separable λ_0 -regular topological space is completely regular.

Proof. Professor J. Novák proved in [4] that every first-countable sequentially regular topological space is regular. The assertion follows at once from the foregoing Theorem 2.

R e f e r e n c e s

- [1] R. FRIČ: Sequential structures and their application to probability theory. Thesis, MÚ ČSAV, Praha, 1972.

- [2] F.B. JONES: Moore spaces and uniform spaces. Proc.Amer. Math.Soc.9(1958),483-486.
- [3] V. KOUTNÍK: On sequentially regular convergence spaces. Czechoslovak Math.J.17(1967),232-247.
- [4] J. NOVÁK: On convergence spaces and their sequential envelopes. Czechoslovak Math.J.15(1965),74-100.
- [5] J. NOVÁK: On some problems concerning the convergence spaces and groups. General Topology and its Relations to Modern Analysis and Algebra(Proc.Kanpur Topological Conf.,1968).Academia,Prague,1971, 219-229.

MÚ ČSAV v Praze
Praha 1, Žitná 25

VŠD v Žiline
Žilina, Marx-Engelsa 25

Československo

(Oblatum 20.4.1972)