

Jaromír J. Koliha

The product of relatively regular operators

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 16 (1975), No. 3, 531--539

Persistent URL: <http://dml.cz/dmlcz/105645>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1975

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## THE PRODUCT OF RELATIVELY REGULAR OPERATORS

J.J. KOLIHA, Parksville

**Abstract:** The product  $T_1 T_2$  of relatively regular bounded linear operators between Banach spaces is shown to be relatively regular iff the product  $QP$  is relatively regular, where  $Q$  is a projection parallel to the null space of  $T_1$ , and  $P$  a projection onto the range of  $T_2$ . The paper gives applications of this result.

**Key words and phrases:** Relatively regular operator, pseudo-inverse, strict pseudo-inverse, projection, spectrum, resolvent.

AMS: Primary 47A99, 47A50

Ref. Ž.: 7.972.53

Secondary 47B99, 57A10

1. The Main Result

Let  $X$  and  $Y$  be complex Banach spaces, and let  $B(X, Y)$  be the space of all bounded linear operators from  $X$  to  $Y$ ;  $B(X, X)$  is written as  $B(X)$ . If  $T \in B(X, Y)$ ,  $N(T)$  and  $R(T)$  denote the null space and the range of  $T$ , respectively. An operator  $S \in B(Y, X)$  is called a pseudo-inverse of  $T \in B(X, Y)$  if it satisfies the equation

$$(1) \quad TST = T ;$$

an operator  $T \in B(X, Y)$  is called relatively regular if it possesses a pseudo-inverse. It is known (Caradus [5], Nashed [9]) that  $T$  is relatively regular iff  $N(T)$  is complemented in  $X$  and  $R(T)$  closed and complemented in  $Y$ . If  $S$

is a pseudo-inverse of  $T$ ,  $ST$  is a projection of  $X$  parallel to  $N(T)$ , and  $TS$  a projection of  $Y$  onto  $R(T)$ , i.e.,  $N(ST) = N(T)$  and  $R(TS) = R(T)$ . If  $T \in B(X, Y)$  is a relatively regular operator, we can find a pseudo-inverse  $S$  of  $T$  which satisfies, in addition to (1), the equation

$$(2) \quad STS = S$$

(cf. Caradus [5], p. 9). In this paper we call such  $S$  a strict pseudo-inverse of  $T$ .

Caradus [4], [5] recently initiated the study of relative regularity of the product  $T_1T_2$  of two relatively regular operators in  $B(X)$  based on the product  $QP$  of a projection  $Q \in B(X)$  onto  $R(T_2)$  and a projection  $P \in B(X)$  parallel to  $N(T_1)$ , and obtained useful sufficient conditions. The present paper continues in this investigation, and gives a complete solution to the problem of the relation between relative regularity of  $T_1T_2$  and of  $QP$ .

Let  $X$ ,  $Y$  and  $Z$  be complex Banach spaces, and let  $T_1 \in B(Y, Z)$  and  $T_2 \in B(X, Y)$  be two relatively regular operators. Sufficient conditions for the relative regularity of  $T_1T_2$  have been given by various authors (e.g. Atkinson [1], Caradus [4], [5], Koliha [8]). Bouldin [2], [3] has found a necessary and sufficient condition in the case when  $X = Y = Z$  is a Hilbert space. We show that the relative regularity of the product  $T_1T_2$  is equivalent to the relative regularity of the product of two projections in  $B(Y)$ .

Theorem 1: Let  $T_1 \in B(Y, Z)$  and  $T_2 \in B(X, Y)$  be relatively regular operators with pseudo-inverses  $S_1$  and  $S_2$ ,

respectively. Then the operator  $T_1T_2$  is relatively regular  
iff the operator  $S_1T_1T_2S_2 \in B(Y)$  is relatively regular. If  
 $U$  is a pseudo-inverse of  $T_1T_2$ ,  $T_2UT_1$  is a pseudo-inverse  
of  $S_1T_1T_2S_2$ , then  $S_2T_2S_2VS_1T_1S_1$  is a pseudo-inverse of  
 $T_1T_2$ .

Proof: Let us first assume that  $U$  is a pseudo-inverse of  $T_1T_2$ . Then

$$\begin{aligned} S_1T_1T_2S_2(T_2UT_1)S_1T_1T_2S_2 &= S_1T_1(T_2S_2T_2)U(T_1S_1T_1)T_2S_2 \\ &= S_1T_1T_2UT_1T_2S_2 \\ &= S_1T_1T_2S_2, \end{aligned}$$

so that  $T_2UT_1$  is a pseudo-inverse of  $S_1T_1T_2S_2$ . Conversely, suppose that  $V \in B(Y)$  is a pseudo-inverse of  $S_1T_1T_2S_2$ . Let us write  $P = T_2S_2$  and  $Q = S_1T_1$ . Then  $P \in B(Y)$  is a projection onto  $R(T_2)$ ,  $Q \in B(Y)$  projection parallel to  $N(T_1)$ , and  $QPVQP = QP$ . Put

$$(3) \quad W = PVQ.$$

Then

$$(4) \quad PWQ = W \quad \text{and} \quad QWP = QP.$$

From (4) we get  $Q(I - W)P = 0$ . Hence  $I - W$  maps  $R(P) = R(T_2)$  into  $N(Q) = N(T_1)$  in  $Y$ , and  $T_1(I - W)T_2 = 0$ . Thus

$$\begin{aligned} T_1T_2S_2WS_1T_1T_2 &= T_1PWQT_2 = T_1WT_2 \\ &= T_1T_2 - T_1(I - W)T_2 \\ &= T_1T_2, \end{aligned}$$

which shows that  $S_2WS_1 = S_2PVQS_1 = S_2T_2S_2VS_1T_1S_1$  is a pseudo-inverse of  $T_1T_2$ .

Remark. Let  $S_1$  and  $S_2$  be strict pseudo-inverses of  $T_1$  and  $T_2$ , respectively. Then each pseudo-inverse  $V$  of  $S_1T_1T_2S_2$  yields the pseudo-inverse  $S_2VS_1$  of  $T_1T_2$ . It is readily verified that in this case  $S_2VS_1$  is a strict pseudo-inverse whenever  $V$  is a strict pseudo-inverse.

Assume that  $T_1 \in B(Y,Z)$  and  $T_2 \in B(X,Y)$  are relatively regular, so that the spaces  $N(T_1)$  and  $R(T_2)$  are closed and complemented in  $Y$ . Let  $P \in B(Y)$  be a projection onto  $R(T_2)$ ,  $I - Q \in B(Y)$  a projection onto  $N(T_1)$ . Then we can find pseudo-inverses  $S_1$  and  $S_2$  of  $T_1$  and  $T_2$ , respectively, such that  $T_2S_2 = P$  and  $S_1T_1 = Q$  (Caradus [5]). Then, according to Theorem 1,  $T_1T_2$  is relatively regular iff the product  $QP$  of the projections  $Q, P \in B(Y)$  is relatively regular. This proves the following result.

Theorem 2: Let  $T_1 \in B(Y,Z)$  and  $T_2 \in B(X,Y)$  be relatively regular operators. Then the following conditions are equivalent:

(i) There exists a projection  $P \in B(Y)$  onto  $R(T_2)$  and a projection  $Q \in B(Y)$  parallel to  $N(T_1)$  such that the product  $QP$  is relatively regular in  $B(Y)$ .

(ii) For each projection  $P \in B(Y)$  onto  $R(T_2)$  and each projection  $Q \in B(Y)$  parallel to  $N(T_1)$  the product  $QP$  is relatively regular in  $B(Y)$ .

(iii) The product  $T_1T_2$  is relatively regular in  $B(X,Z)$ .

If (i) is satisfied with a pseudo-inverse  $V$  of  $QP$  and if  $S_1, S_2$  are strict pseudo-inverses of  $T_1, T_2$  such that  $T_2S_2 = P$  and  $S_1T_1 = Q$ , then the operator

$$(5) \quad U = S_2 V S_1$$

is a pseudo-inverse of  $T_1 T_2$ , strict whenever  $V$  is strict.

## 2. Special cases

In this section we assume that  $T_1 \in B(Y, Z)$  and  $T_2 \in B(X, Y)$  are relatively regular operators with strict pseudo-inverses  $S_1$  and  $S_2$ , respectively. Unless stated otherwise, we assume that  $P$  and  $Q$  are the projection operators  $P = T_2 S_2$  and  $Q = S_1 T_1$ .

(I) If  $T_1$  (resp.  $T_2$ ) is regular,  $T_1 T_2$  is relatively regular with a strict pseudo-inverse  $S_2 S_1$ , where  $S_1 = T_1^{-1}$  (resp.  $S_2 = T_2^{-1}$ ). This follows from (5) when we observe that under our assumptions  $Q = I$  and  $V = P$  (resp.  $P = I$  and  $V = Q$ ).

(II) If  $QP$  is a projection,  $T_1 T_2$  is relatively regular with a strict pseudo-inverse  $S_2 Q P S_1 = S_2 S_1 T_1 T_2 S_2 S_1$ . Indeed, the projection  $QP$  is relatively regular with pseudo-inverse  $QP$ .

(III) If  $Q$  and  $P$  commute,  $T_1 T_2$  is relatively regular with a strict pseudo-inverse  $S_2 S_1$ . If  $QP = PQ$ , then  $QP$  is a projection, and  $S_2 Q P S_1 = S_2 P Q S_1 = S_2 T_2 S_2 S_1 T_1 S_1 = S_2 S_1$  is a pseudo-inverse of  $T_1 T_2$  by (II). This result has been obtained by Caradus [4]; cf. also [5], p. 36. As a corollary, we obtain the following result (Caradus [5], p. 37): If  $T_1$  and  $T_2$  are relatively regular with either  $N(T_1)$  finite dimensional or  $R(T_2)$  finite codimensional, then  $T_1 T_2$  is relatively regular. This contains as special case a theorem

due to Atkinson [1] on the product of semi-Fredholm operators.

(IV) If  $\lambda = 0$  is a pole of  $(\lambda I - QP)^{-1}$  of order 1,  $T_1 T_2$  is relatively regular with a strict pseudo-inverse  $U = S_2(O_N \oplus (QP_R)^{-1})S_1$ , where  $O_N$  is the zero operator on  $N(QP)$  and  $P_R$  the restriction of  $P$  to  $R(QP)$ . First of all,  $X = N(QP) \oplus R(QP)$  with  $R(QP)$  closed (Taylor [10], p. 306). This means that  $QP$  is relatively regular. Moreover,  $QP_R$  is a bijective operator on the Banach space  $R(QP)$ , and hence continuously invertible on  $R(QP)$  by the open mapping theorem. The operator  $V = O_N \oplus (QP_R)^{-1}$  is a strict pseudo-inverse of  $QP$ , so that  $U = S_2 V S_1$  is a strict pseudo-inverse of  $T_1 T_2$ .

(V) If  $\lambda = 0$  is a pole of  $(\lambda I - QP)^{-1}$  of order 1, and if the spectrum of  $QP$  is contained in the set  $\{\lambda : |\lambda^2 - \alpha^{-1}| < \alpha^{-1}\} \cup \{0\}$  for some  $\alpha > 0$ , then  $T_1 T_2$  is relatively regular with pseudo-inverses  $U_j = S_2 V_j S_1$ ,  $j = 1, 2$ , where

$$(6) \quad V_1 = \sum_{n=0}^{\infty} \alpha (I - \alpha(QP)^2)^n QP,$$

and where  $V_2$  is the projection onto  $R((PQ)^2)$  parallel to  $N((PQ)^2)$  (Caradus [4],[5], Koliha [8]). Theorems 2 and 3 of [8] show that, for any fixed  $\alpha > 0$ , the two conditions on the spectrum of  $QP$  given above are necessary and sufficient for the convergence of the series (6); the sum  $V_1$  is then a strict pseudo-inverse of  $QP$ . From the equations (3) and (4) it follows that  $V_2 = P V_1 Q$  is also a (strict) pseudo-inverse of  $QP$ . Then

$$V_2 = P \left( \sum_{n=0}^{\infty} \alpha (I - \alpha(QP)^2)^n QP \right) Q = \sum_{n=0}^{\infty} \alpha PQ (I - \alpha(PQ)^2)^n PQ$$

$$= \sum_{n=0}^{\infty} \alpha(PQ)^2 (I - \alpha(PQ)^2)^n = I - \lim_{N \rightarrow \infty} (I - \alpha(PQ)^2)^N = I - W,$$

where  $W = \lim_{N \rightarrow \infty} (I - \alpha(PQ)^2)^N$  is the projection onto  $N((PQ)^2)$  parallel to  $R((PQ)^2)$  (cf. [7], Theorem 4). This improves on a result of Caradus [4].

(VI) If there exists  $L \in B(Y)$  such that the operator  $Q(I - PLQ)P$  is relatively regular, then  $T_1 T_2$  is relatively regular. This follows from Theorem 2 and a result of Atkinson [1] that an operator  $T \in B(Y)$  is relatively regular if  $T - TLT$  is relatively regular for some  $L \in B(Y)$ .

(VII) If the operator  $(QP)^2$  is Fredholm, then  $T_1 T_2$  is relatively regular. Atkinson [1] proved that operators  $S$  and  $T$  are relatively regular if  $ST$  is Fredholm. Our result follows on setting  $S = T = QP$  and applying Theorem 2.

(VIII) Let  $Y$  be a Hilbert space,  $P$  the orthogonal projection of  $Y$  onto  $R(T_2)$  and  $Q$  the orthogonal projection of  $Y$  onto  $N(T_1)$ . The operator  $T_1 T_2$  is relatively regular iff the range of  $QP$  (or equivalently of  $PQ$ ) is closed. This follows from the fact that closed subspaces of Hilbert space are complemented. Bouldin's result [2] shows that the range  $R(QP)$  is closed iff the subspaces

$$R(T_2) \text{ and } N(T_1) \cap [N(T_1) \cap R(T_2)]^\perp$$

enclose a positive angle. A pseudo-inverse  $U$  of  $T_1 T_2$  may be obtained from Groetsch's representation theorem [6] as follows: Let  $A$  be the restriction of the operator  $I - PQP \in B(Y)$  to  $R(PQ)$ , let  $\Omega$  be an open subset of the interval  $(-\infty, 1]$  containing the spectrum of  $A$ , and let  $\{S_\alpha\}$  be a



net of continuous real functions on  $\Omega$  such that  
 $\lim_{\alpha} S_{\alpha}(x) = 1/(1-x)$  uniformly on the spectrum of  $A$ .  
 Then

$$U = \lim_{\alpha} S_2 S_{\alpha}(A) P Q S_1$$

is the uniform operator topology of  $B(X, Y)$ .

#### R e f e r e n c e s

- [1] F.V. ATKINSON: On relatively regular operators, Acta Sci. Math. Szeged 15(1953), 38-56.
- [2] R. BOULDIN: The product of operators with closed range, Tohoku Math. J. 25(1973), 359-363.
- [3] R. BOULDIN: The pseudo-inverse of a product, SIAM J. Appl. Math. 24(1973), 489-495.
- [4] S.R. CARADUS: An equational approach to products of relatively regular operators (submitted to Aequationes Math.).
- [5] S.R. CARADUS: Operator theory of the pseudo-inverse, Queen's Papers in Pure and Applied Mathematics, No. 38, Queen's University, Kingston, Ont., 1974.
- [6] C.W. GROETSCH: Representations of the generalized inverse, J. Math. Anal. Appl. 49(1975), 154-157.
- [7] J.J. KOLIHA: Convergent and stable operators and their generalization, J. Math. Anal. Appl. 43(1973), 778-794.
- [8] J.J. KOLIHA: Convergence of an operator series, Research Report No. 19, Department of Mathematics, University of Melbourne, 1974.
- [9] M.Z. NASHED: Generalized inverses, normal solvability and iteration for singular operator equations, in Nonlinear Functional Analysis and Applications, L.B. Rall, Ed., Academic Press, New York, 1971,

pp. 311-359.

[10] A.E. TAYLOR: Introduction to Functional Analysis, J. Wiley, New York, 1958.

Department of Mathematics  
University of Melbourne  
Parkville, Victoria 3052  
Australia

(Oblatum 1.4.1975)