

Jiří Reif

Some remarks on subspaces of weakly compactly generated Banach spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 16 (1975), No. 4, 787--793

Persistent URL: <http://dml.cz/dmlcz/105666>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1975

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

SOME REMARKS ON SUBSPACES OF WEAKLY COMPACTLY GENERATED
BANACH SPACES

J. REIF, Praha

Abstract: Some hereditary properties of weakly compactly generated Banach spaces are shown.

Key words: Weakly compactly generated Banach space, Eberlein compact, densities property of a Banach space.

AMS: 46B10

Ref. Ž. 7.972.22

Introduction. We work through the paper only with real Banach spaces. A Banach space X is said to be weakly compactly generated (WCG) if there exists a weakly compact set $K \subset X$ which generates X , i.e. the closed linear span of K is X .

Recently, Rosenthal [7] has shown that a closed subspace of a WCG space need not be WCG. Such a subspace (even with an unconditional basis) was found in the space $L_1(\mu)$ for a finite measure μ . We have remarked in this paper some properties of WCG spaces which are hereditary to general closed linear subspaces, e.g. a certain densities property (Proposition 5).

We mention our notation. For a Banach space X we denote B_X^* the unit ball of X^* with the w^* topology. For

a topological (completely regular Hausdorff) space T
 $C_b(T)$ denotes the space of real-valued functions on T
under the supremum norm.

By a subspace of a Banach space we mean always a closed linear subspace.

We quote at first a few marked properties of WCG spaces which are (some of them less evidently) kept by general subspaces of WCG spaces.

Proposition 1. Let X be a subspace of a WCG space. Then there holds:

- (i) X has an equivalent norm which is LUR;
- (ii) X has an equivalent norm such that X^* is strictly convex;
- (iii) X has a Markušević basis;
- (iv) if $c_0 \subset X$ then there exists a linear projection P of X onto c_0 with $\|P\| \leq 2$.

Proof: Properties (i) and (ii) are hereditary, (iii) can be proved using the method of [5] and the decomposition of subspaces of WCG spaces in [3]. (iv) holds by the results of [8] and [3].

We use the following easy characterization of subspaces of WCG spaces for the next.

Lemma. A Banach space X is a subspace of a WCG space if and only if the unit ball of X^* with the w^* topology is a continuous image of an Eberlein compact.

Proof: Let X be a subspace of a WCG space Y . The

restriction mapping $R: Y^* \rightarrow X^*$ defined by $Rf = f|_X$ for $f \in Y^*$ is $w^* - w^*$ continuous and $R(B_Y^*) = B_X^*$ by Hahn-Banach theorem. The space B_Y^* (with the w^* topology) is an Eberlein compact ([1]) and B_X^* is a continuous image of it.

On the other hand, let X be a Banach space and B_X^* a continuous image of an Eberlein compact K . Then we can suppose the inclusions $X \subset C(B_X^*) \subset C(K)$ and the latter space is WCG ([1]).

There is observed in [4] that if $T: X \rightarrow Y$ is a linear continuous mapping with the range dense in Y and X is WCG, then so is Y . Indeed, if K is a weakly compact set generating X , then $T(K)$ is a weakly compact set generating Y . We make an analogy to this within subspaces of WCG spaces.

Proposition 2. Let both X, Y be Banach spaces and $T: X \rightarrow Y$ a continuous linear mapping with $\overline{TX} = Y$. Suppose X is a subspace of a WCG space. Then so is Y .

Proof: The mapping $T^*: Y^* \rightarrow X^*$ is $w^* - w^*$ continuous and one-to-one. Accordingly, T^* is a homeomorphism on B_Y^* and we can assume the inclusion $B_Y^* \subset |T^*| \cdot B_X^*$. Since the property "to be a continuous image of an Eberlein compact" is closed hereditary, our assertion is a consequence of the Lemma.

Remark. Each Eberlein compact K has the following property due to Kaplansky: if $A \subset K$ and $x \in \overline{A}$, then there exists a sequence $\{x_n\}_{n=1}^{\infty}$ in A such that $x_n \rightarrow x$. It

is easy to verify that this property is kept by continuous Hausdorff images of Eberlein compacts.

Proposition 3. Let X be a subspace of a WCG space and $K \subset X^*$ a w^* -sequentially closed set which is either bounded or convex. Then K is w^* -closed.

Proof: For any $r > 0$ the set $K \cap \{x \in X^*; \|x\| \leq r\}$ is w^* -closed by the Remark and Lemma.

Corollary. Let f be a convex function on X^* where X is a subspace of a WCG space. Then f is w^* -lower semicontinuous if it is sequentially w^* -lower semicontinuous.

Proposition 4. Let X be a topological Hausdorff completely regular space. Suppose $C_b(X)$ is a subspace of a WCG space. Then there holds:

- (a) X is pseudocompact;
- (b) X is compact if it is normal.

Proof: Suppose X is not pseudocompact. Then there exists an infinite discrete set $\Gamma \subset X$ which is C -embedded into X , i.e. the restriction mapping $R: C_b(X) \rightarrow m(\Gamma)$ (defined by $Rf = f/\Gamma$ for $f \in C_b(X)$) is onto $m(\Gamma)$. The space $m(\Gamma)$ cannot be a subspace of a WCG space by Proposition 1. Consequently, the space $C_b(X)$ cannot be a subspace of a WCG space by Proposition 2, a contradiction. Let X be now moreover normal. Denote B_C^* the unit ball of $C_b^*(X)$ with the w^* topology. As for the Čech-Stone compactification βX of X we can assume $\beta X \subset B_C^*$, βX is a continuous image of an Eberlein compact by the Lemma. Thus βX has the property of Kaplansky from the Remark.

It implies easily (provided X is normal) that X must be compact.

D. Preiss and P. Simon have shown recently that if K is a pseudocompact subset of an Eberlein compact, then K is compact ([6]). Consequently, if for a Hausdorff completely regular space X the space $C_b(X)$ is WCG, then X is compact.

For a topological space X dX (density of X) is the smallest cardinal number \aleph such that there exists a subset A dense in X with $\text{card } A = \aleph$.

The next property and also Corollary 1 are proved in [4] for WCG spaces, but the method used there cannot be utilized in our case.

Proposition 5. Let X be a subspace of a WCG space. Then for the densities of X and X^* we have the equality $dX = d(X^*, w^*)$.

Proof: For any normed linear space there holds $d(X^*, w^*) \leq dX$. Thus for X separable our assertion is evident. So suppose X is a non-separable subspace of a WCG space Y . We can assume $dX = dY$ ([1]). Suppose the inequality $d(X^*, w^*) = dX$ is false, i.e. let A be a w^* -dense subset of X^* with $\text{card } A < dX$. Since X is non-separable we can assume that $\text{card } A \geq \aleph_0$.

Let $\tilde{f} \in Y^*$ be an extension of f for each $f \in A$ and denote $\tilde{A} = \{\tilde{f}; f \in A\}$. By [3] there is a continuous linear projection $P: Y \rightarrow Y$ with $PX \subset X$, $P^* \tilde{f} = \tilde{f}$ for $\tilde{f} \in \tilde{A}$ and $d(PY) \leq \text{card } \tilde{A}$. We define the projection $S: X \rightarrow X$

by the restriction of P on X . Clearly, $S^*f = f$ for $f \in \epsilon A$. Since A is w^* -dense in X^* and S^* is w^* - w^* continuous we have $S^* = id_{X^*}$. Consequently, $S = id_X$ and hence $X \subset PY$. But for the densities we have $d(PY) \leq \text{card } A < < dX$, thus $d(PY) < dX$, a contradiction.

Corollary 1. Let X be a Banach space such that X^* is a subspace of a WCG space. Then $dX = dX^*$.

Proof: For any normed linear space there holds $d(X^{**}, w^*) \leq dX \leq dX^*$, and the first member of the inequality is equal to the last one by Proposition 5.

Corollary 2 (cf. [2]). Let X be as in Corollary 1. Then X has the densities property, i.e. for each subspace $Y \subset X$ there is $dY^* = dY$. Thus X^* has the Radon-Nikodym property.

Proof: Suppose Y is a subspace of X . Then Y^* is a continuous linear image of X^* and thus Y^* is a subspace of a WCG space by Proposition 2. Consequently, $dY = dY^*$ by Corollary 1.

If X has the densities property, then X^* has the Radon-Nikodym property, see e.g. [2].

R e f e r e n c e s

- [1] D. AMIR, J. LINDENSTRAUSS: The structure of weakly compact sets in Banach spaces, Ann. of Math. 88(1968), 35-46.
- [2] J. DIESTEL, J.J. UHL: The Radon-Nikodym theorem for Banach space valued measures, to appear.

- [3] K. JOHN, V. ZIZLER: Smoothness and its equivalents in weakly compactly generated Banach spaces, Journ. Funct. Anal. 15(1974), 1-11.
- [4] J. LINDENSTRAUSS: Weakly compact sets, their topological properties and Banach spaces they generate, Ann. of Math. Stud. 69, Princeton Univ. Press (1972), 235-273.
- [5] J. LINDENSTRAUSS: Decomposition of Banach spaces, Indiana Univ. Journ. 20(1971), 817-819.
- [6] D. PREISS, P. SIMON: A weakly pseudocompact subspace of a Banach space is weakly compact, Comment. Math. Univ. Carolinae 15(1974), 603-611.
- [7] H.P. ROSENTHAL: The heredity problem for weakly compactly generated Banach spaces, Comp. Math. 28(1974), 83-111.
- [8] A. SOBCZYK: Projection of the space (m) on its subspace (c_0) , Bull. Amer. Math. Soc. 47(1941), 938-947.
- [9] G. STEGALL: The Radon-Nikodym property in conjugate Banach spaces, to appear.
- [10] S. TROJANSKI: On locally uniformly convex and differentiable norms in certain non-separable Banach spaces, Studia Math. 37(1971), 173-180.
- [11] W.A. VEECH: Short proof of Sobczyk theorem, Proc. Amer. Math. Soc. 28(1971), 627-628.

Matematický ústav

Karlova universita

Sokolovská 83, 18600 Praha 8

Československo

(Oblatum 3.7. 1975)