

Juhani Nieminen

Tolerance relations on complete lattices

Commentationes Mathematicae Universitatis Carolinae, Vol. 18 (1977), No. 4, 639--645

Persistent URL: <http://dml.cz/dmlcz/105808>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1977

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

TOLERANCE RELATIONS ON COMPLETE LATTICES

Juhani NIEMINEN, Vaasa

Abstract: It is shown that each compatible tolerance relation T on a complete lattice L has a homotopy representation by means of two semicongruences induced by T on L .

Key words: Tolerance, homotopy representation.

AMS: 06A23

Ref. Ž.: 2.724.38

The purpose of this short paper is to show that each compatible tolerance relation on a complete lattice has the property of the homotopy type, i.e. a compatible tolerance relation T on a complete lattice L can be decomposed into two semicongruences on L and, on the other hand, expressed by means of these two semicongruences. The concept of homotopy suitable for the approach here was introduced by Petrescu in [4]. The other observations of this note are based on the characterization of compatible tolerance relations by means of \mathcal{C} -coverings and related mappings given by Chajda, Niederle and Zelinka in [1]. For other properties of tolerance relations on algebras the reader is referred to the recent paper [3] of Chajda and Zelinka and to the references therein.

Let $\mathcal{A} = \langle A, F \rangle$ be an algebra with the support A and

the set F of fundamental operations. A tolerance relation T on the set A is a binary, reflexive and symmetric relation on A . T is compatible with \mathcal{A} , if for any n -ary operation $f \in F$, where n is a positive integer, and for arbitrary elements $a_1, \dots, a_n, b_1, \dots, b_n$ of satisfying $a_i T b_i$ for $i = 1, \dots, n$, we have $f(a_1, \dots, a_n) T f(b_1, \dots, b_n)$.

Let M be a non-empty set. The family $\mathcal{M} = \{M_\gamma, \gamma \in \Gamma\}$, where Γ is a subscript set, is called a covering of M by subsets, if and only if M_γ is for each $\gamma \in \Gamma$ a subset of M and $\bigcup_{\gamma \in \Gamma} M_\gamma = M$. As usually, we suppose that $M_\gamma \neq M_\beta$ for $\gamma \neq \beta, \gamma, \beta \in \Gamma$. A covering $\mathcal{M} = \{M_\gamma, \gamma \in \Gamma\}$ of M is called a τ -covering of M , if and only if \mathcal{M} satisfies the following two conditions

(i) if $\gamma_0 \in \Gamma$ and $\Gamma_0 \subseteq \Gamma$, then $M_{\gamma_0} \subseteq \bigcup_{\gamma \in \Gamma_0} M_\gamma$ and $\bigcap_{\gamma \in \Gamma_0} M_\gamma \subseteq M_{\gamma_0}$;

(ii) if $N \subseteq M$ and N is not contained in any set from \mathcal{M} , then N contains a two-element subset of the same property.

The following lemma shows the connection between tolerance relations on M and τ -coverings of M [1, Thm. 1].

Lemma 1. Let M be a non-empty set. There exists then a one-to-one correspondence between tolerance relations on M and τ -coverings of M such that if T is a tolerance relation on M and \mathcal{M}_T is the τ -covering of M corresponding to T , then any two elements of M are in the relation T if and only if there exists a set from \mathcal{M}_T which contains both of them.

The second lemma [1, Thm. 3] illuminates the properties of compatible tolerances on algebras.

Lemma 2. Let $\mathcal{A} = \langle A, F \rangle$ be an algebra, T a tolerance on \mathcal{A} and \mathcal{M}_T a τ -covering of \mathcal{A} corresponding to T . The tolerance T is compatible with \mathcal{A} , if and only if there exists an algebra $\mathcal{B} = \langle B, G \rangle$ with the following properties:

(i) there exists a one-to-one mapping $\varphi : F \rightarrow G$ such that for any positive integer n and for each $f \in F$ the operation φf is n -ary if and only if f is n -ary;

(ii) there exists a one-to-one mapping $\chi : \mathcal{M}_T \rightarrow B$ such that for each n -ary operation $f \in F$ and for any $n + 1$ elements M_0, M_1, \dots, M_n from \mathcal{M}_T the equality $\varphi f(\chi(M_1), \dots, \chi(M_n)) = \chi(M_0)$ implies that for any n elements a_1, \dots, a_n of A such that $a_i \in M_i$, $i = 1, \dots, n$, the element $f(a_1, \dots, a_n) \in M_0$.

Let $\mathcal{A} = \langle A, F \rangle$ and $\mathcal{B} = \langle B, G \rangle$ be two algebras of the same type. Let n_0 be the maximum number n for which there exists an n -ary operation f on \mathcal{A} and I the interval $[1, n_0]$. A family $\mathfrak{F} = [(\alpha_i)_{i \in I}; \beta]$ of mappings of A into B such that

$\beta(f(a_1, \dots, a_n)) = f(\alpha_1(a_1), \dots, \alpha_n(a_n))$ for every $n \leq n_0$, $a_1, \dots, a_n \in A$, is called a homotopy of \mathcal{A} into \mathcal{B} . The mappings α_i are called components of homotopy \mathfrak{F} and β the principal component of \mathfrak{F} . Moreover, it is shown that each α_i induces an equivalence relation on A [4, Lemma 0.1].

We shall show that the mapping χ relating to a compatible tolerance T on L is a principal component of a homotopy induced by T . The components α_1 and α_2 are generated by semicongruences on L which are induced by the τ -covering \mathcal{M}_T of T . We shall construct α_1 which is given by an equivalence relation $E(\alpha_1)$ being compatible with respect to the

\wedge -operation on L , i.e. by a \wedge -semicongruence. The equivalence relation $E(\alpha_1)$ is constructed by determining the partition ε inducing $E(\alpha_1)$. ε is obtained by modifying the τ -covering \mathcal{M}_T of the compatible tolerance T on L .

Theorem 1. Let T be a compatible tolerance on a complete lattice L , \mathcal{M}_T the τ -covering corresponding to T and let $M_\gamma \in \mathcal{M}_T$. Then the family of sets $\mathcal{M}_T^\wedge = \{M_\sigma^\wedge, \sigma \in \Gamma\}$, where $M_\sigma^\wedge = M_\sigma \setminus \bigcup_{\gamma \neq \sigma} \{M_\gamma \mid M_\gamma \cap M_\sigma \neq \emptyset\}$, the least element $e_{1\sigma}$ of M_σ is greater than the least element $e_{1\gamma}$ of M_γ ($e_{1\sigma} < e_{1\gamma}$), $M_\sigma \in \mathcal{M}_T$, forms a partition of L determining a \wedge -semicongruence on L .

Proof. According to [2, Thm. 1], each $M_\gamma \in \mathcal{M}_T$ is a convex sublattice of L , and as L is complete, there are in M_γ the least and greatest elements $e_{1\gamma}$ and $e_{g\gamma}$, respectively.

According to the definition of M_σ^\wedge , each M_σ^\wedge contains at least $e_{1\sigma}$, whence $M_\sigma^\wedge \neq \emptyset$ for each $\sigma \in \Gamma$. Moreover, the properties of T imply that when $a, b \in M_\sigma^\wedge$ then also $a \wedge b \in M_\sigma^\wedge$. Thus the theorem holds, if we can show that any element $x \in L$ belongs to at least one set M_σ^\wedge of \mathcal{M}_T^\wedge , and $M_\sigma^\wedge \cap M_\gamma^\wedge = \emptyset$ for each pair $\sigma, \gamma \in \Gamma$ when $\sigma \neq \gamma$.

Let $a \in L$ and $\mathcal{M}_T(a)$ be the family of all subsets of \mathcal{M}_T containing the element a . Let $M_\gamma, M_{\sigma\epsilon} \in \mathcal{M}_T(a)$ be such sets that $e_{1\gamma}$ and $e_{1\sigma\epsilon}$ are non-comparable. Let q be the least element of the set $M_\gamma \cap M_{\sigma\epsilon}$; such an element q exists and $q \in M_\gamma \cap M_{\sigma\epsilon}$, since L is complete and as an intersection of two convex sets $M_\gamma \cap M_{\sigma\epsilon}$ is a convex set of L , too. As $q \in M_\gamma, M_{\sigma\epsilon}$, $q \geq e_{1\gamma} \vee e_{1\sigma\epsilon}$ and as $e_{1\gamma} \vee e_{1\sigma\epsilon} \in M_\gamma \cap M_{\sigma\epsilon}$,

$q \leq e_{1\gamma} \vee e_{1\lambda}$, whence $q = e_{1\gamma} \vee e_{1\lambda}$. According to the compatibility of T , any two elements of the interval $[e_{1\gamma} \vee e_{1\lambda}, e_{g\gamma} \vee e_{g\lambda}]$ are in the relation T . Thus $[e_{1\gamma} \vee e_{1\lambda}, e_{g\gamma} \vee e_{g\lambda}] \subseteq M_\lambda \in \mathcal{M}_T(a)$ for some index $\lambda \in \Gamma$. If $e_{1\lambda} \leq e_{1\gamma}$, then $M_\gamma \notin \mathcal{M}_T$ according to the condition (i) for \mathcal{M}_T ; the same holds for M_λ , too. If $e_{1\gamma}$ and $e_{1\lambda}$ are non-comparable, then $(e_{1\gamma} \wedge e_{1\lambda})Te_{g\gamma}$, as $e_{g\gamma} \in M_\gamma, M_\lambda$. Then $e_{1\gamma} > e_{1\gamma} \wedge e_{1\lambda}$, and so there were in \mathcal{M}_T a set containing properly M_γ , which is a contradiction. Hence $e_{1\gamma}, e_{1\lambda} \leq e_{1\lambda} \leq e_{1\gamma} \vee e_{1\lambda}$ and so $e_{1\lambda} = e_{1\gamma} \vee e_{1\lambda}$. Consequently, there is in $\mathcal{M}_T(a)$ for any two sets M_γ, M_λ a third set M_λ such that $e_{1\lambda} = e_{1\gamma} \vee e_{1\lambda}$. As L is complete, there is also an element $\bigvee_{\lambda \in \Gamma} e_{1\lambda} = e_{1\varphi}$, where $e_{1\varphi}$ is the least element of a subset M_φ belonging to the τ -covering \mathcal{M}_T and containing the element a . According to the definition of M_φ and to the maximality of $e_{1\varphi}$ with respect to a , $a \in M_\varphi$, and so any element of L belongs to at least one of the sets $M_\gamma, \gamma \in \Gamma$.

If $M_\gamma \cap M_\lambda \neq \emptyset, \gamma \neq \lambda$, then we can prove as above that $e_{1\sigma} \vee e_{1\gamma} \in M_\sigma \cap M_\gamma$. But this is the least element of a subset $M_\lambda \in \mathcal{M}_T, e_{1\lambda} > e_{1\sigma}, e_{1\gamma}$, and thus, according to the definitions of M_σ and $M_\gamma, e_{1\lambda} \in M_\sigma, M_\gamma$. This is a contradiction, whence $M_\sigma \cap M_\gamma = \emptyset$ for any pair $\sigma, \gamma \in \Gamma, \sigma \neq \gamma$. This completes the proof.

Let T be a compatible tolerance on a complete lattice L and χ a mapping, $\chi : \mathcal{M}_T \rightarrow B$, induced by T and defined in Lemma 2. As for any $\gamma \in \Gamma$ there exists a unique subset

$M_{\mathcal{F}}^{\wedge} \in \mathcal{M}_{\mathcal{T}}^{\wedge}$ of L , we can define a mapping $\alpha_1: L \rightarrow B$ as follows: for any $a \in L$, $\alpha_1(a) = \chi(M_{\mathcal{F}})$ if and only if $a \in M_{\mathcal{F}}^{\wedge}$ in $\mathcal{M}_{\mathcal{T}}^{\wedge}$.

By using the dual proof of Theorem 1, we can show the existence of a partition $\mathcal{M}_{\mathcal{T}}^{\vee}$ of L determining a \vee -semi-congruence on L . As above, we define the mapping $\alpha_2: L \rightarrow B$ induced by $\mathcal{M}_{\mathcal{T}}^{\vee}$ and χ : for any $a \in L$, $\alpha_2(a) = \chi(M_{\mathcal{F}})$ if and only if $a \in M_{\mathcal{F}}^{\vee}$ in $\mathcal{M}_{\mathcal{T}}^{\vee}$. Now we are able to state our main theorem

Theorem 2. Let L be a complete lattice, \mathcal{T} a compatible tolerance on L , $\mathcal{M}_{\mathcal{T}}$ the corresponding τ -covering of L and χ the mapping, $\chi: L \rightarrow B$, induced by \mathcal{T} , where B is the carrier set of the algebra $\mathcal{B} = \langle B, G \rangle$ defined in Lemma 2. Then the triple $\xi = [\alpha_1, \alpha_2; \chi]$ determines a homotopy of L into $\mathcal{B} = \langle B, G \rangle$.

Proof. As χ is defined only on the family $\mathcal{M}_{\mathcal{T}}$, we have to define χ on L such that it gives the desired homotopy property. For the two operations of L we define:

$\chi(f(a_1, a_2)) = \chi(M_0)$ which is obtained from $\varphi f(\chi(M_1), \chi(M_2))$, where $a_1 \in M_1^{\wedge}$ and $a_2 \in M_2^{\vee}$ (see Lemma 2 (ii)). As $a = a \vee a = a \wedge a$ in L , we obtain $\chi(a) = \chi(f(a, a))$ which is already defined. By using this definition for $\chi: L \rightarrow B$ it obviously holds that $\chi(f(a_1, a_2)) = \varphi f(\alpha_1(a_1), \alpha_2(a_2))$ for any $a_1, a_2 \in L$, where φf can be substituted by f as L and \mathcal{B} are of the same type. This completes the proof.

References

- [1] I. CHAJDA, J. NIEDERLE and B. ZELINKA: On existence con-

ditions for compatible tolerances, Czech. Math. J. 26(1976), 304-311.

- [2] I. CHAJDA and B. ZELINKA: Tolerance relations on lattices, Časop. přest. mat. 99(1974), 394-399.
- [3] I. CHAJDA and B. ZELINKA: Lattices of tolerances, Časop. přest. mat. 102(1977), 10-24.
- [4] A. PETRESCU: On the homotopy of universal algebras (I), Rev. Roum. Math. Pures et Appl. 22(1977), 541-551.

Vaasa School of Economics

Inst. Math.

Raastuvankatu 31, 65100 Vaasa 10

Finland

(Oblatum 2.6. 1977)