

Pavel Pudlák

Symmetric embedding of finite lattices into finite partition lattices

Commentationes Mathematicae Universitatis Carolinae, Vol. 20 (1979), No. 1, 183--187

Persistent URL: <http://dml.cz/dmlcz/105912>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1979

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

SYMMETRIC EMBEDDING OF FINITE LATTICES INTO FINITE PARTITION LATTICES
P. PUDLÁK

Abstract: It has been shown that every finite lattice can be embedded into a finite partition lattice. Here we show some additional properties which such an embedding can have.

Key words: Finite lattice, partition lattice, symmetric graph, matching.

AMS: 06A20, 05C99 .

For a finite lattice L define the dimension function on L , $d:L \rightarrow \mathbb{N}$, $d(x)$ = the length of the longest maximal chain between 0 and x . Let Δ denote the kernel of d , let $x \sim y$ denote that there is $\alpha \in \text{Aut}(L)$ such that $x = \alpha(y)$. It is known that in a partition lattice $\Pi(X)$ two partitions are in the relation \sim iff they are of the same type iff they are isomorphic. The partition \sim of L is a refinement of Δ .

Let $\varphi:L \rightarrow \Pi(X)$ and let θ be the co-image of $\Delta_{\Pi(X)}$, (or $\sim_{\Pi(X)}$), i.e.

$x \theta y$ iff $d(x) = d(y)$,

(or $x \theta y$ iff $\exists \alpha \in \text{Aut}(L) \quad x = \alpha(y)$).

Then, clearly, θ satisfies the following two properties

(1) $x \theta y, x \neq y \Leftrightarrow x = y$, i.e. every class of θ is a co-chain,

x) This is a part of the CSc dissertation of the author.

(2) for no $x, y, z, t \in L$, $x \theta y$, $z \theta t$, $x < z$, $y > t$.

Theorem. If a finite lattice L and a partition θ of L satisfy (1), (2), then there is an embedding of L into some finite $\Pi(X)$ such that

$$\begin{aligned} x \theta y &\Rightarrow \varphi(x) \sim \varphi(y), \\ \neg x \theta y &\Rightarrow \neg \varphi(x) \Delta \varphi(y). \end{aligned}$$

Corollary.

1) For every finite lattice, there is an embedding into a finite partition lattice which preserves Δ .

2) The same for \sim .

Problem. Let L be a finite lattice and $d': L \rightarrow \mathbb{N}$ an arbitrary mapping such that $d'(x) < d'(y)$ whenever $x < y$. Is there always an embedding $\varphi: L \rightarrow \Pi(X)$, X finite, such that, for $y \neq 0$,

$$\frac{d'(x)}{d'(y)} = \frac{d(\varphi(x))}{d(\varphi(y))},$$

where d is the dimension on $\Pi(X)$?

Proofs

Lemma 1: Let $(L_\alpha)_{\alpha \in I}$ be a system of lattices with the following properties:

- 1) $|L_\alpha \cap L_\kappa| \leq 1$, for $\alpha \neq \kappa$,
- 2) if $x \in L_\alpha \cap L_\kappa$ and $y \in L_\beta \cap L_\gamma$ then $x = y$ or x and y are incomparable,
- 3) if G is the symmetric graph on I , in which (α, κ) is an edge iff $|L_\alpha \cap L_\kappa| = 1$, then G does not contain cycles of length < 5 .

Then adding the biggest and the smallest element to $\bigcup_I L_\alpha$, we obtain a lattice.

Proof: The proof of this lemma is just a tedious verification of basic properties of a lattice, we leave it to the reader. (Condition 3) enables us to treat the case such that for some $x \in L_\lambda$, $y \in L_\kappa$, where distance of λ, κ in G is 2, there is a nontrivial upper (or lower) bound z . Then we can derive that z must be in L_λ , where λ is uniquely determined by the fact that (λ, λ) and (λ, κ) are edges of G .)

Lemma 2: For every $k \geq 1$, there is a symmetric graph G such that

- 1) G is bipartite,
- 2) G can be decomposed into k disjoint matchings,
- 3) G does not contain cycles of length < 10 .

Proof: In [3] a graph $G_{n,m}$ is constructed for all $m, n \geq 2$, which can be decomposed into n disjoint Hamiltonian cycles, does not contain cycles of length $< m$, and is bipartite. Since $G_{n,m}$ is bipartite, the Hamiltonian cycles can be decomposed into matchings, then we can omit superfluous matchings. (Use of the result [3] was suggested by V. Röd1 .)

Let $C, D \subseteq L$ be two co-chains in a lattice L . We shall say that they are non-crossing iff for no $x, y \in C$ and $z, t \in D$, $x < z$, $y > t$. A partition Θ of L satisfies (1).(2) iff the classes of Θ are pairwise non-crossing co-chains.

Lemma 3: Let C_1, \dots, C_n be a system of non-crossing co-chains of a finite lattice L . Then there is a finite lattice K , and a system of embeddings $\varphi_\lambda : L \rightarrow K$, $\lambda \in I$, and for every i , $1 \leq i \leq n$, $x, y \in C_i$, there is a permutation π of the set of indexes I such that

$$\varphi_\lambda(x) = \varphi_{\pi(\lambda)}(y) \text{ for every } \lambda \in I.$$

Proof:

1) $n = 1$. Let $k = |C_1|$ and let $G = (Z, R)$ be the graph of Lemma 2 for k . Let $Z = Z_1 \cup Z_2$ and $R = \bigcup_{x \in C_1} R_x$ be the decompositions given by 1), 2) of Lemma 2. Take a system of distinct copies of L , say, L_z , $z \in Z_1$, such that they are also distinct from Z_2 . Then glue together x_z of L_z with κ , for every $x \in C_1$ and $(z, \kappa) \in R_x$. Since G does not contain cycles of length < 10 , we can use Lemma 1 to obtain a lattice K . For $x, y \in C_1$, the permutation π can be defined putting $\pi(z)$ equal to the unique $\kappa \in Z_1$ such that there is $\lambda \in Z_2$, $(z, \lambda) \in R_x$, and $(\lambda, \kappa) \in R_y$.

2) $n > 1$. By induction over n , using 1). We have only to add to the induction hypothesis the condition that any co-chain non-crossing with C_1, \dots, C_n is mapped by φ_z , $z \in I$, on a co-chain in K .

Proof of the Theorem: Let L, θ satisfy conditions (1), (2), L finite. Let C_1, \dots, C_n be all the classes of the partition θ . Extend L to L' and C_i to C'_i , $i = 1, \dots, n$, in such a way that for every two different C'_i, C'_j there are $x_0 \in C'_i, y_0 \in C'_j$, x_0 comparable with y_0 . Let K be the lattice given by Lemma 3 for L', C'_1, \dots, C'_n , let $\psi: K \rightarrow \Pi(X)$ be an embedding of K into a finite partition lattice. Take a system of sets X_z , $z \in I$ of the same cardinality as X , and let $\psi_z: K \rightarrow \Pi(X_z)$, $z \in I$, be some isomorphic copies of $\psi: K \rightarrow \Pi(X)$. Finally, define $\varphi: L \rightarrow \Pi(Y)$, $Y = \bigcup_I X_z$, by

$$\varphi(x) = \bigcup_I \psi_z(\varphi_z(x)).$$

Clearly, φ is an embedding. Now, let $x, y \in C'_i$, then $\varphi_z(x) = \varphi_{\pi(z)}(y)$ for some permutation π and every $z \in I$. Since ψ_z and $\psi_{\pi(z)}$ are isomorphic, we have

$$\psi_2 \varphi_2(x) \sim \psi_{\mathcal{M}(2)} \varphi_2(x) = \psi_{\mathcal{M}(2)} \varphi_{\mathcal{M}(2)}(y).$$

Thus there is a 1-1 correspondence between isomorphic parts of $\varphi(x)$ and $\varphi(y)$, which proves $\varphi(x) \sim \varphi(y)$.

On the other hand, if x, y belong to different classes C'_i, C'_j , we have $x_0 \in C'_i, y_0 \in C'_j, x_0, y_0$ comparable. Then, of course, $\varphi(x_0)$ and $\varphi(y_0)$ must have different dimension. Therefore $\varphi(x)$ and $\varphi(y)$ have different dimension.

The only thing that remains to do now is to take the restriction of φ to L .

R e f e r e n c e s

- [1] Pavel PUDLÁK, Jiří TŮMA: Every finite lattice can be embedded in a finite partition lattice, Algebra Universalis - to appear.
- [2] Pavel PUDLÁK: Representations of finite lattices (in Czech), CSc dissertation, Mathematical Institute of the Czechoslovak Academy of Sciences, Prague, 1978.
- [3] V. Neumann-Lara: k -Hamiltonian graphs with given girth, Coll. Math. Soc. János Bolyai, Infinite and Finite Sets, Keszthely, 1973, 1133-1140.

Matematický ústav

Č S A V

Žitná 25, 11567 Praha 1

Československo

(Oblatum 6.12.1978)