

Mirko Navara; Pavel Pták
On the integration on σ -classes

Commentationes Mathematicae Universitatis Carolinae, Vol. 21 (1980), No. 3, 627

Persistent URL: <http://dml.cz/dmlcz/106026>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1980

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ANNOUNCEMENTS OF NEW RESULTS

ON THE INTEGRATION ON σ -CLASSES

M. Navara, P. Pták (Technical University of Prague, Suchbátarova 2, 16627 Praha 6, Československo), received 9.4. 1980

We announce a partial solution to two problems posed by S. Gudder in the paper "A generalized measure and probability theory for the physical sciences", Foundations of Probability Theory, Statistical Inference and Statistical Theories of Science, Vol. III, 121-141(1976). Suppose (Ω, C) is a σ -class and f and g are finitely valued measurable functions on (Ω, C) . Suppose m is a probability measure on (Ω, C) .

1. If $f \leq g$ then $\int_{\Omega} f \, dm \leq \int_{\Omega} g \, dm$. The same is valid if f, g have countable number of values which converge to zero.
2. If any of f, g has at most five values and $f+g$ is measurable, then $\int_{\Omega} f \, dm + \int_{\Omega} g \, dm = \int_{\Omega} (f+g) \, dm$.

SOME SEQUENTIAL PROPERTIES

(M. Hušek, Charles University, Sokolovská 83, 18600 Praha 8, Československo), received 29.4. 1980

R. Frič and V. Koutník asked in their contribution on Top. Colloq. in Budapest, August 1978, whether there are sequentially complete completely regular spaces X, Y such that $C^*(X) = C_s^*(X)$ (C_s^* stands for sequentially continuous maps), $C^*(Y) = C_s^*(Y)$, $C^*(X)$ and $C^*(Y)$ are isomorphic and X, Y are not homeomorphic. The space $X = [0,1]^{\omega_1}$ and the Σ_1 -product $Y = \{ \{x_{\alpha}\} \in X \mid \{ \alpha \mid x_{\alpha} \neq 0 \} \leq \omega \}$ solve the problem (the required properties follow from Mazur's results in Fund. Math. 39(1952), 229-238). This pair also solves their Problem 5 whether for $X \subset \beta Y$, $\beta X = \beta Y$ one has $X \subset \sigma Y$ (here σY is the sequential closure of Y in βY). Similarly one can construct disjoint sets X, Y in $[0,1]^{\omega_1}$ such that $C^*(X) =$