

Antonín Kučera

On recursive measure of classes of recursive sets

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 23 (1982), No. 1, 117--121

Persistent URL: <http://dml.cz/dmlcz/106136>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1982

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**ON RECURSIVE MEASURE OF CLASSES OF RECURSIVE SETS**  
**A. KUČERA**

**Abstract:** It is shown that any class of recursive sets  $\{\varphi_{h(n)} : n \in \mathbb{N}\}$  where  $h$  is a function of degree  $\underline{a}$  such that  $\underline{a} \cup \underline{Q} \neq \underline{Q}$  has  $\underline{Q}$ -measure zero ( $\underline{Q}$ -measure is a recursive analogue of the product measure on  $2^{\mathbb{N}}$ ).

**Key words:** Recursive set, recursively enumerable set, degree.

**Classification:** 03D30, 03F60

---

It is known that the recursive sets are not uniformly recursive. C. Jockusch [4, Theorem 9] observed that there is a function  $h$  of degree  $\leq \underline{a}$  such that  $\mathcal{S}_{h(0)}, \mathcal{S}_{h(1)}, \dots$  are precisely the recursive sets iff  $\underline{a} \cup \underline{Q}' \geq \underline{Q}$ . In this paper we prove that any class of recursive sets  $\{\varphi_{h(n)} : n \in \mathbb{N}\}$  where  $h$  is a function of degree  $\leq \underline{a}$  such that  $\underline{a} \cup \underline{Q}' \neq \underline{Q}$  even has  $\underline{Q}$ -measure zero. The concept of  $\underline{Q}$ -measure is an effective analogue of the product measure on  $2^{\mathbb{N}}$ . It was introduced by O. Demuth [1] for constructive real numbers and plays the important role in constructive mathematical analysis (see, e.g., [2]).

Our notation and terminology are standard. In particular we use the letters  $i, j, k, n$  for elements of  $\mathbb{N} = \{0, 1, \dots\}$ . We identify subsets of  $\mathbb{N}$  with their characteristic functions.

A string is a finite sequence of 0's and 1's. Strings may also be viewed as functions from finite initial segments of  $N$  into  $\{0,1\}$ . We use the letters  $\sigma, \tau$  for strings,  $lh(\sigma)$  is the length of  $\sigma$  and  $\sigma * \tau$  is the string which results from concatenating  $\sigma$  and  $\tau$ . A subset  $A$  of  $N$  extends  $\sigma$  ( $A \supseteq \sigma$ ) if the characteristic function of  $A$  extends  $\sigma$ . We assume that the set of all strings is effectively Gödel-numbered so that we can apply notions of recursion theory to strings. For functions  $f, g$  we say that  $f$  dominates  $g$  if  $f(n) \geq g(n)$  for all but finitely many  $n$ . Let  $\varphi_n$  be the  $n$ -th partial recursive function in some standard enumeration of all partial recursive functions.

We shall use the Martin's result [6] that there is a function  $f$  of degree  $\underline{a}$  which dominates every recursive function iff  $\underline{a}' \geq \underline{0}''$ . We shall also use the following straightforward modification of the result.

Lemma: For any degree  $\underline{b}$  and for any class  $\mathcal{A} = \{\varphi_{h(n)} : n \in N\}$  of recursive functions where  $h$  is a function of degree  $\leq \underline{b}'$  there is a function  $f$  of degree  $\leq \underline{b}$  which dominates all functions of  $\mathcal{A}$ .

We shall use a special case of the concept of  $\mathcal{Q}$ -measure (see [1]).

Definition: A class  $\mathcal{A}$  of subsets of  $N$  has  $\mathcal{Q}$ -measure zero if there exist a recursive sequence  $R_0, R_1, \dots$  of r.e. sets of strings and a recursive sequence  $y_0, y_1, \dots$  of constructive real numbers (i.e. recursive reals) such that for every  $n$

1) the real number  $\sum_{\sigma \in R_n} 2^{-lh(\sigma)}$  is equal to  $y_n$  and  $y_n \leq 2^{-n}$ ,

2) for any set  $A$ ,  $A \in \mathcal{R}$ , there is a string  $\sigma$ ,  $\sigma \in R_n$ , such that  $\sigma \subseteq A$ .

It should be noted two important facts in the definition:

i)  $\sum_{\sigma \in R_n} 2^{-lh(\sigma)}$  is required to be equal to a constructive real number for every  $n$ ,

ii)  $y_0, y_1, \dots$  is required to form a recursive sequence.

Zaslavskij and Cejtin [8] proved that the class of all recursive sets has  $\mathcal{Q}$ -measure equal to 1. More information on the role of  $\mathcal{Q}$ -measure and some survey of constructive mathematical analysis can be found in [2].

**Theorem:** If  $\mathfrak{a}$  is a degree such that  $\mathfrak{a} \cup \mathcal{Q}' \neq \mathcal{Q}$  then any class of recursive sets  $\{\varphi_{h(n)} : n \in \mathbb{N}\}$  where  $h$  is a function of degree  $\leq \mathfrak{a}$  has  $\mathcal{Q}$ -measure zero.

**Proof.** It follows from [8] or from [5] that there is a r.e. set  $S_0$  of strings such that

1)  $\sum_{\sigma \in S_0} 2^{-lh(\sigma)}$  is less than  $\frac{1}{2}$ ,

2) for every recursive set  $A$  there exists a string  $\sigma$ ,  $\sigma \in S_0$ , such that  $\sigma \subseteq A$ ,

(i.e. there is a recursive binary tree  $T$  without infinite recursive branches such that the usual product measure on  $2^{\mathbb{N}}$  of the class of all infinite branches of  $T$  is greater than  $\frac{1}{2}$ ). It should be noted that the real number  $\sum_{\sigma \in S_0} 2^{-lh(\sigma)}$  is recursive in  $\mathcal{Q}'$  but it cannot be equal to any constructive real number (see [8]).

Let  $S_0, S_1, \dots$  be a recursive sequence of r.e. sets of strings such that for every  $n$   $S_{n+1} = \{\sigma * \tau : \sigma \in S_n \text{ \& } \tau \in S_0\}$ .

Let  $\{\sigma_{n,k} : k \in \mathbb{N}\}$  be a recursive enumeration of  $S_n$  for every  $n$  (all  $S_n$  are, of course, infinite). It is easy to verify that  $\sum_{\sigma \in S_n} 2^{-lh(\sigma)} < 2^{-(n+1)}$  for all  $n$ .

Further, for any recursive set  $A$  we can effectively find a recursive function  $\alpha$  such that for all  $n$   $A \supseteq \sigma_{n, \alpha(n)}$ . So, let  $g$  be a recursive function such that if  $\varphi_n$  is a recursive set then  $\varphi_n \supseteq \sigma_{k, \varphi_g(n)}(k)$  for all  $k, n$ . Now let  $\underline{a}$  be a degree such that  $\underline{a} \cup \underline{0}' \neq \underline{0}''$  and  $h$  be a function of degree  $\leq \underline{a}$  such that  $\{\varphi_{h(n)} : n \in \mathbb{N}\}$  is a class of recursive sets. We use the function  $g$  described above to form the class of recursive functions  $\mathcal{B} = \{\varphi_{gh(n)} : n \in \mathbb{N}\}$ . The function  $gh$  is obviously of degree  $\leq \underline{a}$ . By the theorem of Friedberg [3] (or [7] § 13.3) there is a degree  $\underline{b}$  such that  $\underline{b}' = \underline{a} \cup \underline{0}'$ . By the lemma there is a function  $f$  of degree  $\leq \underline{b}$  which dominates all functions of the class  $\mathcal{B}$ . Since  $\underline{b}' \neq \underline{0}''$ , there is a recursive function  $\sigma'$  which  $f$  fails to dominate. Thus, for all  $n$ ,  $\varphi_{gh(n)}(k) \leq \sigma'(k)$  for infinitely many  $k$ . By the properties of  $g$  we have  $\varphi_{h(n)} \supseteq \sigma_{k, \varphi_{gh(n)}}(k)$  for all  $k, n$ . Let  $R_0, R_1, \dots$  be a recursive sequence of r.e. sets of strings such that for every  $n$   $R_n = \{\sigma_{k,j} : k \geq n \text{ \& } j \leq \sigma'(k)\}$ . It follows that for all  $i, n$  there is a string  $\sigma' \in R_n$  such that  $\varphi_{h(i)} \supseteq \sigma'$ . Further, it is easy to construct a recursive sequence of constructive real numbers  $y_0, y_1, \dots$  such that for all  $n$   $\sum_{\sigma \in R_n} 2^{-lh(\sigma)}$  is equal to  $y_n$  and  $y_n \leq 2^{-n}$ . Thus, the class  $\{\varphi_{h(n)} : n \in \mathbb{N}\}$  has  $\underline{0}$ -measure zero.

### R e f e r e n c e s

- [1] O. DEMUTH: The Lebesgue measurability of sets in constructive mathematics, Comment. Math. Univ. Carolinae 10(1969), 463-492 (in Russian).
- [2] O. DEMUTH and A. KUČERA: Remarks on constructive mathematical analysis, Logic Colloquium '78 (Boffa, van Dalen, McAloon editors), North-Holland, Amsterdam, 1979, 81-129.
- [3] R.M. FRIEDBERG: A criterion for completeness of degrees of unsolvability, J. Symbol. Logic 22(1957), 159-160.
- [4] C.G. JOCKUSCH: Degrees in which the recursive sets are uniformly recursive, Canad. J. Math. 24(1972), 1092-1099.
- [5] C.G. JOCKUSCH and R.I. SOARE:  $\Pi_1^0$  classes and degrees of theories, Trans. Amer. Math. Soc. 173(1972), 33-56.
- [6] D.A. MARTIN: Classes of recursively enumerable sets and degrees of unsolvability, Z. Math. Logik Grundlagen Math. 12(1966), 295-310.
- [7] H. ROGERS: Theory of recursive functions and effective computability, McGraw-Hill, New York, 1967.
- [8] I.D. ZASLAVSKIJ and G.S. CEJTIN: On singular coverings and related properties of constructive functions, Trudy Mat. Inst. Steklov. 67(1962), 458-502; English transl. Amer. Math. Soc. Transl.(2)98(1971), 41-89.

Matematicko-fyzikální fakulta, Universita Karlova,  
Malostranské nám. 25, Praha 1, Czechoslovakia

(Obtatum 9.9. 1981)