

Piotr Biler

Asymptotic behaviour in time of solutions to some equations generalizing the Korteweg-de Vries-Burgers equation

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 26 (1985), No. 1, 177--180

Persistent URL: <http://dml.cz/dmlcz/106355>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1985

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**ASYMPTOTIC BEHAVIOUR IN TIME OF SOLUTIONS TO SOME  
EQUATIONS GENERALIZING THE KORTEWEG-DE VRIES-BURGERS  
EQUATION**  
Piotr BILER

**Abstract:** We summarize the results of a more detailed paper concerning the decay estimates for the solutions to equations describing the propagation of nonlinear waves which generalize the Korteweg-de Vries and Burgers equation.

**Key words:** Generalized Korteweg-de Vries and Burgers equation, propagation of nonlinear waves, decay in time of solutions.

Classification: 35Q20, 35B40

-----

J.C. Saut has considered in [2] a class of model equations describing propagation of nonlinear waves which generalize the Korteweg-de Vries and Burgers equations. He has proved several theorems on the existence, uniqueness and regularity of solutions of the Cauchy problem for equations of the type

$$u_t + \sum_{i=1}^m \frac{\partial}{\partial x_i} [f(t,u) + \sigma H(x,u)] + \varepsilon Bu = g$$

where  $x \in \mathbb{R}^n$ ,  $u = u(x,t)$  is a real function,  $H$ ,  $B$  are the (real) pseudodifferential operators describing dispersive and dissipative properties of the medium and  $f$  is a polynomially bounded function of  $u$ .

We prove, using the ideas of the papers [3],[4], some

-----  
This paper was presented in written form on the International Spring School on Evolution Equations, Dobřichovice by Prague, May 21-25, 1984.

theorems on the decay in time of the solutions (in  $L^p$  norms) to one-dimensional equations of more special structure

$$(*) \quad u_t + f(u)_x + \sigma(Hu)_x + \varepsilon D^s u = 0$$

where  $D^s u(x) = \int |\xi|^s \hat{u}(\xi) e^{ix\xi} d\xi$ ,  $s \in \mathbb{R}^+$ ,

$Hu(x) = \int p(\xi) \hat{u}(\xi) e^{ix\xi} d\xi$  with an even positive symbol  $p$  of polynomial growth.

In the proofs we use energy inequalities for  $(*)$ , interpolation of Sobolev spaces and elementary properties of the fundamental solution of the linearized equation.

Theorem 1. ( $\sigma \neq 0$ ,  $\varepsilon > 0$ ; dispersion and dissipation effects are included)

- a) If  $|f'(u)| \leq C(|u|^p + 1)$  for some  $p < 2(s-1)$ ,  $s \geq 2$ ,  $u_0 \in H^s$ , then  $\lim_{t \rightarrow \infty} |u(t)|_\infty = 0$ .
- b) The optimal decay rate (identical as for the linearized equation) is obtained assuming that  $f$  is sufficiently flat at the origin:

If also  $|f'(u)| \leq C|u|^q$  for some  $q > 2s + 1$  in a neighbourhood of 0 and  $u_0 \in L^1$ , then

$$|u(t)|_\infty = O((1+t)^{-1/s}) \text{ and } |u(t)|_2 = O((1+t)^{-1/2s}).$$

Theorem 2. ( $\varepsilon > 0$ ,  $\sigma = 0$ ; pure dissipative case)

- a) The assumption in Th. 1a) plus  $u_0 \in L^1$  implies that  $|u(t)|_2 = O((1+t)^{-1/2s})$ .
- b) If  $|f'(u)| \leq C|u|^q$  for some  $q \geq 2s - 1$  and small  $|u|$ , then  $|u(t)|_\infty = O((1+t)^{-1/s})$ .

The pure dispersion case ( $\varepsilon = 0$ ,  $\sigma \neq 0$ ) leads to energetically neutral equations:  $|u(t)|_2 = \text{const}$ . They can have special wave-like solutions - solitons - which do not decay when

$t$  tends to infinity. Since now it is more difficult to estimate the fundamental solution of the linearized equation, we restrict our attention to the case of homogeneous symbols  $p(\xi) = |\xi|^{r-1}$ ,  $r \geq 3$ , and we consider only small solutions of  $(*)$  (with initial conditions small enough to do not support the solitons).

Theorem 3.

- a) If  $|f'(u)| \leq C|u|^q$ ,  $q > r + 1$  in a neighbourhood of  $u = 0$  and  $\|u_0\|_1 + \|u_0\|_{(r-1)/2}$  is small then  
 $|u(t)|_\infty = O((1 + |t|)^{-1/r})$  for  $|t| \rightarrow \infty$ .
- b) A better (than obtained by a simple interpolation) result on the decay of  $L^p$  norms of the solution is:  
 If  $q > (r + (r^2 + 4r)^{1/2})/2$  then  
 $|u(t)|_{2(q+1)} = O((1+t)^{-(1-1/(q+1))/r})$ .

The space-periodic solutions of  $(*)$  in the case of dissipation ( $\varepsilon > 0$ ) decay exponentially when  $t$  tends to infinity. Similarly as for the Navier-Stokes equations (cf. [1]) the solutions are asymptotically equal to solutions of the linearized equation. Namely we can prove the following

Theorem 4.

- a) If  $|f'(u)| \leq C(|u|^p + |u|)$  for some  $p < 2(s-1)$ ,  $s \geq 2$ , then  
 $\Lambda = \lim_{t \rightarrow \infty} (D^s u(t), u(t)) / |u(t)|_2^2$  exists and  $\Lambda = \Lambda(u_0)$   
 is an eigenvalue of  $D^s$ .

Moreover

- b) If  $\sigma = 0$  then  $\lim_{t \rightarrow \infty} e^{\Lambda t} u(t)$  exists and it is a non-zero eigenfunction of  $D^s$  corresponding to  $\Lambda$ .

#### R e f e r e n c e s

- [1] C. FOIAS, J.C. SAUT: Asymptotic behavior, as  $t \rightarrow +\infty$ ,

of solutions of Navier-Stokes equations and nonlinear spectral manifolds, Indiana Univ. Math. J. (to appear).

- [2] J.C. SAUT: Sur quelques généralisations de l'équation de Korteweg-de Vries, J. Math. pures appl. 58(1979), 21-61.
- [3] M.E. SCHONBEK: Decay of solutions to parabolic conservation laws, Comm. P.D.E. 5(1980), 449-473.
- [4] W.A. STRAUSS: Nonlinear scattering at low energy, J. Funct. Anal. 41(1981), 110-133.

Institute of Mathematics, University of Wrocław, pl. Grunwaldzki 2/4, 50-385 Wrocław, Poland

(Oblatum 25.5. 1984)