

Aleš Drápal; Tomáš Kepka
Group distances of Latin squares

Commentationes Mathematicae Universitatis Carolinae, Vol. 26 (1985), No. 2, 275--283

Persistent URL: <http://dml.cz/dmlcz/106367>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1985

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

GROUP DISTANCES OF LATIN SQUARES
Aleš DRÁPAL and Tomáš KEPKA

Abstract: Some results concerning the distances between the tables of finite groups and latin squares are proved.

Key words: Group, latin square.

Classification: 05B15

For an integer $n \geq 2$, let $gdist(n)$ denote the least non-zero number of changes in the Cayley table of an n -element group to obtain another latin square. These numbers play an important rôle in the problem concerning the largest possible number of associative triples of elements in finite non-associative quasigroups (see [2]). The purpose of this short note is to develop a technique which might be useful in finding some lower bounds for the numbers $gdist(n)$.

1. Preliminaries. Throughout this note, the terminology, notation, etc., of [3] is used.

Recall that \mathcal{R} denotes the category of reduced partial groupoids and \mathcal{T} the full subcategory of \mathcal{R} consisting of reduced balanced cancellative partial groupoids.

A homomorphism f of a partial groupoid K into a partial groupoid L is called complete if for all $(x,y) \in M(L)$ such that $x,y,xy \in f(K)$ there exists a pair $(a,b) \in M(K)$ with $f(a) = x$ and

$f(b) = y$ (then $f(ab) = xy$). Obviously, every strong homomorphism is complete.

A partial groupoid L is called a (complete, strong) partial subgroupoid of a partial groupoid K if $L \subseteq K$ and this inclusion is a (complete, strong) homomorphism.

Let $K \in \mathcal{R}$. We shall say that K is trivial if $\text{card } B(K) = \text{card } C(K) = \text{card } D(K) = 1$. In this case, $1 \leq \text{card } K \leq 3$ and $\text{card } K = 3$, provided K is balanced. A homomorphism f of K into $L \in \mathcal{R}$ is called trivial if $f[K]$ is a trivial partial groupoid. In this case, $f[K]$ is a strong partial subgroupoid of L , provided L is balanced.

Let $K \in \mathcal{R}$ and $d \in K$. Put $r(d) = r(K, d) = \text{card } \{a, b, c\}; a, b, c \in K, ab = c, d \in \{a, b, c\}$. Since K is reduced, $r(d) \geq 1$.

Let $K, L \in \mathcal{R}$. We shall say that K is an immediate (strongly) open extension of L if L is a (strong) complete partial subgroupoid of K and $r(K, d) = 1$ for every $d \in K - L$. Further, we shall say that K is an (strongly) open extension of L if there exists a finite sequence $K_0 \subseteq K_1 \subseteq \dots \subseteq K_n$ such that $n \geq 1, K_0 = L, K_n = K$ and K_{i+1} is an immediate (strongly) open extension of K_i for each $0 \leq i < n$.

A partial groupoid $K \in \mathcal{T}$ is called (strongly) open if it is non-trivial and it is a (strongly) open extension of a trivial partial subgroupoid $L \in \mathcal{T}$.

1.1. Lemma. Let $K \in \mathcal{T}$ and let $a, b, c \in K$ be such that $ab = c$. Then $L = \{a, b, c\}$ is a three-element strong partial subgroupoid of K and L is a trivial partial groupoid.

Proof. Obvious.

1.2. Lemma. Let $K \in \mathcal{T}$ be such that $m(K) \leq 3$. Then:
(i) $r(a) = 1$ for at least one $a \in A(K)$.

(ii) K is strongly open, provided it is non-trivial.

Proof. Easy.

1.3. Lemma. Let $K \in \mathcal{J}$ be such that $m(K) = 4$. Then exactly one of the following three cases takes place:

- (i) $r(a) = 1$ for at least one $a \in A(K)$ and K is strongly open.
- (ii) $r(a) \geq 2$ for every $a \in A(K)$, $r(d) = 1$ for at least one $d \in D(K)$, K is open and K is not strongly open.
- (iii) $r(a) \geq 2$ for every $a \in A(K)$, $r(d) \geq 2$ for every $d \in D(K)$, K is not open and $H(K)$ is a cyclic group of order 2.

Proof. Easy.

1.4. Lemma. Let $K, L \in \mathcal{J}$ be such that K is an open extension of L and let f be a homomorphism of L into a division groupoid G . Then f can be extended to a homomorphism of K into G .

Proof. We can assume that K is an immediate open extension of L . However, then the result is clear.

1.5. Lemma. Let $K \in \mathcal{J}$ be open and let G be a non-trivial division groupoid. Then there exists at least one non-trivial homomorphism of K into G .

Proof. If $m(K) = 2$ then the result is obvious. Suppose that $m(K) \geq 3$. Then there is a strong partial subgroupoid L of K such that $m(L) = 2$ and K is an open extension of L . Now, the result follows from 1.4.

1.6. Lemma. Let f be a homomorphism of a partial groupoid K into a group G and let $(a, b) \in N(K)$. Then there exists a homomorphism g of K into G such that $g(a) = g(b) = g(ab) = 1$. Moreover, g is non-trivial, provided f is so.

Proof. Put $g(a) = f(a)^{-1}f(e)$, $g(d) = f(d)f(b)^{-1}$ and $g(e) = f(a)^{-1}f(e)f(b)^{-1}$ for all $a \in B(K)$, $d \in C(K)$ and $e \in D(K)$.

1.7. Lemma. Let f be a non-trivial homomorphism of a partial groupoid $K \in \mathcal{T}$ into a group G and let H be a normal subgroup of G . Then there exists either a non-trivial homomorphism of K into H or a non-trivial homomorphism of K into G/H .

Proof. With respect to 1.6, we can assume that 1 is contained in all the sets $f(B(K))$, $f(C(K))$, $f(D(K))$. Denote by g the natural homomorphism of G onto G/H . If gf is a trivial homomorphism then $f(K) \subseteq H$.

1.8. Lemma. Let $K \in \mathcal{T}$ and G be a group. Then there exists a non-trivial homomorphism of K into G iff there exists a non-trivial homomorphism of $H(K)$ into G .

Proof. Choose $x = (a, b) \in M(K)$ and consider the congruence $s = s_x$ by [3, Lemma 2.2], the natural homomorphism q of K onto $L = K/s$, the isomorphism h of $G(L)$ onto $H(K)$ by [3, Lemma 5.2] and the modificative homomorphism g of L into $G(L)$ by [3, Proposition 3.1]. Now, let f be a non-trivial homomorphism of K into G . With regard to 1.6, we can assume that $f(a) = f(b) = 1$. Then $s \subseteq \ker f$, and hence $f = kq$, k being a non-trivial homomorphism of L into G . We have $k = pg$ for a homomorphism p of $G(L)$ into G and ph^{-1} is a non-trivial homomorphism of $H(K)$ into G . Conversely, let k be a non-trivial homomorphism of $H(K)$ into G . Put $f = khgq$. Then f is a homomorphism of K into G and $f(a) = f(b) = f(ab) = 1$. On the other hand, the group $k(H(K))$ is generated by $f(K)$ and it is non-trivial. Consequently, f is non-trivial.

1.9. Lemma. Let $K \in \mathcal{T}$ be non-trivial, $ab = c$ for some $a, b, c \in K$ and let G be a non-trivial division groupoid. Suppose

that either $r(a) = r(b) = 1$ or $r(a) = r(c) = 1$ or $r(b) = r(c) = 1$. Then there exists at least one non-trivial homomorphism of K into G .

Proof. It is divided into several parts.

(i) $r(a) = r(b) = r(c) = 1$. Let $x, y \in G$ be such that $x \neq y$. Define a mapping f of K into G by $f(u) = f(v) = x$, $f(w) = xx$, $f(a) = y$, $f(b) = y$ and $f(c) = yy$ for all $u \in B(K)$, $v \in C(K)$, $w \in D(K)$, $u \neq a$, $v \neq b$ and $w \neq d$. Then f is a non-trivial homomorphism of K into G .

(ii) $r(a) = r(b) = 1$ and $r(c) \geq 2$. Let $x, y \in G$ be such that $x \neq y$. There exists $z \in G$ such that $yz = xx$. Now, define f by $f(u) = f(v) = x$, $f(w) = xx$, $f(a) = y$, $f(b) = z$ for all $u \in B(K)$, $v \in C(K)$ and $w \in D(K)$, $u \neq a$, $v \neq b$.

(iii) $r(a) = r(c) = 1$ and $r(b) \geq 2$. Let $x, y \in G$, $x \neq y$. Define f by $f(u) = f(v) = x$, $f(w) = xx$, $f(a) = y$ and $f(c) = yx$ for all $u \in B(K)$, $v \in C(K)$ and $w \in D(K)$, $u \neq a$, $w \neq c$.

(iv) $r(b) = r(c) = 1$ and $r(a) \geq 2$. In this case, we can proceed similarly as in (iii).

2. Homomorphisms into groups. Let G be a non-trivial group. A partial groupoid K is said to be G -flat (or only flat) if every homomorphism of K into G is trivial.

Let $n \geq 2$ be an integer. We denote by $z(n) = z(G, n)$ the minimum of all $m(K)$ where $K \in \mathcal{T}$ is flat and there exists a non-trivial homomorphism of K into an n -element group.

2.1. Lemma. Let $K \in \mathcal{T}$ be flat.

(i) If f is a homomorphism of K into $L \in \mathcal{T}$ then $f[K]$ is flat.

(ii) K is not open.

(iii) If K is an open extension of $L \in \mathcal{T}$ then L is flat.

Proof. Use 1.4 and 1.5.

2.2. Lemma. Suppose that G is a torsionfree group and let $K \in \mathcal{J}$ be such that $H(K)$ is a torsion group. Then K is flat.

Proof. This follows immediately from 1.8.

2.3. Lemma. Let $K \in \mathcal{J}$ be non-trivial and flat and let $a, b, c \in K$ be such that $ab = c$. Then either $r(a) \geq 2$, $r(b) \geq 2$ or $r(a) \geq 2$, $r(c) \geq 2$ or $r(b) \geq 2$, $r(c) \geq 2$.

Proof. This follows immediately from 1.9.

2.4. Proposition. Let $n \geq 2$ be an integer and let $K \in \mathcal{J}$ be a partial groupoid such that $m(K) = z(n)$. Suppose that there exists a non-trivial homomorphism f of K into an n -element group H . Then $r(a) \geq 2$ for every $a \in K$.

Proof. Assume, on the contrary, that $r(a) = 1$ for some $a \in K$. There are three different elements $x, y, z \in K$ such that $xy = z$ and $a \in \{x, y, z\}$. Now, with respect to 2.3, the following cases can arise:

(i) $r(x) = 1$, $r(y) \geq 2$ and $r(z) \geq 2$. Put $L = K - \{x\}$. Then $L \in \mathcal{J}$, L is a strong partial subgroupoid of K , $m(L) = m(K) - 1$, K is an open extension of L and L is flat. According to 1.7, we can assume that $1 \in f(B(L)) \cap f(C(L)) \cap f(D(L))$. Since $f|_L$ is trivial, $f(L) = 1$. Then $f(x) = f(x)1 = f(x)f(y) = f(xy) = f(z) = 1$ and f is trivial, a contradiction.

(ii) $r(x) \geq 2$, $r(y) = 1$ and $r(z) \geq 2$. We can proceed similarly as in (i).

(iii) $r(x) \geq 2$, $r(y) \geq 2$ and $r(z) = 1$. Again, we can proceed similarly as in (i) (in this case, $L = K - \{z\}$ is a complete partial subgroupoid of K).

2.5. Lemma. Suppose that G is a torsionfree group. Then $4 \leq z(n) \leq 2n$ for every $n \geq 2$.

Proof. By 2.1(ii) and 1.2(ii), $m(K) \geq 4$ for every non-trivial flat partial groupoid $K \in \mathcal{T}$. Hence $4 \leq z(n)$. Further, consider the partial groupoid $Z = Z(n, \circ)$ defined in [4, § 7]. Then $m(Z) = 2n$ and $H(Z)$ is a cyclic group of order n . Consequently, Z is flat by 2.2 and $z(n) \leq 2n$.

2.6. Proposition. Suppose that G is a torsionfree group. Then for every $n \geq 2$, $z(n) = 4$ iff n is even.

Proof. First, let $z(n) = 4$. Then there are $K \in \mathcal{T}$ and a group H such that K is flat, $m(K) = 4$, H contains just n elements and there exists a non-trivial homomorphism of K into H . The partial groupoid K is not open, and so $H(K)$ is a two-element group by 1.3(iii). By 1.8, there is a non-trivial homomorphism of $H(K)$ into H . In particular, n is even. Now, let n be even. Then we can proceed conversely.

2.7. Proposition. Let $n \geq 3$ be odd. Then $z(n)$ is equal to the minimum of all $z(p)$, p being a prime dividing n .

Proof. The result follows from 1.7 and the fact that n is prime, provided there is a simple group of order n .

3. Homomorphisms into ordered partial groupoids. In this section, let G be a cancellative reduced partial groupoid linearly ordered by an ordering \leq , i.e. \leq is a linear ordering defined on G and $ab \leq cd$ whenever $(a,b), (c,d) \in M(G)$, $a \leq c$ and $b \leq d$.

3.1. Lemma. Let $I = (K(\circ), K(*))$ be a couple of finite simple companions. Then every homomorphism of $K(\circ)$ into G is trivial.

Proof. Let f be a homomorphism of $K = K(\circ)$ into G . There is an element $x \in f(B(K))$ such that $y \leq x$ for any $y \in f(D(K))$. Put

$N = \{(a,b) \in M(K); f(a \circ b) = x\}$ and define a relation r on N by $((a,b), (c,d)) \in r$ iff $f(a) = f(c)$ and $f(b) = f(d)$. Since G is cancellative, each of the two equalities implies the other. Obviously, r is an equivalence and we denote by N_1, \dots, N_k the blocks of r . Without loss of generality, we can assume that $f(a_1) < f(a_2) < \dots < f(a_k)$, $(a_i, b_i) \in N_i$. Now, we are going to prove that N_1 is an admissible subset of $M(K)$ in the sense of [4, § 5]. Let $(a,b) \in N_1$. Put $P = \{(u,v) \in M(K); f(u \ast v) = x, f(u) = f(a)\}$, $Q = \{(u,v) \in M(K); f(u \ast v) = x, f(v) = f(b)\}$. The rest of the proof is divided into several parts.

(i) If $(u,v) \in P$ and $u \ast v = u \circ w$ then $f(u \circ w) = x$, $(u,w) \in N_1$. Conversely, if $(u,w) \in N_1$ and $u \circ w = u \ast v$ then $(u,v) \in P$. Hence we have injective mappings of P into N_1 and of N_1 into P , so that $\text{card } P = \text{card } N_1$.

(ii) Similarly as in (i) we can show that $\text{card } Q = \text{card } N_1$.

(iii) Let $(u,v) \in Q$. We have $u \ast v = w \circ v = u \circ z$, $f(a)f(b) = x = f(u \ast v) = f(u \circ z) = f(u)f(z)$, so that $(u,z) \in N$ and $f(a) \leq f(u)$. On the other hand, $x = f(a)f(b) \leq f(u)f(v)$, since $f(b) = f(v)$, hence $x = f(u)f(v) = f(u \circ v)$, $f(u) = f(a)$ and $(u,v) \in N_1$. We have proved that $Q \subseteq N_1$. Now, it is easy to see that $Q \subseteq P$.

(iv) By (i), (ii) and (iii), we have $P = Q = N_1$. Consequently, N_1 is an admissible subset of $M(K)$. Since the couple I is simple, $N_1 = M(K)$ and f is trivial.

3.2. Corollary. Let $K \in \mathcal{J}$ be a primary groupoid and let G be a linearly ordered non-trivial group. Then K is G -flat.

4. The main result

4.1. Proposition. Let G be a linearly ordered non-trivial group. Then, for every $n \geq 2$, $z(G,n) \leq \text{gdist}(n)$.

Proof. The result is an immediate consequence of 3.2 and [3, Proposition 7.5].

4.2. Proposition. Let G be a linearly ordered non-trivial group and $n \geq 2$ and integer. Then there is a prime p dividing n such that $z(G, p) \leq \text{gdist}(n)$.

Proof. The result follows from 4.1 and 2.7.

R e f e r e n c e s

- [1] J. DÉNES and A.D. KEEDWELL: Latin Squares and Their Applications, Akadémiai Kiadó, Budapest, 1974.
- [2] A. DRÁPAL: On quasigroups rich in associative triples, Discrete Math. 44(1983), 251-165.
- [3] A. DRÁPAL and T. KEPKA: Group modifications of some partial groupoids, Annals of Discr. Math. 18(1983), 319-332.
- [4] A. DRÁPAL and T. KEPKA: Exchangeable partial groupoids I, Acta Univ. Carolinae 24(1983), 57-72.

Matematicko-fyzikální fakulta, Univerzita Karlova, Sokolovská 83,
186 00 Praha 8, Czechoslovakia

(Oblatum 8.10. 1984)