

Alessandro Caterino

Special lattices of compactifications

Commentationes Mathematicae Universitatis Carolinae, Vol. 26 (1985), No. 3, 515--523

Persistent URL: <http://dml.cz/dmlcz/106390>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1985

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

SPECIAL LATTICES OF COMPACTIFICATIONS
Alessandro CATERINO

Abstract. Given any compactification αX of a Tychonoff space X , let $f_\alpha : \beta X \rightarrow \alpha X$ denote the canonical quotient map from the Stone-Cech compactification of X onto αX . It is known that the complete upper semi-lattice $K(X)$ of all compactifications of X becomes a lattice whenever the set $F^*(\alpha X) = \{f_\alpha^{-1}(p) : |f_\alpha^{-1}(p)| > 1\}$ is finite for all $\alpha X \in K(X)$. In this paper we give some necessary and sufficient conditions, in terms of X and $\beta X - X$, for $F^*(\alpha X)$ to be finite for all $\alpha X \in K(X)$.

AMS(1980) Subject Class. Primary: 54D35, 54D40. Secondary: 54C45, 54G05, 54G10.

Key-words. Lattice, compactification, remainder, C^* -embedded, cf -space, P -space, F -space.

Introduction. Let X be a Tychonoff space. Denote by $K(X)$ the family of T_2 -compactifications of X . Two compactifications αX and γX are considered equivalent if there is a homeomorphism between αX and γX , which leaves X pointwise fixed; we do not distinguish between equivalent compactifications in $K(X)$. $K(X)$ is partially ordered by the relation: $\alpha_1 X \leq \alpha_2 X$ if there is a continuous map from $\alpha_2 X$ onto $\alpha_1 X$, which leaves X pointwise fixed. It is known that $K(X)$ is always a complete upper semi-lattice and that it is a complete lower semi-lattice (hence a complete lattice) iff X is locally compact (cf. [M]).

In general $K(X)$ is not a lattice, for example when X is first countable but not locally compact (cf. [FV]).

In this paper we study questions related to the problem of when $K(X)$ is a lattice.

In the following, we will use the term compactification instead of T_2 -compactification.

If $\alpha X \in K(X)$, βX will denote the Stone-Cech compactification of X and $f_\alpha : \beta X \longrightarrow \alpha X$ the canonical quotient map. Define the β -family of αX to be $F(\alpha X) = \{f_\alpha^{-1}(p) : p \in \alpha X - X\}$ and set $F^*(\alpha X) = \{F \in F(\alpha X) : |F| > 1\}$. Recall that any family of bounded continuous functions, $S \subset C^*(X)$, which separates points from closed sets, generates a compactification $\alpha_S X = \overline{e_S(X)}$, where $e_S : X \longrightarrow \prod_{f \in S} K_f$, $K_f = \overline{f(X)}$, is the topological embedding defined by $e_S(x) = \{f(x)\}_{f \in S}$.

Moreover, observe that, if $F_1, \dots, F_n \subset \beta X - X$ are disjoint compact sets with $|F_i| > 1$, then the quotient space αX of βX , obtained by shrinking each compact F_i to a point, is a compactification of X and one has $F^*(\alpha X) = \{F_1, \dots, F_n\}$. Obviously αX coincides with the compactification generated by $S = \{f \in C^*(X) : f|_{F_i} \text{ is constant } \forall i=1, \dots, n\}$, where f^β is the extension of f to βX .

Some topological spaces have the property that all their compactifications are obtained as previously described, that is $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$. In this case, it is easy to prove that $K(X)$ is a lattice. In fact, if αX and γX are compactifications of X , then $\alpha X \wedge \gamma X$ is generated by the family of continuous functions

$$\{f \in C^*(X) : f^\beta|_F \text{ is constant } \forall F \in F^*(\alpha X) \cup F^*(\gamma X)\}.$$

In ([C], th.5.6; see also [FV], proof of th. 1) it is pointed out that if $\beta X - X$ is discrete and C^* -embedded in βX , then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$. More generally, one obtains the same result when $\beta X - X$ is a P -space and $Cl_{\beta X}^1(\beta X - X)$ is an F -space (cf. [U]). Recall that a P -space is a space in which every cozero-set is C -embedded and an F -space is a space in which every cozero-set is C^* -embedded (for equivalent definitions cf. 4J, 14.25, 14.29, 14N in [GJ]).

Among the results of the present paper is the following proposition generalizing the above mentioned results: if $\beta X - X$ is a cf -space (that is a space whose compact sets are finite) and every countable discrete subset of $\beta X - X$

is C^* -embedded in βX , then $F_\alpha^*(\alpha X)$ is finite for all $\alpha X \in K(X)$.

The same conclusion is achieved if the following three conditions are satisfied:

a) $\beta X - X$ is C^* -embedded in βX , b) $\beta X - X$ is countably normal (we say that a space is countably normal if any two disjoint countable closed sets are completely separated), and c) every infinite subset of $\beta X - X$ contains an infinite discrete and closed subset of $\beta X - X$.

An application of the last proposition is obtained when $\beta X - X$ is an MI-space (that is, dense in itself and whose dense subsets are open), countably normal and C^* -embedded in βX .

Moreover, we prove that $\beta X - X$ is a cf-space if $F_\alpha^*(\alpha X)$ is finite for all $\alpha X \in K(X)$ and, under additional hypotheses on X or $\beta X - X$, we give some equivalent conditions for $F_\alpha^*(\alpha X)$ to be finite for all $\alpha X \in K(X)$.

We will denote with N and R the sets of natural numbers and real numbers, respectively.

1. All spaces we deal with are Tychonoff. Let αX be a compactification of a space X and let $f_\alpha : \beta X \longrightarrow \alpha X$ be the canonical quotient map. A subset A of βX is said to be saturated (relative to f_α) when $A = f_\alpha^{-1}(f_\alpha(A))$. Given $F \subset A \subset \beta X$, where $F = f_\alpha^{-1}(p)$ with $p \in \alpha X$ and A is an open subset of βX , then, since f_α is a closed map, there exists an open saturated subset U of βX such that $F \subset U \subset A$.

LEMMA 1. Let αX be a compactification of X , $G = \{F_\lambda\}_{\lambda \in \Lambda} \subset C.F^* = F^*(\alpha X)$ and let $A = \{x_\lambda\}_{\lambda \in \Lambda}$ with $x_\lambda \in F_\lambda$ for every $\lambda \in \Lambda$. Then

$$\left(Cl_{\beta X} \left(\bigcup_{\lambda \in \Lambda} F_\lambda \right) \right) - S = \left(Cl_{\beta X} A \right) - S$$

where $S = \bigcup_{F \in F^*} F$.

Proof.

Obviously $\left(Cl_{\beta X} A \right) - S \subset \left(Cl_{\beta X} \left(\bigcup_{\lambda \in \Lambda} F_\lambda \right) \right) - S$. Conversely if $x \notin \left(Cl_{\beta X} A \right) - S$ then we can suppose, without loss of generality, that $x \notin S$. Let $V \subset \beta X$ be an open set such that $x \in V$, $V \cap A = \emptyset$. Then there exists an open saturated subset U of βX with $x \in U \subset V$. It is clear that $U \cap F_\lambda = \emptyset$ for all $\lambda \in \Lambda$, since U is saturated and $x_\lambda \notin U$ for all $\lambda \in \Lambda$. Thus $x \notin \left(Cl_{\beta X} \left(\bigcup_{\lambda \in \Lambda} F_\lambda \right) \right) - S$.

COROLLARY 2. Let $G = \{F_\lambda\}_{\lambda \in \Lambda} \subset F^*(\alpha X)$ and let $A = \{x_\lambda\}_{\lambda \in \Lambda}$, $B = \{y_\lambda\}_{\lambda \in \Lambda}$, $x_\lambda, y_\lambda \in F_\lambda$ for every $\lambda \in \Lambda$. Then

$$(Cl_{\beta X} A) - S = (Cl_{\beta X} B) - S .$$

PROPOSITION 3. If $\beta X - X$ is a cf-space and every countable discrete subset of $\beta X - X$ is C^* -embedded in βX , then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$.

Proof.

Suppose that, for some $\alpha X \in K(X)$, $F^*(\alpha X)$ is infinite. It follows that $D = f_\alpha(F^*(\alpha X))$ is infinite. If $T = \{p_n\}$ is countably infinite discrete subset of D , set $F_n = f_\alpha^{-1}(p_n)$ for every $n \in \mathbb{N}$, and let $A = \{x_n\}$, $B = \{y_n\}$ where $x_n, y_n \in F_n$, $x_n \neq y_n$. Since T is discrete in $\alpha X - X$, it follows that $S = A \cup B$ is a discrete subset of $\beta X - X$. In fact, for every $n \in \mathbb{N}$, there is an open set V_n of $\alpha X - X$ such that $p_m \in V_n$ iff $m = n$.

Then setting $U_n = f_\alpha^{-1}(V_n) \cap F_n$, one has $U_n \cap F_k = \emptyset$ for every $k \neq n$, otherwise $p_k \in V_n$.

By assumption S is C^* -embedded in βX , hence A and B , which are completely separated in S , are completely separated in βX . Thus we have

$Cl_{\beta X} A \cap Cl_{\beta X} B = \emptyset$; moreover it follows from Corollary 2 that $Cl_{\beta X} A \cap X = Cl_{\beta X} B \cap X$. We conclude that both A and B have no cluster points in X , hence $Cl_{\beta X} A \subset \beta X - X$. This is a contradiction, because $\beta X - X$ was supposed to be a cf-space.

As a consequence of the above proposition we obtain the known results :

COROLLARY 4. ([FV]) If $\beta X - X$ is discrete and C^* -embedded in βX , then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$.

COROLLARY 5. ([U]) If $\beta X - X$ is a P-space and $Cl_{\beta X}(\beta X - X)$ is an F-space, then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$.

Proof. Every countable subset of a P-space is closed and discrete, so every P-space is a cf-space (cf. 4K in [GJ]). Also every countable subset of an F-space is C^* -embedded (cf. 14N in [GJ]). Then apply the Tietze-Urysohn theorem.

We give now another sufficient condition for $F^*(\alpha X)$ to be finite for

all $\alpha X \in K(X)$.

PROPOSITION 6. Let X be a space such that :

- a) $\beta X - X$ is C^* -embedded in βX
- b) $\beta X - X$ is countably normal
- c) every infinite subset of $\beta X - X$ contains an infinite discrete and closed subset of $\beta X - X$.

Then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$

Proof.

First observe that, if $T \subset \beta X - X$ is infinite, then there exists a countably infinite subset of T , which is closed and discrete.

Now suppose that there is an $\alpha X \in K(X)$ such that $F^*(\alpha X)$ is infinite. Then there exists a countably infinite set $A' \subset \bigcup \{F : F \in F^*(\alpha X)\}$, which is closed and discrete in $\beta X - X$. Since every $F \in F^*(\alpha X)$ is compact, then $A' \cap F$ is finite for all $F \in F^*(\alpha X)$. Thus, one can suppose that A' meets every $F \in F^*(\alpha X)$ in at most one point. If $A' = \{x_n\}$, let $F_n = f_\alpha^{-1}(f_\alpha(x_n))$ for every $n \in \mathbb{N}$. Then consider a countably infinite set $B \subset \bigcup_{n \in \mathbb{N}} F_n - \{x_n\}$ closed and discrete in $\beta X - X$. As above, we can suppose

that $|B \cap F_n| \leq 1$ for all $n \in \mathbb{N}$. If $B = \{y_{n_j}\}$ with $y_{n_j} \in F_{n_j}$, let $A = \{x_{n_j}\}$. By arguments similar to those in Proposition 3, we obtain that A has no cluster points in X and so it is closed in βX . This is a contradiction since A is an infinite discrete set.

COROLLARY 7. Let $\beta X - X$ be an MI-space, countably normal and C^* -embedded in βX , then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$.

Proof.

It is easy to prove that, every infinite subset of a Hausdorff MI-space Y contains a countably infinite closed and discrete subset. In fact, if $T \subset Y$ is infinite, consider a copy N of \mathbb{N} in T . N has no interior points, otherwise, since N is discrete, such points would be isolated in Y . Thus $Y - N$ is dense in Y , hence open. We conclude that N is closed and discrete in Y .

Next, we will give an example in which $\beta X - X$ is C^* -embedded in βX ,

and it is neither a P-space nor an MI-space, but satisfies the hypotheses of Proposition 3 or 6.

Recall that a space is said to be extremally disconnected if every open set has an open closure. It is said to be basically disconnected if every cozero-set has an open closure. One can also give an equivalent definition of an F-space as being a space in which disjoint cozero-sets are completely separated.

Clearly, the following implications hold :

extremally disconnected \implies basically disconnected \implies F-space .

Let $\Sigma = N \cup \{\sigma\}$ and let U be a free ultrafilter on N . In Σ define the following topology : a subset A of Σ containing σ is open iff $A = U \cup \{\sigma\}$, $U \in U$, also all subsets of Σ that do not contain σ are to be open.

It is easy to prove that Σ is a normal, extremally disconnected space (and so an F-space), but it is not a P-space, nor an MI-space (cf. 4M in [GJ]). Since Σ is an F-space, then every subset of Σ is C^* -embedded (cf. 14N in [GJ]). It is known that, if Y is a Tychonoff space, then there is a space X such that $\beta X - X$ is homeomorphic to Y and is C^* -embedded in βX (cf. [C] Cor.4.18). We apply this result to the case $Y = \Sigma$.

Now we show that every infinite subset of Σ contains an infinite closed and discrete subset of Σ , so Σ is a cf-space.

Let T' be an infinite subset of Σ and let $T = T' - \{\sigma\}$. If $T = \{x_n\}$, set $A = \{x_{2n}\}$ and $B = \{x_{2n+1}\}$. Now if $A \in U$, then $A \cup (N - T) \in U$. Otherwise $N - A \in U$. In the former case, we obtain that B is closed and discrete. In the latter A is closed and discrete.

2. As we have seen in Proposition 3 and 6, the condition that $\beta X - X$ is a cf-space ensures, together with other conditions, that $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$. Now we want to prove that this latter condition implies that $\beta X - X$ is a cf-space.

PROPOSITION 8. If $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$ then $\beta X - X$ is a cf-space.

Proof.

Let K be a compact subset of $\beta X - X$ and suppose that K is infinite.

Then there is a countably infinite discrete subset of K , which we denote by $B = \{x_n\}$. If $\{r_n\}$ is the sequence of real numbers with $r_{2n-1} = r_{2n} = 1/n$, $n \in \mathbb{N}$, then the map $g : Cl_{\beta X} B \longrightarrow \mathbb{R}$ defined by $g(x_n) = r_n$ and $g(x) = 0$, if $x \in (Cl_{\beta X} B) - B$, is continuous and thus it has a continuous extension h to βX .

Now consider the family A of subsets of $\beta X - X$ defined as follows :

$$A = \{h^{-1}(p) \cap K : p \in \mathbb{R}\}$$

and let

$$S = \{f \in C^*(X) : f|_A \text{ is constant } \forall A \in A\}.$$

The family S separates points from closed sets of X . In fact if $C \subset X$ is a closed set and $x \in X$, $x \notin C$, let F be a closed set in βX such that $F \cap X = C$. Then there exists $f \in C^*(\beta X)$ such that $f(x) = 0$ and $f(K \cup F) = 1$. The map $f|_X$ belongs to S and separates x from C .

The family, S , thus generates a compactification $\alpha X = \alpha_S X = \overline{e_S(X)}$; moreover, if f_α is the canonical quotient map, one has $f_\alpha = e_S^\beta = \prod_{f \in S} f^\beta$.

Now we show that $F^*(\alpha X) = \{A \in A : |A| > 1\}$ and so $F^*(\alpha X)$ is infinite.

If $x, y \in A$, for some $A \in A$, then obviously $f_\alpha(x) = f_\alpha(y)$.

Conversely, suppose that $x, y \in \beta X - X$ do not belong to the same $A \in A$. If at least one of the two points, say x , does not belong to K , then there is a continuous map $s : \beta X \longrightarrow \mathbb{R}$ such that $s(x) = 0$ and $s(K \cup \{y\}) = 1$. We have $s|_X \in S$, and $(s|_X)^\beta(x) = s(x) \neq s(y) = (s|_X)^\beta(y)$, therefore $f_\alpha(x) \neq f_\alpha(y)$. Suppose then $x, y \in K$ and $h(x) \neq h(y)$, that is x and y do not belong to the same $A \in A$. The map $\bar{h} = h|_X \in S$. Obviously one has $\bar{h}^\beta(x) = h(x) \neq h(y) = \bar{h}^\beta(y)$, and so again we have $f_\alpha(x) \neq f_\alpha(y)$.

We note that the condition that $\beta X - X$ be a cf-space is not enough to imply that $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$. In fact, it is possible to construct a space X such that $\beta X - X = \mathbb{N} \approx \mathbb{N}$ and $Cl_{\beta X} \mathbb{N} = \omega \mathbb{N}$, where $\omega \mathbb{N}$ is the one-point compactification of \mathbb{N} . The conclusion follows from the following fact : if there is a sequence in $\beta X - X$ converging to a point of X , then $K(X)$ is not a lattice (cf. example 4.7 in [T]).

The following corollaries are easy consequences of Proposition 3 and 8.

COROLLARY 9. Let $\beta X - X$ be locally compact and C^* -embedded in βX .
Then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$ if and only if $\beta X - X$ is
discrete.

COROLLARY 10. Let X be locally compact.
Then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$ if and only if $\beta X - X$ is finite.

COROLLARY 11. Let $C_l(\beta X - X)$ be an F-space.
Then $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$ if and only if $\beta X - X$ is a
cf-space.

Remark: The hypotheses of Corollary 11 are satisfied, for instance, if X is an F-space or $\beta X - X$ is an F-space, C^* -embedded in βX (cf. in [GJ] 14.25 (9) and (10) and 14.26 that every C^* -embedded subspace of an F-space is itself an F-space).

We give now another necessary condition for $F^*(\alpha X)$ to be finite for all $\alpha X \in K(X)$.

PROPOSITION 12. If $F^*(\alpha X)$ is finite for all $\alpha X \in K(X)$, then X is pseudo-
compact.

Proof. If X were not pseudocompact, then it would contain a C-embedded copy N of \mathbb{N} , in particular a closed C^* -embedded copy of \mathbb{N} . Then we would have $\beta N - N \subset \beta X - X$ and so $\beta X - X$ would not be a cf-space.

Note that a pseudocompact space can contain a closed C^* -embedded copy of \mathbb{N} and so the converse of the above proposition is false. For example, the space $\Lambda = \beta \mathbb{R} - (\beta \mathbb{N} - \mathbb{N})$ is pseudocompact and \mathbb{N} is closed C^* -embedded in Λ (cf. 6P in [GJ]).

COROLLARY 13. If X is realcompact and not compact, then there exists
 $\alpha X \in K(X)$ with $F^*(\alpha X)$ infinite.

We conclude with an open question: is there a space X such that $\beta X - X$ is a cf-space, C^* -embedded in βX and X has a compactification αX with $F^*(\alpha X)$ infinite?

R E F E R E N C E S

- [C] R.CHANDLER, Hausdorff compactifications, Dekker, New York (1976).
- [FV] J.VISLISENI, J.FLEKSMOIER, The power and the structure of the lattice of all compact extensions of a completely regular space, Soviet Math. 6 (1965), 1423-1425.
- [GJ] L.GILMAN, M.JERISON, Rings of continuous functions, Van Nostrand, Princeton (1960).
- [K] M.R.KIRCH, A class of spaces in which compact sets are finite, Amer. Math. Monthly 76 (1969), 42.
- [M] K.MAGILL, The lattice of compactifications of a locally compact space, Proc. Lond. Math. Soc. 18 (1968), 231-244.
- [S] P.L.SHARMA, The Lindelöff property in MI-spaces, Ill. Journ. of Math. 25 (1981), 644-648.
- [T] F.C.TZUNG, Sufficient conditions for the set of Hausdorff compactifications to be a lattice, Pacif. J. Math. 77 (1978), 565-573.
- [U] Y.UNLU, Lattices of compactifications of Tychonoff spaces, Gen. Top. and its appl. 9 (1978), 41-57.

Alessandro CATERINO
Dipartimento di Matematica
Università di PERUGIA
Via Vanvitelli, 1
06100 PERUGIA (ITALY)

(Oblatum 5.11. 1984)