

Jouni Luukkainen

Canonical Lipschitz structures on compact Hilbert cube manifolds

Commentationes Mathematicae Universitatis Carolinae, Vol. 26 (1985), No. 4, 661--664

Persistent URL: <http://dml.cz/dmlcz/106405>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1985

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**CANONICAL LIPSCHITZ STRUCTURES ON COMPACT
HILBERT CUBE MANIFOLDS**
Jouni LUUKKAINEN

Abstract: Let d be one of the Lipschitz homogeneous metrics on the Hilbert cube Q discovered by Väisälä and Hohti. We show that every compact Q -manifold is homeomorphic to a Lipschitz (Q,d) -manifold of the form $M \times (Q,d)$ where M is a compact PL n -submanifold of \mathbb{R}^n for some n and that such an $M \times (Q,d)$ is unique up to a Lipschitz homeomorphism.

Key words: Hilbert cube, Lipschitz homeomorphism, Lipschitz manifold, Q -manifold.

Classification: 57N20

1. Existence. A homeomorphism $f: (X,d) \rightarrow (Y,d')$ of metric spaces is a *Lipschitz homeomorphism* if there is $L \geq 1$ such that

$$d(x,y)/L \leq d'(f(x),f(y)) \leq Ld(x,y) \quad (x,y \in X).$$

Let s be a sequence $s_1 \geq s_2 \geq \dots$ of positive real numbers converging to zero such that $(\mathbb{N} = \{1,2,\dots\})$

$$R(s) = \sup \{s_k/s_{k+1} : k \in \mathbb{N}\} < \infty.$$

Let Q_s denote the Hilbert cube $Q = [-1,1]^{\mathbb{N}}$ equipped with the compatible metric d ,

$$d(x,y) = \sup \{s_k |x_k - y_k| : k \in \mathbb{N}\}.$$

Definition 1. A *Lipschitz Q_s -manifold* is a separable metric space whose every point has a neighborhood Lipschitz homeomorphic to Q_s .

There is an essentially equivalent alternative definition based on atlases; cf. [7, 3.3-3.7]. Definition 1 is natural because A. Hohti [4, 5.3] has proved that every connected Lipschitz Q_s -manifold (and thus, in particular, Q_s itself)

is homogeneous with respect to Lipschitz homeomorphisms and because, on the other hand, J. Väisälä [12, 3.5] has proved that this is never true of Q_s if the condition $R(s) < \infty$ is not satisfied. The model cube Q_s is natural also because it is an absolute extensor for Lipschitz maps [8, Theorem 1], which implies, as in [7, 5.12], that every Lipschitz Q_s -manifold is an absolute neighborhood extensor for locally Lipschitz maps.

Example 2. For the cartesian product of finitely many metric spaces use any of the standard Lipschitz equivalent metrics. Define a *Lipschitz n-manifold* ($n \in \mathbb{N} \cup \{0\}$) by means of the model cube $I^n = [-1, 1]^n$ (see [7]). Since $R(s) < \infty$, the natural homeomorphism

$$I^n \times Q_s \rightarrow Q_s, \quad (x, y) \sim (x_1, \dots, x_n, y_1, y_2, \dots),$$

is a Lipschitz homeomorphism. Hence, if M is a Lipschitz n -manifold, $M \times Q_s$ is a Lipschitz Q_s -manifold.

Note that every PL homeomorphism of compact polyhedra in \mathbb{R}^n is a Lipschitz homeomorphism [7, 2.18] and that, thus, every PL manifold in \mathbb{R}^n is a Lipschitz manifold. (For PL topology we refer to [10].) It now immediately follows from well-known results on (topological) Q -manifolds [2] that there exists a Lipschitz Q_s -manifold structure on every compact Q -manifold:

Proposition 3. *If X is a compact Q -manifold, there is a compact PL n -manifold M in \mathbb{R}^n for some n such that X is homeomorphic to the Lipschitz Q_s -manifold $M \times Q_s$.*

Proof. By [2, 36.2] there is a compact polyhedron P in some \mathbb{R}^n such that X is homeomorphic to $P \times Q$. Choose a regular neighborhood M of P in \mathbb{R}^n . Then M is a compact PL n -manifold. Since P and M are simple homotopy equivalent, $P \times Q$ and $M \times Q$ are homeomorphic by [2, 29.4]. Hence, X is homeomorphic to $M \times Q_s$, which is a Lipschitz Q_s -manifold by the above. \square

Proposition 3 is an observation of Hohti and it is published with his permission. Hohti has since constructed a Lipschitz Q_s -manifold structure on every Q -manifold [5].

2. Uniqueness. We next show that a Lipschitz Q_s -manifold structure on a compact Q -manifold X induced by a homeomorphism $X \approx M \times Q_s$ as in Proposition 3 is unique up to a Lipschitz homeomorphism and may thus be called *canonical*.

Theorem 4. Let $M_i \subset \mathbb{R}^{n_i}$ be a compact PL n_i -manifold, $i=1,2$, such that $M_1 \times Q_S$ and $M_2 \times Q_S$ are homeomorphic. Then $M_1 \times Q_S$ and $M_2 \times Q_S$ are Lipschitz homeomorphic.

Proof. By [2, 38.1] (and the proof of [2, 29.5]), M_1 and M_2 are simple homotopy equivalent. Hence, by [13, Theorem 25], if we choose a sufficiently large $n \geq \max(n_1, n_2)$ and identify \mathbb{R}^{n_i} with $\mathbb{R}^{n_i} \times 0 \subset \mathbb{R}^n$, every regular neighborhood of M_i in \mathbb{R}^n is PL homeomorphic to every regular neighborhood of M_2 in \mathbb{R}^n . Choose a regular neighborhood N_i of M_i in \mathbb{R}^{n_i} . Then $N_i' = N_i \times I^{n-n_i}$ is a regular neighborhood of M_i in \mathbb{R}^n , because it collapses onto N_i and, thus, onto M_i . Hence, N_1' and N_2' are PL homeomorphic. Since M_i is PL homeomorphic to N_i , it follows that $M_1 \times I^{n-n_1}$ and $M_2 \times I^{n-n_2}$ are PL homeomorphic. Since Q_S is Lipschitz homeomorphic to $I^{n-n_i} \times Q_S$, this implies that $M_1 \times Q_S$ and $M_2 \times Q_S$ are Lipschitz homeomorphic. \square

It is not known whether every two compact Lipschitz Q_S -manifolds are Lipschitz homeomorphic if they are homeomorphic. This problem is equivalent to the problem whether every compact Lipschitz Q_S -manifold is Lipschitz homeomorphic to a Lipschitz Q_S -manifold with a canonical structure. Our final result shows that this is the case for some manifolds of Example 2.

Theorem 5. Suppose that M is a compact Lipschitz manifold and that either $\dim M = n$ or $\dim M = n-1$, $n \neq 4$, and $\partial M = \emptyset$. Suppose also that M can be topologically embedded into \mathbb{R}^n . Let $p = 6$ if $n = 4$ or 5 and let $p = n$ otherwise. Then there is a compact PL p -manifold N in \mathbb{R}^p such that $M \times Q_S$ is Lipschitz homeomorphic to $N \times Q_S$.

Proof. Suppose first that M is an n -manifold. We reduce the case $n = 4$ or 5 to the case $n = 6$ replacing M by $M \times I^{6-n}$. It suffices to find a PL n -manifold N in \mathbb{R}^n homeomorphic to M , because then M and N are Lipschitz homeomorphic by the generalization [11, 4.8] of a theorem of D. Sullivan. Choose a manifold $S \subset \mathbb{R}^n$ homeomorphic to M and an open collar $c: \partial S \times [0,1) \rightarrow S$ of ∂S in S . Then $T = S \setminus c[\partial S \times [0,1/2]]$ is homeomorphic to S and $c[\partial S \times (0,1)]$ is an open bicollar neighborhood of $\partial T = c[\partial S \times \{1/2\}]$ in \mathbb{R}^n . Hence, by [6, I, 5.1 and 4.1] if $n \geq 6$ or by classical results [9] if $n \leq 3$, there is a homeomorphism $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $N = fT$ is a PL submanifold of \mathbb{R}^n .

Suppose now that M is an $(n-1)$ -manifold. By [1, p. 61] if $n \geq 5$ or by classical results if $n \leq 3$, there is a locally flat embedding $f: M \rightarrow \mathbb{R}^n$. If S is a component of fM , by [3, 27.10] $\mathbb{R}^n \setminus S$ consists of two components, whose closures are n -manifolds with boundary S . Hence, there is an embedding

$g: M \times I^1 \rightarrow \mathbb{R}^n$. Thus, replacing M by $M \times I^1$ reduces the situation to the first case of the theorem. \square

References

- [1] ANCEL, F.D., and J.W. CANNON: The locally flat approximation of cell-like embedding relations, *Ann. of Math.* (2) 109 (1979), 61-86.
- [2] CHAPMAN, T.A.: Lectures on Hilbert cube manifolds, CBMS Regional Conference Series in Mathematics 28, American Mathematical Society, Providence, Rhode Island, 1976.
- [3] GREENBERG, M.J.: Lectures on algebraic topology, W.A. Benjamin, Inc., New York - Amsterdam, 1967.
- [4] HOHTI, A.: On Lipschitz homogeneity of the Hilbert cube, *Trans. Amer. Math. Soc.* (to appear).
- [5] HOHTI, A.: Lipschitz manifolds from Q -manifolds, manuscript.
- [6] KIRBY, R.C., and L.C. SIEBENMANN: Foundational essays on topological manifolds, smoothings, and triangulations, *Annals of Mathematics Studies* 88, Princeton University Press and University of Tokyo Press, Princeton, New Jersey, 1977.
- [7] LUUKKAINEN, J., and J. VÄISÄLÄ: Elements of Lipschitz topology, *Ann. Acad. Sci. Fenn. Ser. A I Math.* 3 (1977), 85-122.
- [8] McSHANE, E.J.: Extension of range of functions, *Bull. Amer. Math. Soc.* 40 (1934), 837-842.
- [9] MOISE, E.E.: *Geometric topology in dimensions 2 and 3*, Springer-Verlag, New York - Heidelberg - Berlin, 1977.
- [10] ROURKE, C.P., and B.J. SANDERSON: *Introduction to piecewise-linear topology*, Springer-Verlag, Berlin - Heidelberg - New York, 1972.
- [11] TUKIA, P., and J. VÄISÄLÄ: Lipschitz and quasiconformal approximation and extension, *Ann. Acad. Sci. Fenn. Ser. A I Math.* 6 (1981), 303-342.
- [12] VÄISÄLÄ, J.: Lipschitz homeomorphisms of the Hilbert cube, *Topology Appl.* 11 (1980), 103-110.
- [13] WHITEHEAD, J.H.C.: Simplicial spaces, nuclei and m -groups, *Proc. London Math. Soc.* (2) 45 (1939), 243-327.

University of Helsinki, Department of Mathematics, Hallituskatu 15,
SF-00100 Helsinki 10, Finland

(Oblatum 29.3. 1985)