

Věra Trnková

Full embeddings into the categories of Boolean algebras

Commentationes Mathematicae Universitatis Carolinae, Vol. 27 (1986), No. 3, 535--541

Persistent URL: <http://dml.cz/dmlcz/106474>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1986

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

FULL EMBEDDINGS INTO THE CATEGORIES
OF BOOLEAN ALGEBRAS
Věra TRNKOVÁ

Abstract: We prove that every small thin category can be fully embedded in the category of Boolean algebras and all one-one homomorphisms and also in the category of Boolean algebras and all surjective homomorphisms.

Key words: Boolean algebra, full embeddings of categories.

Classification: 18B15, 06E99

A category is called s-universal if every small category can be fully embedded in it. If \mathcal{K} is s-universal, then every monoid can be represented as the monoid of all endomorphisms of an object of \mathcal{K} (this is the result of a full embedding of a one-object category with the morphism part formed by the given monoid). Many current categories are known to be s-universal. This field of problems is extensively investigated in the monograph [PT].

If \mathcal{K} is an s-universal category, then also every group can be represented as the group of all automorphisms of an object of \mathcal{K} . This fact implies that neither the category

$\mathfrak{B}(1-1)$ of all Boolean algebras and all one-one homomorphisms

nor the category

$\mathfrak{B}(\text{onto})$ of all Boolean algebras and all surjective homomorphisms

is s-universal. In fact, there are groups not representable as

the groups of all automorphisms of a Boolean algebra, e.g. \mathcal{L}_3 , see [MM]. On the other hand, there are large Boolean algebras with the trivial group of automorphisms, [K]. By [M], there are arbitrarily large Boolean algebras such that the identities are their unique one-one endomorphisms. By [LR], for every uncountable cardinal λ , there is a Boolean algebra of the cardinality λ such that the identity is its unique surjective endomorphism. In the final remark of [LR], the existence of a full embedding of every small discrete category (i.e. having only the unit morphisms) into $\mathcal{B}(\text{onto})$ is stated. Here, we investigate full embeddings of small thin categories into the above categories of Boolean algebras (let us recall that a category k is thin if, for every couple of objects A, B of k , there is at most one morphism from A into B). Let us state explicitly that in the proof of the theorem stated below, the constructions of [LR] and [M] are essential and only a small reasoning is added to them. However, the embedding theorem seems to be of some interest in connection with the field of problems investigated in [PT].

Theorem. Every small thin category can be fully embedded into $\mathcal{B}(1-1)$ and into $\mathcal{B}(\text{onto})$.

Proof.

A) Full embeddings into $\mathcal{B}(1-1)$.

1. Let \aleph be a cardinal. Following [S] and [M], let us say that a Boolean algebra B is \aleph -complicated if, for every collection $\{(b_\alpha, a_\alpha) \mid \alpha < \aleph\}$ of pairs of non-zero elements of B such that

- a) $b_\alpha \wedge b_{\alpha'} = 0$ and $a_\alpha \wedge a_{\alpha'} = 0$ whenever $\alpha \neq \alpha'$ and
- b) $a_\alpha \not\leq b_\alpha$ for all $\alpha < \aleph$,

there exists $S \subseteq \aleph$ such that

- $\alpha)$ there exists no $w \in B$ with $a_\alpha \leq w$ for all $\alpha \in S$
 $a_\alpha \wedge w = 0$ for all $\alpha \in \aleph \setminus S,$
 $\beta)$ but there exists $u \in B$ with $b_\alpha \leq u$ for all $\alpha \in S$
 $b_\alpha \wedge u = 0$ for all $\alpha \in \aleph \setminus S.$

By [M], if \aleph is an infinite cardinal such that $2^\aleph = \aleph^{\aleph_0}$, then there exists an atomless \aleph -complicated Boolean algebra C (of the cardinality 2^\aleph) such that for any non-zero $a \in C$ there is a system $\{b_\alpha \mid \alpha < \aleph\}$ of non-zero pairwise disjoint elements of C such that $b_\alpha \leq a$ for all $\alpha < \aleph$.

2. Let C be as above. Let us verify the following assertion: if $c, d \in C$ and there is a one-one homomorphism h of $C \upharpoonright c$ onto $C \upharpoonright d$, then $c = d$ and h is the identity. In fact, suppose that h is not the identity. Then there is a non-zero $a \in C$, $a \leq c$, such that $a \wedge h(a) = 0$. Let $\{b_\alpha \mid \alpha < \aleph\}$ be a system of non-zero pairwise disjoint elements of C with $b_\alpha \leq a$ for all $\alpha < \aleph$. Then $\{b_\alpha, h(b_\alpha) \mid \alpha < \aleph\}$ fulfils the above a) b), hence there exists $S \subseteq \aleph$ and $u \in C$ such that the above $\alpha), \beta)$ are fulfilled (with a_α replaced by $h(b_\alpha)$). However, $w = h(u \wedge a)$ fulfils

$$h(b_\alpha) \leq w \text{ for all } \alpha \in S \text{ and}$$

$$h(b_\alpha) \wedge w = 0 \text{ for all } \alpha \in \aleph \setminus S,$$

which is a contradiction (this verification is analogous to the proof of Theorem 10 in [M]).

3. The conclusion of 1 and 2: For every cardinal \aleph there exists a Boolean space (= compact Hausdorff 0-dim) X such that

(i) every nonvoid clopen (= closed-and-open) subset D of X contains a pairwise disjoint collection $\{D_\alpha \mid \alpha < \aleph\}$ of nonvoid clopen subsets of X ;

(ii) if D, E are clopen subsets of X and f is a continuous map of D onto E , then $D = E$ and f is the identity.

4. Let a small thin category k be given. We may suppose

that it is skeletal (i.e. no two distinct objects of k are isomorphic). First, we fully embed k into the thin category of all infinite subsets of a set T , i.e. $\Psi: k \rightarrow \exp T$ is such that each $\Psi(A)$ is infinite and $k(A, B) \neq \emptyset$ iff $\Psi(A) \subseteq \Psi(B)$. Choose $\aleph \geq \text{card } T$, let X be a Boolean space which fulfils (i) and (ii) in 3. Choose $\{D_t \mid t \in T\}$ a pairwise disjoint collection of nonvoid clopen subsets of X . For every object A of k , denote by $\Phi(A)$ a one-point compactification of the subset $\bigcup_{t \in \Psi(A)} D_t$ of X , the added point is denoted by ξ_A . If $\Psi(B) \supseteq \Psi(A)$, let us define $\varphi_B^A: \Phi(B) \rightarrow \Phi(A)$ by

$$\varphi_B^A(x) = x \text{ for all } x \in \bigcup_{t \in \Psi(A)} D_t$$

$$\varphi_B^A(x) = \xi_A \text{ else.}$$

5. Let A, B be objects of k , let $f: \Phi(B) \rightarrow \Phi(A)$ be a surjective continuous mapping. We want to prove that $\Psi(B) \supseteq \Psi(A)$ and $f = \varphi_B^A$. For every clopen subset D of X such that

$$D \subseteq G = \bigcup_{t \in \Psi(A)} D_t \setminus \{f(\xi_B)\},$$

put $E = f^{-1}(D)$. Then, by (ii), $E = D$ and $f(x) = x$ for all $x \in E$. This implies that $f^{-1}(G) = G$ and $f(x) = x$ for all $x \in G$. Consequently, for every $t \in \Psi(A)$, we have $f(\xi_B) \notin D_t$. Thus $\Psi(B) \supseteq \Psi(A)$ and $f(x) = x$ for all $x \in \bigcup_{t \in \Psi(A)} D_t = G$. The continuity of f implies $f(\xi_B) = \xi_A$ and f sends each D_t with $t \notin \Psi(A)$ on ξ_A because $f^{-1}(G) = G$. Thus $f = \varphi_B^A$. Consequently, the map

$$A \mapsto \text{all clopen subsets of } \Phi(A)$$

defines a full embedding of k into $\mathcal{B}(1-1)$.

B) Full embeddings into $\mathcal{B}(\text{onto})$.

1. The construction of [LR] will be used; let us recall some facts and notation. If λ is an uncountable regular cardinal, $I(\lambda)$ is the ideal of all subsets of λ disjoint from some closed

unbounded subset of λ and $D(\lambda) = P(\lambda)/I(\lambda)$ is the Boolean algebra of all stationary subsets of λ . If X is a topological space, $x \in X$, then $Cf(x, X)$ denotes the set of all regular infinite cardinals μ such that there is a sequence $\{x_i | i < \mu\}$ in X with

$$x = \lim_{i < \mu} x_i, \text{ and, for every } \alpha < \mu,$$

$$\lim_{i < \alpha} x_i \text{ exists and is distinct from } x.$$

We say that $x \in X$ is λ -special if $\lambda \in Cf(x, X)$ and, for every $\{x_i | i < \lambda\}$ and $\{y_i | i < \lambda\}$ as in the definition of $Cf(x, X)$, the set $\{\alpha | \lim_{i < \alpha} x_i = \lim_{i < \alpha} y_i\}$ is closed and unbounded in λ . If x is λ -special, then S_x^λ is the element of $D(\lambda)$ defined as follows: $S_x^\lambda = S'/I(\lambda)$, where

$$S' = \{\alpha | \lambda \in Cf(\lim_{i < \alpha} x_i, X)\},$$

with $\{x_i | i < \lambda\}$ as in the definition of $Cf(x, X)$ (since x is λ -special, S_x^λ is independent of the choice of $\{x_i | i < \lambda\}$).

2. Let λ be an uncountable regular cardinal. In [LR], a complete linear ordering I is constructed such that the set of all its elements having a successor is dense in it and

- (a) for every $x \in I$, either $\lambda \notin Cf(x, I)$ or x is λ -special;
- (b) the set P of all λ -special $x \in I$ with $S_x^I \neq 0$ is dense in I ;
- (c) if $x, y \in P$, $x \neq y$, then $S_x^I \cap S_y^I = 0$.

By [LR], I (with the order topology) is a Boolean space such that the identity is the unique one-one continuous map of I into itself. Since $D(\lambda)$ contains λ pairwise disjoint non-zero elements, we can obtain, by the same construction, a collection $\{I_\gamma | \gamma < \lambda\}$ of linear orderings such that each of them has all the above properties and, moreover,

- (d) if $x \in I_\alpha$, $y \in I_\beta$ are λ -special and $\alpha \neq \beta$, then $S_x^\alpha \cap S_y^\beta = 0$.

Then the following statement is fulfilled:

(*) $\left\{ \begin{array}{l} \text{if } \alpha, \beta < \lambda, K \text{ is a clopen nonvoid subset of } I_\alpha \text{ and the} \\ \text{re is a one-one continuous map } f: K \rightarrow I_\beta, \text{ then } \alpha = \beta \text{ and} \\ f \text{ is the inclusion (i.e. } f(x) = x \text{ for all } x \in K). \end{array} \right.$

In fact, let us suppose that f is not the inclusion. Since $P_\alpha \cap K$ is dense in K (where P_α is the subset of I_α as in (b)), there exists $x \in P_\alpha \cap K$ such that $f(x) \neq x$. By (b), x is λ -special in I_α with $S_x^K = S_x^{I_\alpha} \neq 0$. Since $\lambda \in \text{Cf}(x, I_\alpha)$, λ is also in $\text{Cf}(f(x), I_\beta)$. By (a), $f(x)$ is λ -special in I_β and, clearly, $S_x^K \subseteq S_{f(x)}^{I_\beta}$. However, by (b), (c) and (d), $S_{f(x)}^{I_\beta}$ is either 0 or disjoint from $S_x^{I_\alpha}$, which is a contradiction.

3. Let a small thin skeletal category be given. We embed fully its dual category k^* into the thin category of all infinite subsets of a set I ; denote by $\Psi: k^* \rightarrow \exp I$ the embedding. Find a pairwise disjoint collection $\{I_t \mid t \in I\}$ of Boolean spaces with the above properties and $\lambda = \text{card } I$ (we may suppose that λ is an uncountable regular cardinal). For every object A of k^* , put again

$$\Phi(A) = \{\xi_A\} \cup \bigcup_{t \in \Psi(A)} I_t,$$

where I_t are clopen in $\Phi(A)$ and ξ_A makes a one-point compactification of the union. If $\Psi(A) \subseteq \Psi(B)$, then $\varphi_A^B: \Phi(A) \rightarrow \Phi(B)$, defined by

$$\varphi_A^B(\xi_A) = \xi_B, \quad \varphi_A^B(x) = x \text{ for all } x \in \bigcup_{t \in \Psi(A)} I_t,$$

is a one-one continuous map. Now, let A, B be arbitrary objects of k^* and $f: \Phi(A) \rightarrow \Phi(B)$ be a one-one continuous map. We want to show that then $\Psi(A) \subseteq \Psi(B)$ and $f = \varphi_A^B$. For every $a \in \Psi(A)$, put

$$B_a = \{b \in \Psi(B) \mid f(I_a) \cap I_b \neq \emptyset\}.$$

Since $f(I_a) \setminus \{\xi_B\} \neq \emptyset$, the set B_a is not empty. For every $b \in B_a$,

put $\mathcal{O}_{a,b} = I_a \cap f^{-1}(I_b)$. Then $\mathcal{O}_{a,b}$ is a nonvoid clopen subset of I_a and f defines a one-one continuous map of $\mathcal{O}_{a,b}$ into I_b . By $(*)$, $b = a$ and $f(x) = x$ for all $x \in \mathcal{O}_{a,b}$. Hence $B_a = \{a\}$ and $\mathcal{O}_{a,a}$ is clopen in I_a . Since I_a has no isolated points (see (b)), necessarily $\mathcal{O}_{a,a} = I_a$. Consequently $\Psi(A) \subseteq \Psi(B)$ and $f(x) = x$ for all $x \in \bigcup_{a \in \Psi(A)} I_a$. By the continuity of f , $f(\xi_A) = \xi_B$, consequently $f = \varphi_A^B$. Thus, Φ is a full embedding of k^* into the category of Boolean spaces and one-one continuous maps, hence it determines a full embedding of k into $\mathcal{B}(\text{onto})$.

References

- [K] M. KATĚTOV: Remarks on Boolean algebras, Coll. Math. 2 (1951), 229-235.
- [LR] J. LOATS and M. RUBIN: Boolean algebras without non-trivial onto endomorphism exists in every uncountable cardinality, Proc. Amer. Math. Soc. 72(1978), 346-351.
- [MM] R. MCKENZIE and J.D. MONK: On automorphism groups of Boolean algebras, 10. Infinite and finite sets, Keszthély (Hungary), 1973, 951-988.
- [M] J.D. MONK: A very rigid Boolean algebra, Israel J. of Math. 35(1980), 135-150.
- [PT] A. PULTR and V. TRNKOVÁ: Combinatorial, algebraic and topological representations of groups, semigroups and categories, North-Holland, 1980.
- [S] S. SHELAH: Boolean algebras with few endomorphisms, Proc. Amer. Math. Soc. 74(1979), 135-142.

Matematický ústav, Karlova univerzita, Sokolovská 83, 186 00 Praha 8, Czechoslovakia

(Oblatum 14.4. 1986)