

Abu Saleh Abdun Noor; William H. Cornish
Around a neutral element in a nearlattice

Commentationes Mathematicae Universitatis Carolinae, Vol. 28 (1987), No. 2, 199--210

Persistent URL: <http://dml.cz/dmlcz/106531>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1987

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

AROUND A NEUTRAL ELEMENT IN A NEARLATTICE

A.S.A. NOOR and W.H. CORNISH

Abstract: Nearlattices, or lower semilattices in which any two elements have a supremum whenever they are bounded above, provide an interesting generalization of lattices. In this context, we define different types of elements in a nearlattice S and then for a fixed element n , using the ternary operation J_n , study the behaviour of $S_n = (S; \wedge)$ where $x \wedge y = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$; $x, y \in S$.

Key words: Standard element, neutral element, nearlattice.

Classification: 06A12, 06A99, 06B10

1. Introduction. A nearlattice is a lower semilattice which has the property that any two elements possessing a common upper bound, have a supremum. Cornish and Hickman [1] called this the upper bound property. For detailed literature, we refer the reader to consult [1], [2] and [7].

A nearlattice-congruence Φ on a nearlattice S is a congruence of the underlying lower semilattice such that, whenever $a_1 \equiv b_1$, $a_2 \equiv b_2$ (Φ) and $a_1 \vee a_2$, $b_1 \vee b_2$ exist, $a_1 \vee a_2 \equiv b_1 \vee b_2$ (Φ). In the second section of [4], a fundamental contribution was made by Hickman. Defining a ternary operation j on a nearlattice S by $j(x, y, z) = (x \wedge y) \vee (y \wedge z)$, he showed that the resulting algebras of the type $(S; j)$ form a variety.

Standard and neutral elements, as well as standard ideals in a nearlattice were extensively studied in [2]. An element s in a nearlattice S is called standard if for all $x, y, t \in S$, $t \wedge [(x \wedge y) \vee (x \wedge s)] = (t \wedge x \wedge y) \vee (t \wedge x \wedge s)$. An element n in a nearlattice S is called neutral if it is standard and for any $t, x, y \in S$, $n \wedge [(t \wedge x) \vee (t \wedge y)] = (n \wedge t \wedge x) \vee (n \wedge t \wedge y)$. Clearly, every element of a distributive nearlattice is neutral. An ele-

ment n of a nearlattice S is called superstandard if it is standard and for any $x, y \in S$, $n \wedge [(x \wedge y) \vee (x \wedge n) \vee (y \wedge n)] = (x \wedge n) \vee \vee (y \wedge n)$, whenever $(x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$ exists. Of course, every neutral element is superstandard. But in the pentagonal lattice $\{0, a, b, n, 1\}$ where $0 < a < n < 1$; $0 < b < 1$; $a \wedge b = n \wedge b = 0$ and $a \vee b = n \vee b = 1$, n is superstandard but not neutral. [7] provides an example of a standard element in a lattice which is not superstandard.

An element n in a nearlattice S is called medial if $m(x, n, y) = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$ exists for all $x, y \in S$, while n is called sesquimedial if $J_n(x, y, z) = [(x \wedge n) \vee (y \wedge n)] \wedge [(y \wedge n) \vee (z \wedge n)] \vee \vee j(x, y, z)$ exists for all $x, y, z \in S$ where $j(x, y, z) = (x \wedge y) \vee (y \wedge z)$. Since $J_n(x, y, x) = m(x, n, y)$ for all $x, y \in S$, any sesquimedial element is medial. A nearlattice S is called medial if $m(x, y, z) = (x \wedge y) \vee \vee (y \wedge z) \vee (z \wedge x)$ exists for all $x, y, z \in S$. Of course, every element of a medial nearlattice is sesquimedial (see Lemma 3.1).

Let n be a fixed element of a nearlattice S . By an n -ideal of S , we mean a convex subnearlattice of S containing n . The n -ideal generated by a_1, \dots, a_m is denoted by $\langle a_1, \dots, a_m \rangle_n$. Clearly $\langle a_1, \dots, a_m \rangle_n = \langle a_1 \rangle_n \vee \dots \vee \langle a_m \rangle_n$. When S is a lattice, $\langle a_1, \dots, a_m \rangle_n = \langle a_1 \wedge \dots \wedge a_m \wedge n, a_1 \vee \dots \vee a_m \vee n \rangle_n$. Thus, for a lattice S , the set of finitely generated n -ideals of S is a lattice and its members are simply the intervals $[a, b]$ such that $a \leq n \leq b$, and for such intervals, $[a, b] \vee [a_1, b_1] = [a \wedge a_1, b \vee b_1]$ and $[a, b] \cap [a_1, b_1] = [a \vee a_1, b \wedge b_1]$. The n -ideal generated by a single element is called a Principal n -ideal and the set of Principal n -ideals of S is denoted by $P_n(S)$. When S is a lattice, it is not hard to see that $P_n(S)$ is a lattice if and only if n is complemented in each interval containing it.

For a fixed element n , the binary operation $x \circ y = m(x, n, y) = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$ has been studied by several authors including Jakubík and Kolibiar [5] for distributive lattices, Sholander [8] for distributive medial near lattices and Kolibiar [6] for an arbitrary lattice with n as a neutral element in it. Sholander [8] showed that for a distributive medial nearlattice S , $(S; \circ)$ is a semilattice. On the other hand Kolibiar [6] showed that if n is a neutral element in an arbitrary lattice S , $(S; \circ)$ is a semilattice. Recently, Noor [7] extended their work and showed that for a neutral and sesquimedial element n of a near-

lattice S , $S_n = (S; \cap)$ is not only a semilattice, it is a nearlattice. Moreover, the n -ideals of S are precisely the ideals of S_n . According to [7], we refer to S_n as an isotope of S .

In Section 2, we introduce the notion of a nearly neutral element in a nearlattice and then generalize and extend some of the results in [7]. We show that for a medial superstandard element n of a nearlattice S , S_n is a nearlattice wherein $J_n(x, y, z) = j_n(x, y, z)$ if and only if n is nearly neutral and sesquimedial in S . We also show that for a nearly neutral and sesquimedial element of a nearlattice S , n is neutral if and only if the nearlattice congruences of S are precisely the nearlattice congruences of S_n .

In Section 3, introducing the ternary operation $M_n(x, y, z)$, we show that for a sesquimedial neutral element n of a nearlattice S , S is medial if and only if S_n is so.

2. Nearly neutral element of a near lattice. An element n of a nearlattice is called nearly neutral if it is standard and has the property $n \wedge ((t \wedge x \wedge n) \vee (t \wedge y)) = (t \wedge x \wedge n) \vee (t \wedge y \wedge n)$ for all $x, y, t \in S$. Of course, a neutral element is always nearly neutral. Observe that in Figure 1, n is nearly neutral but $n \wedge (a \vee b) > (n \wedge a) \vee (n \wedge b)$ shows that it is not neutral there.

The following result shows that every nearly neutral element is superstandard, but in the pentagonal lattice $\{0, a, b, n, 1\}$ where $0 < a < n < 1$; $0 < b < 1$; $a \wedge b = n \wedge b = 0$; $a \vee b = n \vee b = 1$, n is superstandard but not nearly neutral.

Proposition 2.1. For an element n of a nearlattice S , the following conditions are equivalent.

- (i) For all $x, y, t \in S$,

$$n \wedge ((t \wedge x \wedge n) \vee (t \wedge y)) = (t \wedge x \wedge n) \vee (t \wedge y \wedge n).$$
- (ii) For all $x, y \in S$,

$$n \wedge ((x \wedge n) \vee y) = (x \wedge n) \vee (y \wedge n),$$
 whenever $(x \wedge n) \vee y$ exists.

Moreover, if n is sesquimedial, (i) and (ii) are also equivalent to each of the next two conditions.

- (iii) For all $x, y, z \in S$, $(x \cap y) \wedge n = (x \wedge n) \vee (y \wedge n)$ and

$$J_n(x, y, z) \wedge n = (x \cap y) \wedge (y \cap z) \wedge n,$$
 where $x \cap y = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$.

(iv) For all $x, y, z \in S$, $(x \cap y) \wedge n = (x \wedge n) \vee (y \wedge n)$ and $J_n(x, y, z) \wedge n \notin x \cap y$.

Proof. (i) \Rightarrow (ii). Suppose $(x \wedge n) \vee y$ exists. Then $n \wedge ((x \wedge n) \vee y) = n \wedge [(((x \wedge n) \vee y) \wedge x) \wedge n] \vee [(((x \wedge n) \vee y) \wedge y)] = (x \wedge n) \vee (y \wedge n)$.
(ii) \Rightarrow (i) is trivial.

Suppose now that n is sesquimedial and (i) and (ii) hold. Then $n \wedge (x \cap y) = n \wedge ((x \wedge n) \vee (y \wedge n) \vee (x \wedge y)) = n \wedge [(((x \wedge n) \vee (y \wedge n)) \wedge n) \vee (x \wedge y)] = (x \wedge n) \vee (y \wedge n) \vee (x \wedge y \wedge n) = (x \wedge n) \vee (y \wedge n)$. Also,
 $J_n(x, y, z) \wedge n = n \wedge [(((x \wedge n) \vee (y \wedge n)) \wedge ((y \wedge n) \vee (z \wedge n))) \vee (x \wedge y) \vee (y \wedge z)] =$
 $= n \wedge [((x \cap y) \wedge (y \cap z) \wedge n) \vee (x \wedge y) \vee (y \wedge z)] =$
 $= ((x \cap y) \wedge (y \cap z) \wedge n) \vee (n \wedge ((x \wedge y) \vee (y \wedge z))) = (x \cap y) \wedge (y \cap z) \wedge n$.

Thus (iii) holds.

Clearly (iii) implies (iv).

Finally suppose (iv) holds. Let $x, y \in S$ be such that $(x \wedge n) \vee y$ exists. Then

$J_n(x, y, (x \wedge n) \vee y) = [(((x \wedge n) \vee (y \wedge n)) \wedge (y \wedge n) \vee (n \wedge ((x \wedge n) \vee y)))] \vee (x \wedge y) \vee y =$
 $= (x \wedge n) \vee (y \wedge n) \vee y = (x \wedge n) \vee y$, and so by (iv) $n \wedge ((x \wedge n) \vee y) \notin x \cap y$.
Thus, $n \wedge ((x \wedge n) \vee y) \notin n \wedge (x \cap y) = (x \wedge n) \vee (y \wedge n)$; it follows that $n \wedge ((x \wedge n) \vee y) = (x \wedge n) \vee (y \wedge n)$ and (ii) holds. \square

The following result is found in [7, Th. 2.1].

Proposition 2.2. If n is a standard element of a nearlattice S , then $(S; \subseteq)$ is a partially ordered set and the map $x \rightarrow \langle x \rangle_n$ is an isomorphism of $(S; \subseteq)$ onto $P_n(S)$, where on S , $x \subseteq y$ if and only if $(x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$ exists and is equal to x . \square

Let n be a medial element of a nearlattice S . For any $x, y \in S$ define the binary operation $x \cap y = m(x, n, y) = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$. Recently Noor in [7] proved the following result.

Theorem 2.3. If n is a medial and standard element of a nearlattice S , then S_n is a semilattice if and only if n is superstandard in S .

Moreover, when n is neutral and sesquimedial then S_n is in fact a nearlattice and the n -ideals of S are precisely the ideals of S_n . \square

Our next theorem generalizes and extends the above Theorem. To obtain this, we need the following lemma. (i) is found in

[7; Lemma 2.4], and the proof of (ii) is similar to the proof of (ii) in [7; Lemma 2.4].

Lemma 2.4. In a nearlattice S ,

(i) a subset K of S is an ideal of S if and only if for all $x, y \in K$ and $a \in S$, $j(x, a, y) \in K$.

(ii) If n is a superstandard element of S such that S_n is a nearlattice wherein

$$J_n(x, y, z) = j_n^{S_n}(x, y, z) = (xny) \vee (y nz),$$

then a subset K of S is an n -ideal of S if and only if it contains n and $J_n(x, a, y) \in K$ for any $x, y \in K$ and $a \in S$. \square

Corollary 2.5. Suppose n is a superstandard element of a nearlattice S such that the isotope S_n of S is itself a nearlattice wherein $J_n(x, y, z) = j_n^{S_n}(x, y, z)$. Then the ideals of S_n are precisely the n -ideals of S . \square

Theorem 2.6. If S is a nearlattice and $n \in S$ is medial and superstandard, then the following conditions are equivalent.

(i) n is nearly neutral and sesquimedial in S .

(ii) The isotope $S_n = (S; \circ)$ is a nearlattice wherein

$$J_n^{S_n}(x, y, z) = J_n(x, y, z).$$

(iii) S_n has the upper bound property and n -ideals of S are precisely the ideals of S_n .

(iv) Any finitely generated n -ideal contained in a principal n -ideal is a principal n -ideal.

Proof. (i) \Rightarrow (ii). Suppose n is nearly neutral and sesquimedial in S . Then, clearly

$$J_n(x, y, z) = ((xny) \wedge (y nz) \wedge n) \vee j(x, y, z),$$

and so by [2; Th. 2.4], $J_n(x, y, z) \equiv j(x, y, z) (\Theta_n)$ and

$xny \equiv x \wedge y (\Theta_n)$. Hence $(xny) \wedge J_n(x, y, z) \equiv x \wedge y (\Theta_n)$ and similarly $(y nz) \wedge J_n(x, y, z) \equiv y \wedge z (\Theta_n)$. Therefore,

$$\begin{aligned} [(xny) \wedge J_n(x, y, z)] \vee [(y nz) \wedge J_n(x, y, z)] \vee [n \wedge J_n(x, y, z)] &\equiv \\ &\equiv (x \wedge y) \vee (y \wedge z) \vee (n \wedge j(x, y, z)) = j(x, y, z) (\Theta_n). \end{aligned}$$

Since the left hand side of this congruence exceeds the right hand side, by [2; Th. 2.4],

$$\begin{aligned} &\text{left hand expression} \\ &= j(x, y, z) \vee (n \wedge (\text{left hand expression})) \end{aligned}$$

$$= j(x,y,z) \vee (n \wedge J_n(x,y,z)) = J_n(x,y,z).$$

Thus, $J_n(x,y,z) \in \langle x\cap y, y\cap z \rangle_n$. On the other hand, $(x\cap y) \wedge J_n(x,y,z) \equiv x \wedge y (\Theta_n)$ implies $(x\cap y) \wedge J_n(x,y,z) = (x \wedge y) \vee (n \wedge (x\cap y) \wedge J_n(x,y,z))$, and so $((x\cap y) \wedge J_n(x,y,z)) \vee ((x\cap y) \wedge n) = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n) = x \cap y$. Hence, $x \cap y \in \langle J_n(x,y,z) \rangle_n$ and similarly $y \cap z \in \langle J_n(x,y,z) \rangle_n$. Thus, $\langle x\cap y, y\cap z \rangle_n = \langle J_n(x,y,z) \rangle_n$ and so by Proposition 2.2, $(x\cap y) \cup (y\cap z) = J_n(x,y,z)$.

(ii) \Rightarrow (iii) follows immediately from Corollary 2.5.

(iii) \Rightarrow (iv) is an easy consequence of the isomorphism of $(S_n; \subseteq)$ and $(P_n(S); \subseteq)$, and the upperbound property of S_n .

(iv) \Rightarrow (i). Let $a, b, c \in S$. Since $a \cap b, b \cap c \in b$, $\langle a \cap b, b \cap c \rangle_n \subseteq \subseteq \langle b \rangle_n$ by Proposition 2.2. Thus, by (iv), there exists $t \in S$ such that $\langle a \cap b, b \cap c \rangle_n = \langle t \rangle_n$, and so $(a \cap b) \wedge (b \cap c) \wedge n = t \wedge n$. Now, $a \cap b \subseteq t$ implies $a \cap b = ((a \cap b) \wedge t) \vee (a \cap b) \wedge n \vee (t \wedge n) = ((a \cap b) \wedge t) \vee ((a \cap b) \wedge n)$, and so $a \cap b \equiv (a \cap b) \wedge t (\Theta_n)$. Hence $a \wedge b \equiv a \cap b \equiv (a \cap b) \wedge t \equiv a \wedge b \wedge t (\Theta_n)$.

Similarly, $b \wedge c \equiv b \wedge c \wedge t (\Theta_n)$. This implies

$$j(a,b,c) \equiv (a \wedge b \wedge t) \vee (b \wedge c \wedge t) (\Theta_n)$$

and so $j(a,b,c) = (a \wedge b \wedge t) \vee (b \wedge c \wedge t) \vee (n \wedge j(a,b,c))$. Also, $j(a,b,c) \wedge t \equiv (a \wedge b \wedge t) \vee (b \wedge c \wedge t) (\Theta_n)$, and so

$$j(a,b,c) \wedge t = (a \wedge b \wedge t) \vee (b \wedge c \wedge t) \vee (n \wedge t \wedge j(a,b,c)).$$

Thus, $j(a,b,c) \cap t = (j(a,b,c) \wedge t) \vee (j(a,b,c) \wedge n) \vee (t \wedge n) = j(a,b,c) \vee (t \wedge n)$.

Again, $a \cap b \equiv a \wedge b (\Theta_n)$. So $(a \cap b) \wedge j(a,b,c) \equiv a \wedge b \wedge j(a,b,c) = a \wedge b (\Theta_n)$, and hence $(a \cap b) \wedge j(a,b,c) = (a \wedge b) \vee ((a \cap b) \wedge j(a,b,c) \wedge n)$. This implies $(a \cap b) \cap j(a,b,c) = a \cap b$; that is, $a \cap b \subseteq j(a,b,c)$. Similarly, $b \cap c \subseteq j(a,b,c)$. Hence, $t \subseteq j(a,b,c)$, and so $t = t \wedge j(a,b,c) = j(a,b,c) \vee (t \wedge n) = j(a,b,c) \vee ((a \cap b) \wedge (b \cap c) \wedge n) = J_n(a,b,c)$, as n is superstandard. Hence n is sesquimedial, and $J_n(a,b,c) \wedge n = t \wedge n = (a \cap b) \wedge (b \cap c) \wedge n$. Also $(x \cap y) \wedge n = (x \wedge n) \vee (y \wedge n)$, as n is superstandard. Therefore, by 2.1(iii), n is nearly neutral. \square

The following lemma is due to Hickman [4; Proposition 2.2].

Lemma 2.7. In a nearlattice S , an equivalence relation is a nearlattice congruence if and only if it is a congruence for the algebra $(S; j)$. \square

Now we consider the influence of J_n on congruences. The following theorem is an extension of [7; Lemma 2.6(ii)].

Theorem 2.8. Let n be a sesquimedial, nearly neutral element of a nearlattice S . Then the following conditions are equivalent.

- (i) n is neutral in S ;
- (ii) an equivalence relation on S is a congruence for the algebra $(S; J_n)$ if and only if it is a nearlattice-congruence of S .

Proof. (i) \Rightarrow (ii) is proved in [7; Lemma 2.6(ii)].

(ii) \Rightarrow (i). Define a relation Θ on the nearlattice S by $x \equiv y(\Theta)$ if and only if $x \wedge n = y \wedge n$. This is clearly an equivalence relation on S .

Now suppose $x \equiv y(\Theta)$. Then $x \wedge n = y \wedge n$, and so by 2.1, for any $s, t \in S$, $n \wedge J_n(x, s, t) = (x \wedge n) \wedge (s \wedge t) \wedge n = ((x \wedge n) \vee (s \wedge n)) \wedge ((s \wedge n) \vee (t \wedge n)) = ((y \wedge n) \vee (s \wedge n)) \wedge ((s \wedge n) \vee (t \wedge n)) = n \wedge J_n(y, s, t)$. Thus, $J_n(x, s, t) \equiv J_n(y, s, t)(\Theta)$. Similarly, $J_n(s, x, t) \equiv J_n(s, y, t)(\Theta)$ and $J_n(s, t, x) \equiv J_n(s, t, y)(\Theta)$, and so Θ is a congruence for the algebra $(S; J_n)$. Thus, by (ii), Θ is a nearlattice congruence on S . Now, clearly $x \equiv x \wedge n(\Theta)$ and $y \equiv y \wedge n(\Theta)$ for all $x, y \in S$. So for any $t \in S$, $(t \wedge x) \vee (t \wedge y) \equiv (t \wedge x \wedge n) \vee (t \wedge y \wedge n)(\Theta)$, and hence, $n \wedge [(t \wedge x) \vee (t \wedge y)] = n \wedge [(t \wedge x \wedge n) \vee (t \wedge y \wedge n)] = (t \wedge x \wedge n) \vee (t \wedge y \wedge n)$, which implies n is neutral in S . \square

Combining Theorem 2.6, Lemma 2.7 and the above theorem, we have the following extension of [7, Th. 2.7].

Theorem 2.9. Let n be a nearly neutral sesquimedial element of a nearlattice S . Then n is neutral if and only if the nearlattice congruences of S are precisely the nearlattice congruences of S_n . \square

The following proposition will be needed to prove one of our main results in Section 3. This was known by Kolibiar [6] in case of a bounded lattice with n as a central element.

Proposition 2.10. If n is a nearly neutral sesquimedial element of a nearlattice S with 0 , then 0 is neutral and medial in S_n . Moreover, the double isotope $(S_n)_0$ is precisely S .

If, in addition, n is neutral in S , then 0 is sesquimedial in S_n and $J_0^n(x, y, z) = j_{(S_n)_0}^n(x, y, z) = j(x, y, z) = J_0(x, y, z)$ for all $x, y, z \in S$.

Proof. By 2.6, for all $r, x, y \in S$, $0 \cap ((rx) \cup (ry)) =$
 $= 0 \cap J_n(x, r, y) = n \wedge J_n(x, r, y) = J_n(rx, 0, ry) = (0 \cap rx) \cup (0 \cap ry).$
 Also, $r \cap ((xy) \cup (x0)) = r \cap J_n(y, x, 0) = r \cap ((x \wedge n) \vee (x \wedge y)) = (r \wedge n) \vee$
 $\vee (x \wedge n) \vee (r \wedge x \wedge y)$ as n is nearly neutral and hence standard. On the
 other hand, $(rx \wedge y) \cup (rx \wedge 0) = J_n(y, rx, 0) = ((rx) \wedge n) \vee (y \wedge (rx)) =$
 $= (r \wedge n) \vee (x \wedge n) \vee [y \wedge ((rx) \wedge r \wedge x) \vee ((rx) \wedge n)] = (r \wedge n) \vee (x \wedge n) \vee (r \wedge x \wedge y).$
 That is $r \cap ((xy) \cup (x0)) = (rx \wedge y) \cup (rx \wedge 0)$; consequently 0 is ne-
 utral in S_n .

Now, clearly $x \wedge y, x \wedge 0, y \wedge 0 \in x \wedge y$, and so $(xy) \cup (x0) \cup (y0)$ ex-
 ists and it is $\in x \wedge y$. Thus, 0 is medial in S_n , and so $((S_n)_0; \bar{\wedge})$
 is a semilattice by Theorem 2.3, where

$$x \bar{\wedge} y = (xy) \cup (x0) \cup (y0).$$

Suppose $x \wedge y, x \wedge 0, y \wedge 0 \in s$ for some $s \in S_n$. Then $s \wedge n \in (x0) \wedge n =$
 $= x \wedge n$. Similarly $s \wedge n \in y \wedge n$, and so $s \wedge n \in x \wedge y \wedge n$. Also,

$$\begin{aligned} x \wedge y &= (x \wedge y) \wedge s = ((x \wedge y) \wedge s) \vee ((x \wedge y) \wedge n) \vee (s \wedge n) = \\ &= ((x \wedge y) \wedge s) \vee ((x \wedge y) \wedge n). \end{aligned}$$

Then

$$x \wedge y = (x \wedge y) \wedge (x \wedge y) = (x \wedge y \wedge s) \vee (x \wedge y \wedge n) = (x \wedge y) \wedge s.$$

This implies $x \wedge y \in s$, and hence

$$x \bar{\wedge} y = (xy) \cup (x0) \cup (y0) = x \wedge y;$$

in other words, $(S_n)_0 = S$.

Finally, suppose that n is neutral in S . Since 0 is neutral in S_n ,

$$\begin{aligned} ((x0) \cup (y0)) \cap ((y0) \cup (z0)) &= (x \bar{\wedge} y) \cap (y \bar{\wedge} z) \cap 0 = \\ &= (x \wedge y) \cap (y \wedge z) \cap 0 = [(x \wedge y) \cap (y \wedge z)] \wedge n = \\ &= (x \wedge y \wedge n) \vee (y \wedge z \wedge n) = n \wedge j(x, y, z) \end{aligned}$$

as n is neutral. Also it can be easily shown that $x \wedge y, y \wedge z \in$
 $\in j(x, y, z) = J_0(x, y, z)$. Therefore

$$[((x0) \cup (y0)) \cap ((y0) \cup (z0))] \cup (x \wedge y) \cup (y \wedge z)$$

exists in S_n ; whence 0 is sesquimedial in S_n . The rest follows by
 2.6. \square

It should be noted that the above proposition is not true
 when n is merely nearly neutral. For example, in Figure 2 which
 is the isotope of Figure 1, 0 is not sesquimedial.

3. Medial nearlattices. Recall that a nearlattice S is medial if for all $x, y, z \in S$, $m(x, y, z) = (x\lambda y)\vee(y\lambda z)\vee(z\lambda x)$ exists. A nearlattice S is said to have the three property if, for any $x, y, z \in S$, $x\vee y\vee z$ exists whenever $x\vee y$, $y\vee z$ and $z\vee x$ exist. Nearlattices with the three property were discussed by Evans in [31], where he referred to them as strong conditional lattices. It is easy to see that a nearlattice S has the three property if and only if it is medial.

Lemma 3.1. Every element of a medial nearlattice is sesqui-medial.

Proof. Suppose S is medial and n is any element of S . For any $x, y, z \in S$, $((x\lambda n)\vee(y\lambda n))\wedge((y\lambda n)\vee(z\lambda n))$, $x\lambda y \notin m(x, n, y)$ and $((x\lambda n)\vee(y\lambda n))\wedge((y\lambda n)\vee(z\lambda n))$, $y\lambda z \notin m(y, n, z)$. Thus using the upper bound property and the three property of S , $((x\lambda n)\vee(y\lambda n))\wedge((y\lambda n)\vee(z\lambda n))\vee(x\lambda y)\vee(y\lambda z) = J_n(x, y, z)$ exists in S . \square

Suppose S is a medial nearlattice and $a, b, c \in S$. If $a\vee b$, $b\vee c$, $c\vee a$ exists, we define $m^d(a, b, c) = (a\vee b)\wedge(b\vee c)\wedge(c\vee a)$. Of course, when S is distributive, $m^d(a, b, c) = m(a, b, c)$. For a fixed element n of S , let us introduce a ternary operation M_n , defined by $M_n(x, y, z) = m^d(x\lambda n, y\lambda n, z\lambda n)\vee m(x, y, z)$; $x, y, z \in S$. Notice that $m^d(x\lambda n, y\lambda n, z\lambda n)$ always exists in S . But also we have:

Lemma 3.2. In a medial nearlattice S with $n \in S$, $M_n(x, y, z)$ always exists for all $x, y, z \in S$.

Proof. Notice that $m^d(x\lambda n, y\lambda n, z\lambda n)$, $x\lambda y \notin m(x, n, y)$, $m^d(x\lambda n, y\lambda n, z\lambda n)$, $y\lambda z \notin m(y, n, z)$ and $m^d(x\lambda n, y\lambda n, z\lambda n)$, $z\lambda x \notin m(z, n, x)$. Then by the upper bound property and the three property both $m^d(x\lambda n, y\lambda n, z\lambda n)\vee(z\lambda x)$ and $m^d(x\lambda n, y\lambda n, z\lambda n)\vee(x\lambda y)\vee(y\lambda z)$ exist. Thus a second application of the three property yields the existence of $M_n(x, y, z)$. \square

Note that if n is nearly neutral in a nearlattice S , $M_n(x, y, z) = ((x\lambda n)\wedge(y\lambda n)\wedge(z\lambda n)\wedge n)\vee m(x, y, z)$, and when n is neutral, $M_n(x, y, z)\wedge n = (x\lambda n)\wedge(y\lambda n)\wedge(z\lambda n)\wedge n$. Also if S is a lattice and n is neutral, $M_n(x, y, z) = (m^d(x, y, z)\wedge n)\vee m(x, y, z) = m^d(x, y, z) \wedge (n\vee m(x, y, z))$.

Of course $m(x, y, z)$ and $M_n(x, y, z)$ are symmetric in x, y and z , whereas $j(x, y, z)$ and $J_n(x, y, z)$ are not. Thus, the operations

m and M_n are better behaved and easier to handle than the operations j and J_n respectively.

The following proposition is easily verifiable and so is given without proof.

Proposition 3.3. For an element n of a medial nearlattice S , $M_n(x,y,z) = m(x,y,z)$ for all $x,y,z \in S$ if and only if $(n \setminus)$ is a distributive lattice.

Hence in a distributive medial nearlattice S , $M_n(x,y,z) = m(x,y,z)$ for all $x,y,z \in S$. \square

Now we present the following interesting result which extends Theorem 2.6.

Theorem 3.4. Suppose n is a neutral sesquimedial element of a nearlattice S . Then the following conditions are equivalent.

- (i) S is medial;
- (ii) S_n is a medial nearlattice and $m^{S_n}(x,y,z) = M_n(x,y,z)$ for all $x,y,z \in S$.

Moreover, (i) does not necessarily imply (ii) when n is merely nearly neutral.

Proof. (i) \Rightarrow (ii). Since n is neutral,

$$M_n(x,y,z) \wedge n = (x \cap y) \wedge (y \cap z) \wedge (z \cap x) \wedge n.$$

By [2, Th. 2.4],

$$M_n(x,y,z) \cong m(x,y,z)(\Theta_n)$$

and $x \cap y \cong x \wedge y(\Theta_n)$. Thus, $(x \cap y) \wedge (M_n(x,y,z)) \cong x \wedge y(\Theta_n)$. Similarly,

$$(y \cap z) \wedge M_n(x,y,z) \cong y \wedge z(\Theta_n),$$

and

$$(z \cap x) \wedge M_n(x,y,z) \cong z \wedge x(\Theta_n).$$

Then using the technique of the proof of (i) \Rightarrow (ii) in Theorem 2.6, we obtain $\langle x \cap y, y \cap z, z \cap x \rangle_n = \langle M_n(x,y,z) \rangle_n$, and (ii) follows from the isomorphism of $(S_n; \subseteq)$ and $(P_n(S); \subseteq)$.

(ii) \Rightarrow (i). Adjoin a new 0 in S and form $(S;0)_n$. Then by 2.10, 0 is neutral and medial in $(S;0)_n$. Thus $(S;0)_n$ is medial as S_n is medial. Hence, by (i) \Rightarrow (ii), $((S;0)_n)_0$ is medial. But $((S;0)_n)_0 = (S;0)$ by 2.10, and so S is medial as required.

For the final assertion consider the lattice of Figure 1, where n is nearly neutral but not neutral. But its isotope,

given by Figure 2 is not medial. \square

It is well-known by Kolibiar [6] that if L is a lattice with 0 and 1 and n is central in it, then L_n is also a bounded lattice with n and n' as the smallest and the largest elements respectively, where $x \vee y = m(x, n', y)$ for all $x, y \in L$.

An element n in a lattice L is called central if it is neutral and complemented in each interval containing it.

We conclude this paper with the following extension of Kolibiar's result.

Proposition 3.5. Suppose L is a lattice and $n \in L$ is standard. Then the isotope L_n is a lattice if and only if n is central in L .

Proof. Since n is standard, $(L; \subseteq)$ and $(P_n(L); \subseteq)$ are isomorphic by 2.2. Thus, L_n is a lattice if and only if $P_n(L)$ is a lattice, i.e. if and only if n is complemented in each interval containing it. Consequently, the result follows by [2, Th. 3.5].




Figure 1




Figure 2

References

- [1] William H. CORNISH and R.C. HICKMAN: Weakly distributive semilattices, Acta Math. Acad. Sci. Hungar. 32(1978), 5-16.
- [2] William H. CORNISH and A.S.A. NOOR: Standard elements in a nearlattice, Bull. Austral. Math. Soc. 26(1982), 185-213.
- [3] E. Evans: Median lattices and convex subalgebras, Colloq. Math. Soc. Janos Bolyai, 29, North Holland Amsterdam, New York 1982, 225-240.
- [4] Robert HICKMAN: Join algebras, Commun. Alg. 8(1980), 1653-1685.
- [5] J. JAKUBÍK and M. KOLIBIAR: On some properties of a pair

- of lattices (Russian), Czechoslovak Math. J. 4(1954), 1-27.
- [6] Milan KOLIBIAR: A ternary operation in lattices (Russian), Czechoslovak Math. J. 6(1956), 318-329.
- [7] A.S.A. NOOR: Isotopes of nearlattices, Ganit. J. Bangladesh Math. Soc. 1(1)(1981), 63-72.
- [8] M. SHOLANDER: Medians, lattices and trees, Proc. Amer. Math. Soc. 5(1954), 808-812.

Department of Mathematics
Rajshahi University
Rajshahi
Bangladesh

School of Mathematical Sciences
Flinders University
Bedford Park
South Australia 5042
Australia

(Oblatum 19.11. 1986)