

Ofelia Teresa Alas

On the number of compact subsets in topological groups

Commentationes Mathematicae Universitatis Carolinae, Vol. 28 (1987), No. 3, 565--568

Persistent URL: <http://dml.cz/dmlcz/106568>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1987

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON THE NUMBER OF COMPACT SUBSETS
IN TOPOLOGICAL GROUPS

O.T. ALAS

Abstract: Results on the number of compact subsets in topological groups are proved. Examples are provided.

Key words: Pseudocharacter, boundedness number, weak Lindelöf number.

Classification: 54A25

Notation and terminology. Let (G, τ) be a nondiscrete Hausdorff group, e be its neutral element and \mathcal{K} denote the set of all compact subsets of G . For any set X , $|X|$ denotes the cardinality of X ; and, for any topological space X , $\psi(X)$, $\chi(X)$, $w(X)$, $c(X)$, $wL(X)$ denote the pseudocharacter, character, weight, cellularity, weak Lindelöf number of X , respectively.

1. Number of compact subsets

Definition (due to I. Juhász). The boundedness number of (G, τ) - denoted by $bo(G)$ - is the smallest infinite cardinal number \aleph such that for any open neighborhood V of e , there is a subset A of G , with $|A| \leq \aleph$, so that $V \cdot A = G$.

Notice that this notion is different from total- β -boundedness introduced by Comfort in [3].

Theorem 1. The following inequalities hold $\psi(G) \leq |G| \leq |\mathcal{K}| \leq bo(G)^{\psi(G)}$.

Proof. There is a collection of open symmetric neighborhoods of e , \mathcal{V} , such that $|\mathcal{V}| = \psi(G)$ and $\bigcap \{V \mid V \in \mathcal{V}\} = \{e\}$. For each $V \in \mathcal{V}$ fix a subset A_V of G such that $V \cdot A_V = G$ and $|A_V| \leq bo(G)$. Now the proof follows the one which appears in [1], since $\bigcap_{V \in \mathcal{V}} \left(\bigcup_{x \in A_V} Vx \cdot Vx \right) = \Delta$, the diagonal of $G \times G$,

Partially supported by CNPq.

$|G| \leq \text{bo}(G)^{\psi(G)}$ and any compact subset of G has density not bigger than $\psi(G)$.

Remarks. 1) As a matter of fact, the proof above shows that the set of all closed subsets of G whose densities do not exceed $\psi(G)$ has cardinality not bigger than $\text{bo}(G)^{\psi(G)}$.

2) It is easy to see that $\text{bo}(G) \leq \text{wL}(G) \leq \text{c}(G)$ (hence, $\text{bo}(G)^{\psi(G)} = \text{wL}(G)^{\psi(G)} = \text{c}(G)^{\psi(G)}$) and $\text{w}(G) = \text{bo}(G) \cdot \chi(G)$.

3) If $\text{bo}(G)$ is either a successor cardinal or a singular cardinal, then $\text{o}(G)^{\text{bo}(G)} = \text{o}(G)$, where $\text{o}(G)$ denotes the number of open sets in G .

Corollary 1. If $\text{bo}(G) \leq 2^{\psi(G)}$, then $\psi(G) \leq |\mathcal{K}| \leq 2^{\psi(G)}$.

Lemma. If K is a nonempty compact subset of G , then $\psi(K, G) \leq \psi(G)$.

Proof. Let \mathcal{V} be a collection of symmetric open neighborhoods of e , closed under finite intersections. Furthermore we shall assume that $|\mathcal{V}| = \psi(G)$ and $\bigcap \{\text{cl}(V) \mid V \in \mathcal{V}\} = \{e\}$. Then $\bigcap \{V, K \mid V \in \mathcal{V}\} = K$; indeed, let $y \notin K$, then there is $V \in \mathcal{V}$ such that $V \cap K = \emptyset$ (otherwise, $V \cap K \neq \emptyset, \forall V \in \mathcal{V}$ and since K is compact, $\bigcap \{\text{cl}(V) \cap K \mid V \in \mathcal{V}\}$ would be nonempty, which is impossible). But if $V \cap K = \emptyset$, then $y \notin V, K$.

Corollary 2. If $\text{bo}(G) \leq 2^{\psi(G)}$ and there is a compact subset K of G such that $\psi(K, G) < \psi(G)$, then $|\mathcal{K}| = 2^{\psi(G)}$.

Proof. If there is a nonempty compact subset K of G such that $\psi(K, G) < \psi(G)$, then for each $x \in K$, $\psi(G) = \psi(x, G) \leq \psi(x, K) \cdot \psi(K, G)$. It follows from Čech-Pospíšil's theorem that $|K| \geq 2^{\psi(G)}$, hence $|\mathcal{K}| = 2^{\psi(G)}$.

Theorem 2. (GCH) If G is pseudocompact, then $|\mathcal{K}|^{\aleph_0} = |\mathcal{K}|$.

Proof. Since G is infinite and pseudocompact, then $|G| \geq 2^{\aleph_0}$. We may assume that if α is a cardinal number such that $\alpha \geq 2^{\aleph_0}$, $\text{cf}(\alpha) \neq \aleph_0$, then $\alpha^{\aleph_0} = \alpha$.

From Theorem 1 and since $\text{bo}(G) = \aleph_0$, either $|\mathcal{K}| = 2^{\psi(G)}$ or $|G| = |\mathcal{K}| = \psi(G)$. In the first case there is nothing to prove; let us consider that $|G| = |\mathcal{K}| = \psi(G)$. From van Douwen's theorem 1.1 ([4]) if $\text{cf}(|G|) = \aleph_0$, there is a cardinal $\mu < |G|$, such that $\psi(G) \leq \text{w}(G) \leq 2^{\mu}$. But $|G| \geq 2^{\mu}$, hence $|G| = |\mathcal{K}| = 2^{\mu}$ and the proof is completed.

Lemma. If V is an open symmetric neighborhood of e and $\mathcal{K}(\text{cl}(V))$ denotes the set of all compact subsets of $\text{cl}(V)$, then $|\mathcal{K}| = \text{bo}(G) \cdot |\mathcal{K}(\text{cl}(V))|$.

Proof. It is immediate that $|\mathcal{K}| \geq \text{bo}(G)$ and $|\mathcal{K}| \geq |\mathcal{K}(\text{cl}(V))|$: On the other hand, let B be a subset of G such that $\text{bo}(G) \geq |B|$ and $V \cdot B = G$. For each nonempty finite subset F of B let \mathcal{K}_F denote the set of all compact subsets of G contained in $V \cdot F$. The function from \mathcal{K}_F into $\prod \{\mathcal{K}(\text{cl}(V)y) \mid y \in F\}$ which assigns to each $K \in \mathcal{K}_F$ the point $(\text{cl}(V) \cap K)_{y \in F}$ is injective. But $\mathcal{K} = \cup \{\mathcal{K}_F \mid \emptyset \neq F \subset B, \text{ finite}\}$ and $|\mathcal{K}(\text{cl}(V))| = |\mathcal{K}(\text{cl}(V))|$, hence $|\mathcal{K}| \leq \text{bo}(G) |\mathcal{K}(\text{cl}(V))|$, which completes the proof.

Remark. The GCH cannot be avoided in Theorem 2, since I. Juhász, under CH and using forcing arguments, obtained an HFD subgroup of $\{0,1\}^{\omega_1}$, such that $|\mathcal{K}|^{\aleph_0} \neq |\mathcal{K}|$.

2. Examples

Example 1. ([5] or [2], page 1170.) Under $\aleph_1 = 2^{\aleph_0}$ and $\aleph_2 < 2^{\aleph_1}$, there is a hereditarily separable pseudocompact group G with $|G| = |\mathcal{K}| = \aleph_2$ (which is not a power of 2, but $\aleph_2^{\aleph_0} = \aleph_2$).

Example 2. Let G be the topological subgroup of $\{0,1\}^{\omega_1}$ whose members are the $(x_\alpha)_{\alpha \in \omega_1}$ such that $\{\alpha \in \omega_1 \mid x_\alpha = 1\}$ is countable. G is countably compact, $\psi(G) = \aleph_1$, $|G| = 2^{\aleph_0}$ and $|\mathcal{K}| = 2^{\aleph_1}$. (Notice that the set $\{(x_\alpha)_{\alpha \in \omega_1} \in G \mid x_\alpha = 1 \text{ for at most one } \alpha \in \omega_1\}$ is compact and has just one accumulation point.)

Example 3. Let us consider $\{0,1\}^{\omega_1}$ with the G_σ -topology (each factor with the discrete topology). For each $\beta \in \omega_1$, let y_β be the point $(x_\alpha)_{\alpha \in \omega_1}$ such that $x_\alpha = 1, \forall \alpha < \beta$ and $x_\alpha = 0$, otherwise. Denote by G the topological subgroup generated by the y_β . Then $\text{bo}(G) = \aleph_0$ and $w(G) = \aleph_1$. (Notice that no countable subcollection of $\{\text{pr}_\xi^{-1}(\{0\}) \mid \xi \in \omega_1\}$, where pr_ξ denotes the projection for each $\xi \in \omega_1$, has its union dense in G .)

Example 4. Let α be an infinite cardinal such that $\alpha^{\aleph_0} = \alpha$. Comfort proved that there is a dense countably compact subgroup G_* of $\{0,1\}^{2^\alpha}$ such that $|G_*| = \alpha$. Denoting by G the topological product group $\sum \times G_*$, where \sum denotes the subgroup of $\{0,1\}^\alpha$ such that its elements have at most countably many coordinates different from 0, we have that $|G| = \alpha$, $\psi(G) = \alpha$ and $w(G) = |\mathcal{K}| = 2^\alpha$. Notice that G is countably compact.

Example 5. Let α be an infinite cardinal number, whose cofinality is β and let $(\alpha_i)_{i \in \beta}$ be a strictly increasing family of cardinals such that $\alpha = \sup_{i \in \beta} \alpha_i$. For each $i \in \beta$ let G_i be a discrete topological group with $|G_i| = \alpha_i$. The topological product group $G = \prod_{i \in \beta} G_i$ has a boundedness number equal to α , $\psi(G) = \beta$ and $|K| = \alpha^\beta$.

References

- [1] O.T. ALAS: Inequalities with topological cardinal invariants, Collected papers dedicated to Prof. Edison Farah on the occasion of his retirement (1982), 91-97.
- [2] W.W. COMFORT: Topological groups, Handbook of set-theoretic topology, North-Holland, Amsterdam (1984), 1145-1263.
- [3] W.W. COMFORT and D.L. GRANT: Cardinal invariants, pseudocompactness and minimality: some recent advances in the topological theory of topological groups, Top. Proc. 6(1981), 227-265.
- [4] E. van DOUWEN: The weight of a pseudocompact (homogeneous) space whose cardinality has countable cofinality, Proc. Amer. Math. Soc. 80(1980), 678-682.
- [5] A. HAJNAL and I. JUHÁSZ: A separable normal topological group need not be Lindelöf, Gen. Top. Appl. 6(1976), 199-205.
- [6] I. JUHÁSZ: Cardinal functions in topology - ten years later, Mathematical Centre Tracts 123, Amsterdam, (1980).

Instituto de Matemática e Estatística, Caixa Postal 20570 (Ag. IGUATEMI),
Sao Paulo, Brazil

(Oblatum 31.3. 1987)