

Ryszard Engelking

An elementary proof of Noble's theorem on normality of powers

Commentationes Mathematicae Universitatis Carolinae, Vol. 29 (1988), No. 4, 677--678

Persistent URL: <http://dml.cz/dmlcz/106683>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1988

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

AN ELEMENTARY PROOF OF NOBLE'S THEOREM
ON NORMALITY OF POWERS

Ryszard ENGELKING

Dedicated to Professor M. Katětov on his seventieth birthday

Abstract: We show in a simple way that if all powers of a space are normal, then the space itself is compact.

Key words: Cartesian product, normality, compactness.

Classification: 54B10, 54D15, 54D30

One of the important results in the theory of normality of Cartesian products, originated in 1948 by M. Katětov and A.H. Stone (see [2] and [6]), is the theorem due to N. Noble [4] which states that if all powers of a space are normal, then the space itself is compact. The theorem has been originally obtained in the frame of a general theory developed by N. Noble, and this prompted several authors to propose simpler and more direct proofs (see [1], [3] and [5]). In all these proofs A.H. Stone's theorem on the non-normality of N^{\aleph_1} is applied and, together with a conveniently chosen rather strong topological result, yields Noble's theorem.

It turns out that the Noble theorem can also be established in an elementary way by a variant of the argument A.H. Stone used to prove the non-normality of N^{\aleph_1} .

We shall show that if for a topological space X the power X^m is normal for every m , then X is compact.

Suppose that X is not compact and consider a family $\{F_s\}_{s \in S}$ of closed subsets of X which has the finite intersection property and an empty intersection; denote by m the cardinality of S . The set $F = \prod_{s \in S} F_s \subset X^m = \prod_{s \in S} X_s$, where $X_s = X$ for $s \in S$, is closed and disjoint from the diagonal $\Delta \subset X^m$. Consider an open set U containing F .

Let x_1 be an arbitrary point in F . There exists a finite set $S_1 \subset S$ such

that $p_{S_1}^{-1} p_{S_1}(x_1) \subset U$. Define a point $x_2 \in F$ by letting $p_s(x_2) = a_1$ for $s \in S_1$, where a_1 is an arbitrary point in $\bigcap_{s \in S_1} F_s$, and $p_s(x_2) = p_s(x_1)$ for $s \notin S_1$, and enlarge S_1 to a finite set $S_2 \subset S$ such that $p_{S_2}^{-1} p_{S_2}(x_2) \subset U$. By induction we can define points x_1, x_2, x_3, \dots in F , finite sets $S_1 \subset S_2 \subset S_3 \subset \dots \subset S$ and points a_1, a_2, a_3, \dots in X such that

$$p_s(x_n) = a_{n-1} \text{ for } s \in S_{n-1} \text{ and } p_{S_n}^{-1} p_{S_n}(x_n) \subset U.$$

Since, by A.H. Stone's theorem, X does not contain a closed copy of N , there exists a point $a_0 \in X$ every neighbourhood of which contains infinitely many a_n 's. The points y_1, y_2, y_3, \dots of X^m defined by

$$p_s(y_n) = p_s(x_n) \text{ for } s \in S_n \text{ and } p_s(y_n) = a_0 \text{ for } s \notin S_n$$

belong to U and - as one easily sees - every neighbourhood of the point $y \in \Delta$ all of whose coordinates are equal to a_0 , contains a y_n . Thus $\Delta \cap \bar{U} \neq \emptyset$; since this is in contradiction with the normality of X^m , it follows that X^m is compact.

References

- [1] S.P. FRANKLIN and R.C. WALKER: Normality of powers implies compactness, Proc. Amer. Math. Soc. 36(1972), 295-296.
- [2] M. KATĚTOV: Complete normality of Cartesian products, Fund. Math. 36 (1948), 271-274.
- [3] J. KEESLING: Normality and infinite product spaces, Adv. in Math. 9 (1972), 90-92.
- [4] N. NOBLE: Products with closed projections II, Trans. Amer. Math. Soc. 160(1971), 169-183.
- [5] L. POLKOWSKI: On N. Noble's theorems concerning powers of spaces and mappings, Coll. Math. 41(1979), 215-217.
- [6] A.H. STONE: Paracompactness and product spaces, Bull. Amer. Math. Soc. 54(1948), 977-982.

Institute of Mathematics, University of Warsaw, 00-901 Warsaw, PKiN IX p.
Poland

(Oblatum 24.5. 1988)