

Heikki J. K. Junnila

Around Katětov's metrization theorem

Commentationes Mathematicae Universitatis Carolinae, Vol. 29 (1988), No. 4, 685--694

Persistent URL: <http://dml.cz/dmlcz/106685>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1988

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

AROUND KATĚTOV'S METRIZATION THEOREM

Heikki J.K. Junnila

Dedicated to Professor Miroslav Katětov on his seventieth birthday

ABSTRACT. We discuss some results related to the theorem of Katětov that a compact Hausdorff space X is metrizable if, and only if, X^3 is hereditarily normal, and we prove that X is metrizable if, and only if, X^ω is hereditarily metanormal.

AMS Classification 54E35 54D30 54D18

1. *INTRODUCTION.*

In 1948, M. Katětov [K] proved that a compact Hausdorff space X is metrizable if X^3 is hereditarily normal. Even though this theorem is not very difficult to prove, it is a remarkable result: it provides a topological characterization of metrizability which involves no explicit countability condition whereas the earlier topological metrization theorems required the existence of some kind of a countable structure.

In his paper, Katětov raised the question whether hereditary normality of X^2 , for a compact Hausdorff space X , is enough to make X metrizable. In 1978, P.J. Nyikos [N1] showed that it is consistent with ZFC that the answer to Katětov's question is "no": using the assumption that Martin's Axiom and the negation of the Continuum Hypothesis hold, he provided an example of a non-metrizable compact Hausdorff space X such that the space X^2 is hereditarily normal; recently, G. Gruenhage has obtained an example of such a space under the assumption that the Continuum Hypothesis holds. Both examples are described in detail in [GN]; this paper also contains information on the following problem, which still remains open.

Problem 1 Is it consistent with ZFC that every compact space whose square is hereditarily normal is metrizable?

After Katětov's result, there have appeared other results which take the form: if X is a compact Hausdorff space and X^n has hereditarily some given property, then X is metrizable. Even some earlier results can be presented in that form: Šneĭder proved in 1945 [Š] that a compact Hausdorff space X is metrizable if the diagonal is a G_δ -set in X^2 , and it follows from this result that X is metrizable provided that X^2 is hereditarily Lindelöf. The last-mentioned result was strengthened in 1984 by Gruenhage [G], who proved that a compact Hausdorff space is metrizable provided that X^2 is hereditarily paracompact; under Martin's Axiom and the negation of the Continuum Hypothesis, hereditary paracompactness can be weakened to hereditary collectionwise normality, as was shown by Nyikos in 1981 [N2]. In 1971 P. Zenor [Z] had proved that a compact Hausdorff space X is metrizable provided that X^3 is hereditarily countably paracompact.

Other properties of a compact Hausdorff space X , besides metrizability, can be characterized in terms of hereditary properties of some powers of X . For example, Gruenhage [G] proved that Eberlein compactness of X is equivalent with X^2 being hereditarily σ -metacompact and Corson compactness of X is equivalent with X^2 being hereditarily meta-Lindelöf. There is still much work to do on this area: for example, the following problem, raised in [G], remains open.

Problem 2 Characterize those compact Hausdorff spaces whose square is hereditarily metacompact.

Hereditary metacompactness of X^2 does not characterize Eberlein compact spaces: N.N. Yakovlev [Y] has shown that for the one-point compactification $A(\omega_1)$ of the discrete space on ω_1 , the (Eberlein compact) space $A(\omega_1)^\omega$ is not hereditarily metacompact.

2. A METRIZATION THEOREM.

As mentioned in the introduction, a compact Hausdorff X space is metrizable provided that X^3 is either hereditarily normal or hereditarily countably paracompact. In this section we shall show that metrizability of X follows if the infinite power X^ω satisfies hereditarily a property significantly weaker than normality and countable paracompactness. The property to be considered was introduced by E.K. van Douwen.

Definition 1 [vD] A topological space X is *metanormal* provided that for every discrete family $\{F_n : n \in \omega\}$ of closed subsets of X there exists a family $\{L_n : n \in \omega\}$ of G_δ -subsets of X such that $\bigcap_{n \in \omega} L_n = \emptyset$ and, for each $n \in \omega$, $F_n \subset L_n$.

Note that, besides all normal spaces, all countably metacompact spaces are metanormal.

To prove that certain infinite powers are not hereditarily metanormal, we introduce the following concept.

Definition 2 Let F be a closed subset of a topological space X and let A be an uncountable subset of $X \setminus F$. We say that F *attracts* A provided that every uncountable subset B of A contains a countable subset C such that $Cl(C) \cap F \neq \emptyset$.

If F attracts some subset of X , then we say that F is an *attractive subset* of X .

If the attractive set F consists of one point x , then we say that x is an *attractive point* of X .

Remark For later use, we mention the following alternative characterization of an uncountable set A attracted to a closed set F : whenever I is an uncountable set and $\{V_i : i \in I\}$ is a family of neighborhoods of the set F , then there are only countably many elements $a \in A$ such that the set $\{i \in I : a \in V_i\}$ is countable.

Typical attractive subsets of a compact space are exhibited in the following lemma.

Lemma 1 *The following hold for a subset A of a compact Hausdorff space X :*

1° *If A is uncountable and relatively discrete, then A is attracted by the set $F = Cl(A) \setminus A$.*

2° *Assume that $A = \{x_\alpha : \alpha \in \omega_1\}$ and the following conditions hold:*

(a) $x_\alpha \neq x_\beta$ whenever $\alpha \neq \beta$,

(b) A contains no uncountable relatively discrete subset, and

(c) for each $\beta \in \omega_1$, the set $\{x_\alpha : \alpha \leq \beta\}$ is open in A .

Then A is attracted to the set $F = \bigcap_{\beta \in \omega_1} Cl\{x_\alpha : \alpha \geq \beta\}$.

Proof. 1° follows from the observation that, under the assumption made in 1°, we have that $Cl(B) \cap F \neq \emptyset$ for every infinite $B \subset A$.

2° Since X is compact, we have that $F \neq \emptyset$. Note that it follows from (b) and (c) that A is hereditarily separable. Let C be a countable dense subset of A . Then $F \subset Cl(A) = Cl(C)$ and hence $Cl(C) \cap F \neq \emptyset$. A similar argument shows that every uncountable set $B \subset A$ contains a countable set D with $Cl(D) \cap F \neq \emptyset$. \square

Proposition 1 *A compact Hausdorff space X is hereditarily Lindelöf if, and only if, X contains no attractive set.*

Proof. Necessity of the condition follows directly, since every closed subset of a regular hereditarily Lindelöf space is a G_δ -set.

To prove sufficiency, assume X is not hereditarily Lindelöf. Then X has a subset $A = \{x_\alpha : \alpha \in \omega_1\}$ such that $x_\alpha \neq x_\beta$ whenever $\alpha \neq \beta$ and, for each $\beta \in \omega_1$, the set $\{x_\alpha : \alpha \leq \beta\}$ is open in A . If A contains an uncountable relatively discrete set, then by part 1° of Lemma 1, X contains an attractive set. On the other hand, if every relatively discrete subspace of A is countable, then it follows from part 2° of Lemma 1 that X again contains an attractive set. \square

Now we show that a space can have no attractive points if its countably infinite power is hereditarily metanormal.

Proposition 2 *Let p be an attractive point of a topological space X . Then the subspace $X^\omega \setminus \{p\}^\omega$ of X^ω is not metanormal.*

Proof. Denote the subspace in question by Y . For each $k \in \omega$, let

$$S_k = \{y \in X^\omega : y(k) \neq p \text{ and } y(m) = p \text{ for every } m \neq k\}.$$

Note that $\{S_k : k \in \omega\}$ is a discrete family of closed subsets of Y . Partition the set ω into infinite pieces A_n , $n \in \omega$; for every $k \in \omega$, let $n_k \in \omega$ be defined by the condition that $k \in A_{n_k}$. For every $n \in \omega$, let $F_n = \bigcup_{k \in A_n} S_k$. It follows from the corresponding property of the family $\{S_k : k \in \omega\}$ that $\{F_n : n \in \omega\}$ is a discrete family of closed subsets of Y . We show that there is no family of G_δ -sets in Y as required in the definition of a metanormal space.

For every $n \in \omega$, let L_n be a G_δ -subset of Y such that $F_n \subset L_n$. We show that $\bigcap_{n \in \omega} L_n \neq \emptyset$. For every $n \in \omega$, let G_{ni} , $i \in \omega$, be open subsets of Y such that $\bigcap_{i \in \omega} G_{ni} = L_n$ and, for every $i \in \omega$, $G_{n(i+1)} \subset G_{ni}$.

Let B be an uncountable subset of $X \setminus \{p\}$ which is attracted to p . For all $b \in B$ and $k \in \omega$, denote by y_{bk} that point of Y , whose k^{th} coordinate is b and all the other coordinates equal p . Note that $y_{bk} \in S_k \subset F_{n_k} \subset G_{n_k k}$ and hence there exist neighborhoods V_{bkj} , $j \neq k$, of p in X such that

$$\left[\prod_{j < k} V_{bkj} \right] \times \{b\} \times \left[\prod_{j > k} V_{bkj} \right] \subset G_{n_k k}.$$

Let us now construct a point y which belongs to the set $\bigcap_{n \in \omega} L_n$. By induction on $k \in \omega$, we define points $y_k \in B$ and uncountable sets $E_{kj} \subset B$, for $j > k$, as follows.

By the remark following Definition 2, there exists $y_0 \in B$ such that, for every $j > 0$, the set $D_{0j} = \{b \in B : y_0 \in V_{b_0j}\}$ is uncountable. Note that, for every $j > 0$, the set $E_{0j} = D_{0j} \cap V_{y_0 0j}$ is uncountable.

Let $k > 0$ be such that the points y_l and the uncountable sets E_{lj} have already been defined for all $l < k$ and $j > l$. Then there exists $y_k \in E_{k-1k}$ such that, for each $j > k$, the set $D_{kj} = \{b \in E_{k-1j} : y_k \in V_{bjk}\}$ is uncountable. For each $j > k$, the set $E_{kj} = D_{kj} \cap V_{y_k kj}$ is uncountable, and this completes the inductive step. Note that it follows from the inductive construction that $E_{kj} \subset E_{lj}$ whenever $l < k < j$.

We show that the point $y = (y_k)_{k \in \omega}$ belongs to the set $\bigcap_{n \in \omega} L_n = \bigcap_{n \in \omega} \bigcap_{i \in \omega} G_{ni}$. Let $n \in \omega$ and $i \in \omega$. Then there exists $m \in A_n$ such that $m > i$. For each $k < m$, we have that $y_m \in E_{m-1m} \subset E_{km}$ and hence that $y_k \in V_{y_m mk}$. For each $k > m$, we have that $y_k \in F_{k-1k} \subset E_{mk} \subset V_{y_m mk}$. By the foregoing, we have that

$$y = (y_k)_{k \in \omega} \in \left[\prod_{k < m} V_{y_m mk} \right] \times \{y_m\} \times \left[\prod_{k > m} V_{y_m mk} \right] \subset G_{n_m m} = G_{n_m} \subset G_{ni}. \quad \square$$

Remark A straightforward extension of the above proof shows that the space $Y = X^\omega \setminus \{p\}^\omega$ not only fails to be metanormal, but it does not even have the following weaker property (which should be called *orthonormality*, to conform with van Douwen's terminology): for every discrete family $\{F_n : n \in \omega\}$ of closed subsets and for all open sets

O_n , $n \in \omega$, such that $F_n \subset O_n$ for every $n \in \omega$, there exist G_δ -sets L_n , $n \in \omega$, such that the set $\bigcap_{n \in \omega} L_n$ is open and, for every $n \in \omega$, $F_n \subset L_n \subset O_n$. Note that, besides metanormal spaces, all countably orthocompact spaces satisfy this weaker property.

We shall now indicate a slight extension of Proposition 2.

Corollary *Let K be a compact attractive subset of a topological space X . Then the subspace $X^\omega \setminus K^\omega$ of X^ω is not metanormal.*

Proof. Let Z be the space obtained from X by identifying the set K to a point p , and let f be the corresponding quotient mapping. Note that p is an attractive point of Z ; hence it follows from Proposition 2 that the subspace $Z^\omega \setminus \{p\}^\omega$ of Z^ω is not metanormal.

Since K is compact, f is a perfect mapping. By a theorem of Z. Frolík [F] and N. Bourbaki [B], the "product mapping" $\bar{f} = \langle f, f, \dots \rangle$ from X^ω onto Z^ω is a perfect mapping. We have that

$$\bar{f}^{-1}(Z^\omega \setminus \{p\}^\omega) = X^\omega \setminus K^\omega,$$

and it follows that the restriction of \bar{f} to the subspace $X^\omega \setminus K^\omega$ of X^ω is a perfect mapping onto $Z^\omega \setminus \{p\}^\omega$. It is easy to see that metanormality is preserved under closed mappings. It follows, since $Z^\omega \setminus \{p\}^\omega$ is not metanormal, that neither is $X^\omega \setminus K^\omega$. \square

Using the above corollary, we can easily prove our main result.

Theorem 1 *A compact Hausdorff space X is metrizable if, and only if, the space X^ω is hereditarily metanormal.*

Proof. Necessity of the condition is trivial. To prove sufficiency, assume that X^ω is hereditarily metanormal. Since X^ω is homeomorphic with $(X^2)^\omega$, the latter space is hereditarily metanormal, and it follows from the previous corollary that X^2 contains no attractive set. By Proposition 1, X^2 is hereditarily Lindelöf. It follows by Šneřder's metrization theorem [Š] that X is metrizable. \square

Corollary *A compact Hausdorff space is metrizable if, and only if, the space X^ω is hereditarily countably metacompact.*

We close this section with some more consequences of Proposition 2 and its proof.

Note that, in the proof of Proposition 2, we actually showed that if the space X contains an attractive point p , then $X^\omega \setminus \{p\}^\omega$ contains a family $\{F_n : n \in \omega\}$ of subsets which is closed and discrete in the relative product topology but for which there exist no family $\{L_n : n \in \omega\}$ of G_δ -sets in the relative box topology such that $\bigcap_{n \in \omega} L_n = \emptyset$ and, for every $n \in \omega$, $F_n \subset L_n$; in particular, the box topology of X^ω is not hereditarily metanormal.

Since the one-point compactification $A(\omega_1)$ of the discrete space on ω_1 has the compactifying point as an attractive point, we get the following.

Example $A(\omega_1)^\omega$ is not hereditarily metanormal either in the product topology or in the box topology.

Since $A(\omega_1)^\omega$ (in the product topology) is an Eberlein compact space of weight ω_1 , it follows from results of D. Amir and J. Lindenstrauss (see [D], Chapter 5) that $A(\omega_1)^\omega$ can be embedded in the sequence-space $c_0(\omega_1)$, when the latter space is equipped with the topology of pointwise convergence. On the other hand, if X is any non-ccc compact Hausdorff space, then we can embed $c_0(\omega_1)$ in the space $C_p(X)$ (the set $C(X)$ equipped with the topology of pointwise convergence). As a consequence, we have the following result.

Proposition 3 *If X is a non-ccc compact Hausdorff space, then the space $C_p(X)$ contains a non-metanormal subspace with compact closure.*

In connection with the above result, we should recall the result of H.P. Rosenthal [R] that if X is a ccc compact Hausdorff space, then every compact subspace of $C_p(X)$ is metrizable.

3. THE NON-COMPACT CASE.

The metrization theorems of Katětov and Zenor mentioned in the introduction do not remain true for non-compact spaces. However, for general spaces, the following result obtains.

Theorem [K],[Z] *The following conditions are mutually equivalent for every topological space X :*

- 1° X^ω is perfectly normal.
- 2° X^ω is hereditarily normal.
- 3° X^ω is hereditarily countably paracompact.

The above result might lead one to conjecture that a topological space X is perfect provided that X^ω is hereditarily metanormal or hereditarily countably metacompact. However, this result does not hold even for Lindelöf spaces, as we shall now indicate.

Let $L(\omega_1)$ be the "one-point Lindelöfization" of the discrete space on ω_1 , with the ordinal ω_1 as the "Lindelöfizing" point. Then the neighborhoods of ω_1 in $L(\omega_1)$ are the sets with countable complement while the points $\alpha < \omega_1$ are isolated. The subset $\{\omega_1\}$ is not a G_δ -set in $L(\omega_1)$ and hence $L(\omega_1)$ is not a perfect space.

We shall show that $L(\omega_1)^\omega$ satisfies a base property which implies that $L(\omega_1)^\omega$ is hereditarily metacompact. Recall that a base \mathcal{B} of a space X is of *subinfinite rank* provided that for every infinite subfamily \mathcal{C} of \mathcal{B} , if $\bigcap \mathcal{C} \neq \emptyset$, then \mathcal{C} contains two distinct sets which are related by inclusion. The base \mathcal{B} is *Noetherian* provided that the poset (\mathcal{B}, \subset) has no infinite increasing chains.

Proposition 4 $L(\omega_1)^\omega$ has a Noetherian base of subinfinite rank.

Proof. Let n, f and α be such that $n \in \omega$, $\alpha \in \omega_1$, f is a mapping with $Dom(f) \subset \{0, \dots, n\}$ and $Im(f) \subset \omega_1$, and $\alpha > \max Im(f)$. Then define

$$V_{n,f,\alpha} = \{x \in L(\omega_1)^\omega : f \subset x \text{ and } x(k) \geq \alpha \text{ for every } k \in \{0, \dots, n\} \setminus Dom(f)\}.$$

It is easy to see that the collection of all such $V_{n,f,\alpha}$ forms a base for the topology of $L(\omega_1)^\omega$; we denote this base by \mathcal{B} .

Note that, for all n, f, α and m, g, β , we have that $V_{n,f,\alpha} \subset V_{m,g,\beta}$ if, and only if, $m \leq n$, $g \subset f$ and $\beta \leq \alpha$. Using this observation, it is easy to see that \mathcal{B} is Noetherian.

To show that \mathcal{B} is of subinfinite rank, let $x \in L(\omega_1)^\omega$ and n_k, f_k and α_k , $k \in \omega$, be such that $x \in \bigcap_{k \in \omega} V_{n_k, f_k, \alpha_k}$. Then we can find $i \in \omega$ and $j \in \omega$ such that $i < j$, $n_i \leq n_j$, $\alpha_i \leq \alpha_j$ and $\max Im(f_i) \leq \max Im(f_j)$. We show that $V_{n_j, f_j, \alpha_j} \subset V_{n_i, f_i, \alpha_i}$.

Let us first show that $Dom(f_i) \subset Dom(f_j)$. Let $n \in Dom(f_i)$. Since $f_i \subset x$, we have that $f_i(n) = x(n)$. It follows that $x(n) \leq \max Im(f_i) \leq \max Im(f_j) < \alpha_j$, and it follows further, since $n \leq n_i \leq n_j$ and $x \in V_{n_j, f_j, \alpha_j}$, that $n \in Dom(f_j)$. We have shown that $Dom(f_i) \subset Dom(f_j)$. Since $x \in V_{n_i, f_i, \alpha_i} \cap V_{n_j, f_j, \alpha_j}$, we have that $f_i \cup f_j \subset x$, and it follows, since $Dom(f_i) \subset Dom(f_j)$, that $f_i \subset f_j$. Now the inclusion $V_{n_j, f_j, \alpha_j} \subset V_{n_i, f_i, \alpha_i}$ follows by the observation made in the preceding paragraph of this proof. We have shown that \mathcal{B} is of subinfinite rank. \square

W.F. Lindgren and Nyikos [LN] have observed that every space with a Noetherian base of subinfinite rank is (hereditarily) metacompact. Hence the following result obtains.

Corollary $L(\omega_1)^\omega$ is hereditarily metacompact.

Without proof, we mention another hereditary covering property enjoyed by the space $L(\omega_1)^\omega$: it is not difficult to show that this space is also hereditarily screenable.

Using a result of K. Alster, we obtain the following generalization of Proposition 4: if X is a scattered P-space of weight ω_1 , then X^ω has a Noetherian base of subinfinite rank. This follows from Proposition 4, because Alster showed in [A] that any Lindelöf space X with the stated properties can be embedded in $L(\omega_1)^\omega$, and an examination of Alster's proof shows that this result holds also for non-Lindelöf spaces.

REFERENCES

- [A] K. Alster, A class of spaces whose cartesian product with every hereditarily Lindelöf space is Lindelöf, *Fund. Math.* **114** (1981) 173–181.
- [B] N. Bourbaki, *Topologie Générale*, ch I & II (third ed.) Hermann, Paris, 1961.
- [D] J. Diestel, *Geometry of Banach Spaces—Selected Topics*, Lecture Notes in Math. **485** (Springer–Verlag, 1975).
- [vD] E.K. van Douwen, Covering and separation properties in box products, in: G.M. Reed, ed., *Surveys in General Topology*, Academic Press, New York, 1980.
- [F] Z. Frolík, On the topological product of paracompact spaces, *Bull. Polon. Acad. Sci., Sér. Sci. Math. Astr. Phys.* **8** (1960) 747–750.
- [G] G. Gruenhage, Covering properties on $X^2 \setminus \Delta$, W -sets and compact subsets of Σ -products, *Topology and its Appl.* **17** (1984) 287–304.
- [GN] G. Gruenhage and P.J. Nyikos, Normality in X^2 for compact X , preprint.
- [K] M. Katětov, Complete normality of cartesian products, *Fund. Math.* **35** (1948) 271–274.
- [LN] W.F. Lindgren and P.J. Nyikos, Spaces with bases satisfying certain order and intersection properties, *Pacific J. Math.* **66**(1976)455–476.
- [N1] P.J. Nyikos, A compact non-metrizable space P such that P^2 is completely normal, *Topology Proc.* **2** (1977) 359–364.
- [N2] P.J. Nyikos, Axioms, theorems, and problems related to the Jones lemma, in L.F. McAuley and M.M. Rao, eds., *General Topology and Modern Analysis*, Academic Press, New York, 1981, pp. 441–449.
- [R] H.P. Rosenthal, On injective Banach spaces and the spaces $L^\infty(\mu)$ for finite measures μ , *Acta Math.* **124**(1970)205–248.
- [Š] V.E. Šneĭder, Continuous images of Souslin and Borel sets. Metrization theorems. *Dokl. Akad. Nauk SSSR* **50** (1945) 77–79. (In Russian)
- [Y] N.N. Yakovlev, On bicomacta in Σ -products and related spaces, *Comment. Math. Univ. Carolinae* **21** (1980) 263–283.
- [Z] P. Zenor, Countable paracompactness in product spaces, *Proc. Amer. Math. Soc.* **30** (1971) 199–201.

Department of Mathematics
 University of Helsinki (Oblatum 1.7. 1988)
 Hallituskatu 15
 00100 Helsinki 10
 Finland