

Commentationes Mathematicae Universitatis Carolinae

Ghassan Alherk

On the non- $l_n^{(1)}$ and locally uniformly non- $l_n^{(1)}$ properties, and l^1 copies in Musielak-Orlicz spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 31 (1990), No. 3,
435--443

Persistent URL: <http://dml.cz/dmlcz/106879>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1990

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*
<http://project.dml.cz>

On the non- $l_n^{(1)}$ and locally uniformly non- $l_n^{(1)}$ properties, and l^1 copies in Musielak—Orlicz spaces

GHASSAN ALHERK

Abstract. It is proved that a Musielak—Orlicz space $L^\Phi(\mu)$ over a non-atomic measure is locally uniformly non- $l_n^{(1)}$ if and only if Φ satisfies the Δ_2 -condition. Moreover, there are given some criteria in order that Musielak—Orlicz space be non- $l_n^{(1)}$ as well as in order that it contains an isometric copy of l^1 . These results generalize the results of [1], [2], [4] and [8].

Keywords: Non- $l_n^{(1)}$ space, locally uniformly non- $l_n^{(1)}$ space, Musielak—Orlicz space, Luxemburg norm, Δ_2 -condition

Classification: 46E30, 46B25

1. Introduction.

At the beginning, let us give some terminology and definitions concerning Musielak—Orlicz spaces and geometry of Banach spaces. In the whole paper, (T, Σ, μ) denotes a positive non-atomic measure space, N denotes the set of all natural numbers, R denotes the reals, R_+ denotes the non-negative reals, χ_A denotes the characteristic function of a set $A \in \Sigma$.

A function $\Phi : T \times R \rightarrow [0, +\infty]$ is said to be a Musielak—Orlicz function if $\Phi(t, \cdot)$ is even, convex, vanishing and continuous at zero, left continuous on the whole R_+ and not identically equal to the 0 function on R for μ -a.e. $t \in T$, and such that $\Phi(\cdot, u)$ is a Σ -measurable function for all $u \in R$.

A Musielak—Orlicz function Φ such that $\Phi(t_1, u) = \Phi(t_2, u)$ for all $t_1, t_2 \in T$ and $u \in R$ is called an Orlicz function. For a given Musielak—Orlicz function Φ and a measure μ , the Musielak—Orlicz space $L^\Phi(\mu)$ is defined as the set of all equivalence classes of Σ -measurable functions x from T into R such that $I_\Phi(\lambda x) = \int_T \Phi(T, \lambda x) d\mu < +\infty$ for a certain $\lambda > 0$ depending on x . If Φ is an Orlicz function, then $L^\Phi(\mu)$ is called an Orlicz space (see [11], [12], [13] and [14]). We denote by $E^\Phi(\mu)$ the subspace of $L^\Phi(\mu)$ defined as the set of all $x \in L^\Phi(\mu)$ such that $I_\Phi(\lambda x) < +\infty$ for every $\lambda > 0$.

A Musielak—Orlicz space $L^\Phi(\mu)$ equipped with the Luxemburg norm

$$\|x\|_\Phi = \inf\{r > 0 : I_\Phi \frac{x}{r} \leq 1\}$$

is a Banach space (see [11], [12] and [13]). For any Musielak—Orlicz function Φ , the function Φ^* , complementary to Φ in the sense of Young, is defined by the formula

$$\Phi^*(t, u) = \sup_{v \geq 0} \{u|v - \Phi(t, v)\}$$

for all $t \in T$ and $u \in R$.

We say that a Musielak—Orlicz function Φ satisfies the Δ_2 -condition if there exist a constant $k \geq 2$, a null set $T_0 \in \Sigma$ and a Σ -measurable non-negative function h with $\int_T \Phi(t, h(t)) d\mu < +\infty$ such that $\Phi(t, 2\mu) \geq k\Phi(t, \mu)$ for any $t \in T \setminus T_0$ and $\mu \geq h(t)$ (see [9]).

Every Musielak—Orlicz function which satisfies the Δ_2 -condition has finite values.

A normed space $(X, \|\cdot\|)$ is said to be locally uniformly non- $l_n^{(1)}$ ($n \in N, n \geq 2$) if for every $x_1 \in X$ with $\|x_1\| = 1$ there exists $\delta(x_1)$ in the interval $(0, 1)$ such that for all norm-one elements x_2, \dots, x_n in X , the inequality $\|x_1 \pm \dots \pm x_n\| \leq n(1 - \delta(x_1))$ holds for a certain choice of signs ± 1 (see [16]).

A normed space $(X, \|\cdot\|)$ is called non- $l_n^{(1)}$ ($n \in N, n \geq 2$) if for any norm-one element x_1, \dots, x_n in X , we have $\|x_1 \pm \dots \pm x_n\| < n$ for a certain choice of signs ± 1 (see [3]).

Now, we shall give some lemmas which will be used in this paper.

Lemma 1. (see [4]) *The space l^∞ is not non- $l_n^{(1)}$.*

Lemma 2. (see [8]) *The space l^∞ contains an isometric copy of l^1 .*

Theorem 3. (see [3]) *A normed space $(X, \|\cdot\|)$ is non- $l_n^{(1)}$ if and only if it does not contain any isometric copy of $l_n^{(1)}$.*

2. Results.

For a Musielak—Orlicz function Φ that has only finite values, define

$$g(t) = \sup\{u \in R_+ : \Phi(t, \cdot) \text{ is linear in the interval } [0, u]\};$$

obviously, g is a Σ -measurable function, and $g(t) = +\infty$ whenever $\Phi(t, \cdot)$ is linear on the whole R_+ .

Theorem 4. *A Musielak—Orlicz space $L^\Phi(\mu)$ equipped with the Luxemburg norm is non- $l_n^{(1)}$ ($n \in N, n \geq 2$) if and only if:*

- a) Φ satisfies the Δ_2 -condition,
- b) $\int_T \Phi(t, g(t)) d\mu < n$.

PROOF: Sufficiency. Let $\|x_1\|_\Phi = \dots = \|x_n\|_\Phi = 1$. In virtue of the Δ_2 -condition, we get $I_\Phi(x_1) = \dots = I_\Phi(x_n) = 1$ (see [7]). Now, we shall prove that for all $u_1, \dots, u_n \in R$ and μ -a.e. $t \in T$, we have

$$(*) \quad \sum_{i=1}^n \Phi(t, u_i) > \Phi(t, g(t)) \text{ implies } \Phi\left(t, \frac{u_1 \pm \dots \pm u_n}{n}\right) < \frac{1}{n} \sum_{i=1}^n \Phi(t, u_i),$$

for a certain choice of signs ± 1 . For this purpose we shall consider two cases:

I. $\max |u_i| > g(t)$. For the choice of signs ± 1 such that

$$|u_1 \pm \dots \pm u_n| \leq \max_{1 \leq i \leq n} |u_i|,$$

we get

$$\begin{aligned} \Phi\left(t, \frac{u_1 \pm \dots \pm u_n}{n}\right) &\leq \Phi\left(t, \frac{\max |u_i|}{n}\right) < \left(\frac{1}{n} \Phi(t, \max |u_i|)\right) = \\ &= \frac{1}{n} \max \Phi(t, u_i) \leq \frac{1}{n} \sum_{i=1}^n \Phi(t, u_i). \end{aligned}$$

II. $\max |u_i| \leq g(t)$. Then at least two numbers among $\Phi(t, u_i)$ $i = 1, \dots, n$ are positive. In the opposite case, we have $\sum_{i=1}^n \Phi(t, u_i) = \Phi(t, u_k) \leq \Phi(t, g(t))$, where $1 \leq k \leq n$, which contradicts to the assumption in condition (*). Thus, we get

$$\max \Phi(t, u_i) / \sum_{i=1}^n \Phi(t, u_i) < 1.$$

For a certain choice of signs ± 1 , we have $|u_1 \pm \dots \pm u_n| \leq \max_{1 \leq i \leq n} |u_i|$. Therefore,

$$\begin{aligned} \Phi\left(t, \frac{u_1 \pm \dots \pm u_n}{n}\right) &\leq \Phi\left(t, \frac{\max |u_i|}{n}\right) = \frac{1}{n} \Phi(t, \max |u_i|) = \\ &= \frac{1}{n} \max \Phi(t, u_i) < \frac{1}{n} \sum_{i=1}^n \Phi(t, u_i). \end{aligned}$$

For this choice of signs ± 1 , combining the cases I and II, we get (*). Define

$$A = \left\{ t \in T : \sum_{i=1}^n \Phi(t, x_i(t)) > \Phi(t, g(t)) \right\}.$$

Then, in virtue of (*), we have

$$\Phi\left(t, \frac{x_1(t) \pm \dots \pm x_n(t)}{n}\right) < \frac{1}{n} \sum_{i=1}^n \Phi(t, x_i(t))$$

for all $t \in A$ and a certain choice of sign ± 1 . Therefore,

$$\sum_{\pm 1} \Phi\left(t, \frac{x_1(t) \pm \dots \pm x_n(t)}{n}\right) < \frac{2^{n-1}}{n} \sum_{i=1}^n \Phi(t, x_i(t)).$$

Integrating this inequality on both sides over A , we get

$$\sum_{\pm 1} I_{\Phi}\left(\frac{(x_1 \pm \dots \pm x_n)\chi_A}{n}\right) < \frac{2^{n-1}}{n} \sum_{i=1}^n I_{\Phi}(x_i \chi_A).$$

Hence, we obtain

$$\begin{aligned} 2^{n-1} - \sum_{\pm 1} I_{\Phi} \left(\frac{(x_1 \pm \cdots \pm x_n)}{n} \right) &= \frac{2^{n-1}}{n} \sum_{i=1}^n I_{\Phi}(x_i) - \sum_{\pm 1} I_{\Phi} \left(\frac{x_1 \pm \cdots \pm x_n}{n} \right) \geq \\ &\geq \frac{2^{n-1}}{n} \sum_{i=1}^n I_{\Phi}(x_i \chi_A) - \sum_{\pm 1} I_{\Phi} \left(\frac{(x_1 \pm \cdots \pm x_n) \chi_A}{n} \right). \end{aligned}$$

Hence, in virtue of the previous inequality, we get

$$\sum_{\pm 1} I_{\Phi} \left(\frac{x_1 \pm \cdots \pm x_n}{n} \right) < 2^{n-1}.$$

Then, for a certain choice of signs ± 1 , we have

$$I_{\Phi} \left(\frac{x_1 \pm \cdots \pm x_n}{n} \right) < 1.$$

Thus, in virtue of the Δ_2 -condition, it follows that

$$\left\| \frac{x_1 \pm \cdots \pm x_n}{n} \right\|_{\Phi} < 1$$

for a certain choice of signs ± 1 . The proof of sufficiency is finished.

Necessity. If Φ does not satisfy the Δ_2 -condition, then $L^{\Phi}(\mu)$ contains an isometric copy of l^{∞} (see [5], [6]), so $L^{\Phi}(\mu)$ is not non- $l_n^{(1)}$ (see Lemma 1).

Now, assume that Φ satisfies the Δ_2 -condition and the condition (b) does not hold, i.e. $\int_T \Phi(t, g(t)) d\mu \geq n$. In virtue of the Δ_2 -condition, $\Phi(t, \cdot)$ is continuous for μ -a.e. $t \in T$. If $g(t) < +\infty$ for μ -a.e. $t \in T$, then there are pairwise disjoint sets $A_1, A_2, \dots, A_n \in \Sigma$ such that

$$\int_{A_1} \Phi(t, g(t)) d\mu = \cdots = \int_{A_n} \Phi(t, g(t)) d\mu = 1$$

Define $x_i = g \chi_{A_i}$ for $i = 1, 2, \dots, n$. We have $I_{\Phi}(x_i) = 1$, and

$$I_{\Phi} \left(\frac{x_1 \pm \cdots \pm x_n}{n} \right) = \sum_{i=1}^n I_{\Phi} \left(\frac{x_i}{n} \right) = \frac{1}{n} \sum_{i=1}^n I_{\Phi}(x_i) = 1$$

for any choice of signs ± 1 . Thus, we have

$$\left\| \frac{x_1 \pm \cdots \pm x_n}{n} \right\|_{\Phi} = 1$$

for any choice of signs ± 1 . It means that $L^{\Phi}(\mu)$ is not non- $l_n^{(1)}$.

If $g(t) = +\infty$ for $t \in A$, where $A \in \Sigma$ and $\mu(A) > 0$, then $\Phi(t, u) = P(t)|u|$ for every $t \in A$ and $u \in \mathbb{R}_+$, where P is a Σ -measurable function positive on A . Define on $\Sigma \cap A$ a new non-atomic measure ν by

$$\nu(B) = \int_B P(t) d\mu \quad (\forall B \in \Sigma \cap A).$$

Then $L^{\Phi}(\mu, A) = L^1(\nu, A)$, and therefore $L^{\Phi}(\mu)$ is not non- $l_n^{(1)}$ (see [4]). The proof is finished. ■

Now, we shall give a criterion in order that a Musielak—Orlicz space $L^{\Phi}(\mu)$ contains an isometric copy of l^1 .

Theorem 5. *A Musielak—Orlicz space $L^\Phi(\mu)$ equipped with the Luxemburg norm contains an isometric copy of l^1 if and only if:*

- c) Φ does not satisfy the Δ_2 -condition, or
- d) $I_\Phi(g) = +\infty$, where g is the function defined before Theorem 4.

PROOF : Sufficiency. If Φ does not satisfy the Δ_2 -condition, then $L^\Phi(\mu)$ contains an isometric copy of l^∞ (see [5], [6]) and, in view of Lemma 2, it contains an isometric copy of l^1 . Now, assume that Φ satisfies condition (d) and $(g(t) < +\infty$ for μ -a.e. $t \in T$. We can assume that Φ satisfies the Δ_2 -condition. The measure ν_μ defined on Σ by the formula

$$\nu_\mu(A) = I_\Phi(g\chi_A)$$

is non-atomic and infinite.

Therefore, there exists a sequence $(A_k)_{k=1}^\infty$ of pairwise disjoint sets in Σ such that $I_\Phi(g\chi_{A_k}) = 1$ for every $k \in N$. Denote $a_k = g\chi_{A_k}$ and define an operator P from l^1 into $L^\Phi(\mu)$ by

$$Py = \sum_{k=1}^\infty c_k a_k \quad (\forall y = (c_k) \in l^1).$$

P is linear and it is easily seen that $Py \in E^\Phi(\mu)$ for any $y \in l^1$. In fact, taking into account that $\Phi(t, \cdot)$ is linear on the interval $[0, g(t)]$, we get $\Phi(t, \alpha g(t)) = |\alpha| \Phi(t, g(t))$ for every $|\alpha| \leq 1$. Given $\lambda > 0$, choose $n_0 \in N$ in such a manner that $\lambda|c_k| \leq 1$ for $n \geq n_0$. We have

$$\begin{aligned} I_\Phi(\lambda Py) &= \sum_{k=1}^{n_0-1} \int_{A_k} \Phi(t, \lambda c_k a_k(t)) d\mu + \sum_{k=n_0}^\infty \int_{A_k} \Phi(t, \lambda c_k a_k(t)) d\mu \\ &= \sum_{k=1}^{n_0-1} \int_{A_k} \Phi(t, \lambda c_k a_k(t)) d\mu + \sum_{k=n_0}^\infty \lambda |c_k| \int_{A_k} \Phi(t, a_k(t)) d\mu \\ &= \sum_{k=1}^{n_0-1} \int_{A_k} \Phi(t, \lambda c_k a_k(t)) d\mu + \lambda \sum_{k=n_0}^\infty |c_k| < +\infty. \end{aligned}$$

Now, we shall prove that P is an isometry. We have

$$\begin{aligned} I_\Phi \left(\frac{Py}{\|y\|_{l^1}} \right) &= \int_T \Phi \left(t, \frac{Py}{\|y\|_{l^1}} \right) d\mu = \sum_{k=1}^\infty \int_{A_k} \Phi \left(t, \frac{c_k a_k(t)}{\|y\|_{l^1}} \right) d\mu \\ &= \sum_{k=1}^\infty \frac{|c_k|}{\|y\|_{l^1}} \int_{A_k} \Phi(t, a_k) d\mu = \sum_{k=1}^\infty \frac{|c_k|}{\|y\|_{l^1}} = 1 \end{aligned}$$

Hence,

$$\left\| \frac{Py}{\|y\|_{l^1}} \right\|_\Phi = 1, \text{ i.e. } \|Py\|_\Phi = \|y\|_{l^1}.$$

Assume now that $g(t) = +\infty$ for $t \in A$, where $A \in \Sigma$ and $\mu(A) > 0$. Then $L^\Phi(\mu, A) = L^1(\nu, A)$, where ν is defined as in the proof of Theorem 4. Since $L^1(\nu, A)$ contains an isometric copy of l^1 (see [8]), $L^\Phi(\mu, A)$ contains an isometric copy of l^1 .

Necessity. Assume that none of the conditions (c) and (d) is satisfied. This means that Φ satisfies the Δ_2 -condition and $\int_T \Phi(t, g(t)) d\mu < +\infty$. Therefore, there is $k \in N, k \geq 2$, such that $\int_T \Phi(t, g(t)) d\mu \leq k$. In view of Theorem 4, $L^\Phi(\mu)$ is non- $l_n^{(1)}$ for all $n > k, n \in N$. In virtue of Theorem 3, $L^\Phi(\mu)$ contains no isometric copy of l^1 . The proof is finished. ■

Theorem 6. *Let Φ be a Musielak—Orlicz function such that $\Phi(t, \cdot)$ is linear in no neighbourhood of 0 in R_+ for μ -a.e. $t \in T$. Then the Musielak—Orlicz space $L^\Phi(\mu)$ equipped with the Luxemburg norm is locally uniformly non- $l_n^{(1)}$ if and only if Φ satisfies the Δ_2 -condition.*

PROOF : Sufficiency. Let $\|x_1\|_\Phi = \dots = \|x_n\|_\Phi = 1$. Then, in virtue of the Δ_2 -condition, we have $I_\Phi(x_1) = \dots = I_\Phi(x_n) = 1$ (see [7]). Let $c > 0$ be such that the set

$$A_1 = \{t \in T : c^{-1} \leq \Phi(t, x_1(t)) \leq c\}$$

satisfies the condition $I_\Phi(x_1 \lambda_{A_1}) \geq \frac{7}{8}$. Let $m > 0$ be such that $\frac{c}{m} \leq \frac{1}{8(n-1)}$, and define

$$A_i = \{t \in T : \Phi(t, x_i(t)) \leq m\} \quad \text{for } i = 2, \dots, n$$

we have

$$m\mu(T \setminus A_i) < I_\Phi(x_i \lambda_{T \setminus A_i}) \leq 1.$$

Thus,

$$\mu(T \setminus A_i) \leq \frac{1}{m} \quad \text{for } i = 2, \dots, n.$$

Hence, we get

$$I_\Phi(x_1 \lambda_{A_1 \setminus A_i}) \leq c\mu(A_1 \setminus A_i) \leq \frac{c}{m} \leq \frac{1}{8(n-1)}.$$

Denoting $D = \bigcap_{i=2}^n A_i$, we have

$$\begin{aligned} \frac{7}{8} &\leq I_\Phi(x_1 \lambda_{A_1}) = I_\Phi(x_1 \lambda_{A_1 \setminus D}) + I_\Phi(x_1 \lambda_D) \\ &= I_\Phi(x_1 \lambda_{\bigcup_{i=2}^n (A_1 \setminus A_i)}) + I_\Phi(x_1 \lambda_D) \\ &\leq \frac{1}{8(n-1)}(n-1) + I_\Phi(x_1 \lambda_D), \end{aligned}$$

whence $I_\Phi(x_1 \lambda_D) \geq \frac{3}{4}$. Define

$$P(t) = \sup \left\{ \frac{n\Phi(t, \frac{u}{n})}{\Phi(t, u)} : \Phi(t, u) \in [c^{-1}, m] \right\}.$$

In virtue of the assumption that $\Phi(t, \cdot)$ is linear in no neighbourhood of 0 in R_+ , we get $0 < P(t) < 1$ for μ -a.e. $t \in D$. Hence, we have $\Phi(t, \frac{x}{n}) \leq \frac{P(t)}{n} \Phi(t, x)$ for μ -a.e. $t \in D$, and all x satisfying $\Phi(t, x) \in [c^{-1}, m]$. Define

$$B_k = \left\{ t \in D : P(t) \leq 1 - \frac{1}{k} \right\}.$$

By Σ -measurability of P , it follows that $B_k \in \Sigma$ for $k = 1, 2, \dots$. There is $l \in N$ such that $I_\Phi(x_l \chi_{B_l}) \geq \frac{1}{2}$. Denote $\sigma = 1 - \frac{1}{l}$ and $B = B_l$. Now, we shall prove that for every $t \in B$, we have

$$(**) \quad \sum_{\pm 1} \Phi \left(t, \frac{x_1(t) \pm \dots \pm x_n(t)}{n} \right) \leq \frac{2^{n-1} - 1 + \sigma}{n} \sum_{i=1}^n \Phi(t, x_i(t)).$$

For at least one choice of signs ± 1 , such that $|x_1(t) \pm \dots \pm x_n(t)| \leq \max_{1 \leq i \leq n} |x_i(t)|$, we have

$$(1) \quad \Phi \left(t, \frac{x_1(t) \pm \dots \pm x_n(t)}{n} \right) \leq \Phi \left(\frac{\max |x_i(t)|}{n} \right) \leq \frac{\sigma}{n} \Phi(t, \max |x_i(t)|) = \frac{\sigma}{n} \max \Phi(t, x_i(t)) \leq \frac{\sigma}{n} \sum_{i=1}^n \Phi(t, x_i(t)),$$

for every $t \in B$. For the remaining $2^{n-1} - 1$ choice of signs ± 1 , by the convexity of Φ , we have

$$(2) \quad \Phi \left(t, \frac{x_1(t) \pm \dots \pm x_n(t)}{n} \right) \leq \frac{1}{n} \sum_{i=1}^n \Phi(t, x_i(t)), \text{ for every } t \in B.$$

Combining (1) and (2), we get (**). Integrating the inequality (**) both-sides over B , we get

$$\sum_{\pm 1} I_\Phi \left(\frac{(x_1 \pm \dots \pm x_n) \chi_B}{n} \right) \leq \frac{2^{n-1} - 1 + \sigma}{n} \sum_{i=1}^n I_\Phi(x_i \chi_B).$$

Hence, we obtain

$$\begin{aligned} & 2^{n-1} - \sum_{\pm 1} I_\Phi \left(\frac{x_1 \pm \dots \pm x_n}{n} \right) \\ &= \frac{2^{n-1}}{n} \sum_{i=1}^n I_\Phi(x_i) - \sum_{\pm 1} I_\Phi \left(\frac{x_1 \pm \dots \pm x_n}{n} \right) \\ &\geq \frac{2^{n-1}}{n} \sum_{i=1}^n I_\Phi(x_i \chi_B) - \sum_{\pm 1} I_\Phi \left(\frac{(x_1 \pm \dots \pm x_n) \chi_B}{n} \right) \\ &\geq \frac{2^{n-1}}{n} \sum_{i=1}^n I_\Phi(x_i \chi_B) - \frac{2^{n-1} - 1 + \sigma}{n} \sum_{i=1}^n I_\Phi(x_i \chi_B) \\ &= \frac{1 - \sigma}{n} \sum_{i=1}^n I_\Phi(x_i \chi_B) \geq \frac{1 - \sigma}{n} I_\Phi(x_l \chi_B) \\ &\geq \frac{1 - \sigma}{2n} = \eta. \end{aligned}$$

Thus, we have

$$\sum_{\pm 1} I_{\Phi} \left(\frac{(x_1 \pm \cdots \pm x_n)}{n} \right) \leq 2^{n-1} - \eta = 2^{n-1}(1 - q),$$

where $q = \eta/2^{n-1}$ and it depends only on x_1 . Therefore, for a certain choice of signs ± 1 , we get

$$I_{\Phi} \left(\frac{x_1 \pm \cdots \pm x_n}{n} \right) \leq 1 - q.$$

In virtue of the Δ_2 -condition, we have

$$\left\| \frac{x_1 \pm \cdots \pm x_n}{n} \right\|_{\Phi} \leq 1 - \beta(q)$$

for a certain choice of signs ± 1 , where β is a function from $(0, 1)$ into $(0, 1)$ such that $\|x\| \leq 1 - \beta(q)$, whenever $I_{\Phi}(x) \leq 1 - q$ (see [1]).

Necessity. If Φ does not satisfy the Δ_2 -condition, then $L^{\Phi}(\mu)$ contains an isometric copy of l^{∞} (see [5], [6]). Therefore, in view of Lemma 1, $L^{\Phi}(\mu)$ is not locally uniformly non- $l_n^{(1)}$. The proof of the Theorem 6 is finished. ■

REFERENCES

- [1] Bombal F., *On l_1 subspaces of Orlicz vector-valued function spaces*, Math. Proc. Comb. Phil. Soc. **101**, 107 (1987), 107–112.
- [2] Fuentes F. and Hernandez F.L., *On weighted Orlicz sequence spaces and their subspaces*, Rocky Mount. Math. J. **18**, 3 (1988), 585–599.
- [3] Grzaslewicz R., Hudzik H. and Orlicz W., *Uniformly non- $l_n^{(1)}$ property in some normed spaces*, Bull. Acad. Polon. Sci. Math. **34**, 3-4 (1986), 161–171.
- [4] Hudzik H., *Locally uniformly non- $l_n^{(1)}$ Orlicz spaces*, Proceed. of the 13th Winter School on Abstract Analysis, Srni, January 20-27, 1985, Supplemento ai Rendiconti del Circolo Matematico di Palermo Ser. II, num. 10 (1985), 49–56.
- [5] Hudzik H., *Uniform convexity of Musielak—Orlicz spaces with Luxemburg's norm*, Commentationes Math. **23** (1983), 21–32.
- [6] Hudzik H., *On some equivalent conditions in Musielak—Orlicz spaces*, Commentationes Math. **24** (1984), 57–64.
- [7] Hudzik H., *Strict convexity of Musielak—Orlicz spaces with Luxemburg's norm*, Bull. Acad. Polon. Sci. Math. **29**, 5-6 (1981), 235–247.
- [8] Hudzik H., *Orlicz spaces containing a copy of L^1* , Math. Japonica.
- [9] Hudzik H. and Kaminska A., *On uniformly convexifiable and B-convex Musielak—Orlicz spaces*, Commentationes Math. **25** (1985), 59–75.
- [10] Hudzik H., Kaminska A., Kurc W., *Uniformly non- $l_n^{(1)}$ Musielak—Orlicz spaces*, Bull. Acad. Polon. Sci. Math. **35**, 7-8 (1987), 441–448.
- [11] Krasnoselski M.A. and Ruticki Ia.B., *Convex functions and Orlicz spaces*, Groningen 1961 (translation).
- [12] Luxemburg W.A.J., *Banach function spaces*, Thesis, Delft 1955.
- [13] Musielak J., *Orlicz spaces and modular spaces*, Lecture Notes in Math., Springer-Verlag, 1034 (1983).

- [14] Musielak J., Orlicz W., *On modular spaces*, Studia Math. **18** (1959), 49–65.
- [15] Milnes H.W., *Convexity of Orlicz spaces*, Pacific J. Math. (1957), 1451–1486.
- [16] Schaffer J.J., *Geometry of spheres in normed spaces*, Lecture Notes in Math., Springer-Verlag, **20** (1976) .

University of Aleppo, Faculty of Sciences, Department of Mathematics, Aleppo, Syria
Institute of Mathematics, A. Mickiewicz University, 60-769 Poznań, Poland

(Received November 9, 1989)