

Alexander Abian

Compactification of the domains of certain rings of functions

*Archivum Mathematicum*, Vol. 22 (1986), No. 2, 61--64

Persistent URL: <http://dml.cz/dmlcz/107247>

## Terms of use:

© Masaryk University, 1986

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## COMPACTIFICATION OF THE DOMAINS OF CERTAIN RINGS OF FUNCTIONS\*)

ALEXANDER ABIAN  
(Received December 2, 1983)

**Abstract.** Let  $R$  be a ring (with unit  $u$ ) of functions (pointwise addition and multiplication) from a set  $X$  into a ring  $Y$  with the special property that every proper ideal of  $R$  is contained in a proper ideal  $J$  of  $R$  for which  $R/J$  is isomorphic to a subring of  $Y$ . We show that  $X$  can be extended to a set  $X'$  and every element  $f$  of  $R$  can be extended to a function  $f'$  from  $X'$  into  $Y$  such that the resulting set  $R' = \{f' \mid f \in R\}$  is a ring of functions from  $X'$  into  $Y$  and the correspondence  $f \rightarrow f'$  is a ring isomorphism from  $R$  onto  $R'$ . Moreover, if the image (under the isomorphism)  $u'$  of the unit  $u$  is covered by the elements of a subset  $E'$  of  $R'$  (i.e., for every  $z \in X'$  if  $u'(z) \neq 0$  then  $g'(z) \neq 0$  for some  $g' \in E'$ ) then  $u'$  is already covered by finitely many elements of  $E'$ .

**Key words.** Ideal of a Ring. Compact domain.  
**MS Classification.** Primary 54 D 35.

In what follows all the ring-theoretic statements which are made in connection with a set of functions (with a common domain) refer to the pointwise addition and multiplication of the elements of that set.

Also, in what follows, we let  $R$  stand for a ring of functions from a set  $X$  (i.e.,  $X$  is the common domain of the elements of  $R$ ) into a ring  $Y$  such that  $R$  has a unit  $u$  and  $R$  satisfies that following condition:

*For every proper ideal  $I$  of  $R$  there exists a proper ideal  $J$  of  $R$  such that*

$$(1) \quad I \subseteq J \text{ and } R/J \text{ is isomorphic to a subring of } Y.$$

Next, we introduce the notion of *covering* of an element of a ring of function with a subset of that ring:

**Definition.** Let  $E'$  be a set of functions from a set  $X'$  into a ring  $Y$  (with zero 0). Then a function  $f'$  from  $X'$  into  $Y$  is said to be covered by the elements of  $E'$  if and only if for every  $z \in X'$

$$(2) \quad f'(z) \neq 0 \quad \text{implies} \quad g'(z) \neq 0 \quad \text{for some} \quad g' \in E'$$

i.e.,  $f'$  is covered by the elements of  $E'$  if and only if whenever  $f'$  does not vanish at a point, so does an element of  $E'$  at that point.

\*) This work partly supported by the Iowa State University SHRI.

**Lemma.** *Let  $R'$  be a ring of functions from a set  $X'$  into a ring  $Y$  and let  $E'$  be a subset of  $R'$ . If an element  $f'$  of  $R'$  is covered by no finite number of elements of  $E'$  then  $f'$  is not an element of the ideal  $I'$  (of  $R'$ ) generated by  $E'$ .*

**Proof.** Since  $f'$  is covered by no finite number of elements of  $E'$ , in view of (2), we see that for every finite number of elements  $e'_0, \dots, e'_n$  of  $E'$  there exists  $z \in X'$  such that  $e'_i(z) = 0$  for every  $i \in n$  whereas  $f'(z) \neq 0$ . Thus,  $f' \notin I'$  as desired.

Based on the above we prove:

**Theorem.** *Let  $R$  be a ring of functions from a set  $X$  into a ring  $Y$  such that  $R$  has a unit  $u$  and  $R$  satisfies (1). Then  $X$  can be extended to a set  $X'$  and every element  $f$  of  $R$  can be extended to a function  $f'$  from  $X'$  into  $Y$  such that:*

- (i) *the resulting set  $R' = \{f' \mid f \in R\}$  is a ring of functions  $f'$  from  $X'$  into  $Y$ .*
- (ii) *the correspondence  $f \rightarrow f'$  is a ring isomorphism from  $R$  onto  $R'$ .*
- (iii) *if the image (under the above isomorphism)  $u'$  of the unit  $u$  of  $R$  is covered by the elements of a subset  $E'$  of  $R'$  then  $u'$  is already covered by finitely many elements of  $E'$ .*

**Proof.** Clearly, the set

$$(3) \quad J_x = \{f \mid (f \in R \quad \text{and} \quad f(x) = 0)\} \quad \text{for every} \quad x \in X$$

is an ideal of  $R$  such that  $R/J_x$  is isomorphic to a subring of  $Y$ .

Let

$$(4) \quad \{J_v \mid v \in V\}$$

be the set of all the ideals  $J_v$  of  $R$  such that  $R/J_v$  is isomorphic to a subring of  $Y$  and such that  $J_v \notin \{J_x \mid x \in X\}$  where  $J_x$  is given by (3).

With every  $J_v$  which appears in (4) we associate a unique isomorphism  $i_v$  from  $R/J_v$  onto a subring of  $Y$ . Thus, for every  $f \in R$  and every  $v \in V$ , we have:

$$(5) \quad i_v([f]) \quad \text{is a unique element of } Y$$

where  $[f]$  is the coset of  $R/J_v$  of which  $f$  is an element.

Now, let us consider the set  $X'$  given by:

$$(6) \quad X' = X \cup V$$

where  $V$  is as in (4).

Clearly,  $X'$  is an extension of  $X$ . Also, by (3), (4), (6) we see that

$$(7) \quad \{J_z \mid z \in X'\}$$

is the set of *all* the ideals  $J$  of  $R$  such that  $R/J$  is isomorphic to a subring of  $Y$ .

In view of (5), to every  $f \in R$  we correspond a function  $f'$  from  $X'$  into  $Y$  defined as:

$$(8) \quad f' = f \quad \text{on } X \quad \text{and} \quad f'(v) = i_v([f]) \quad \text{for every } v \in V$$

Clearly, from (4), (6), (8) it follows that  $f'$  is an extension of  $f$  to  $X'$ .  
Again, from (4) and (8) we see that

$$J_v = \{f \mid (f \in R) \text{ and } f'(v) = 0\} \quad \text{for every } v \in V$$

which by (3) and (7) implies that

$$(9) \quad J_z = \{f \mid (f \in R) \text{ and } f'(z) = 0\} \quad \text{for every } z \in X'$$

From (8) it follows that the correspondence  $f \rightarrow f'$  is one-to-one and that

$$(10) \quad f' + g' = (f + g)' \quad \text{and} \quad f' \cdot g' = (f \cdot g)'$$

for every element  $f$  and  $g$  of  $R$ .

Thus, from (10) we see that the set  $R'$  (of functions from  $X'$  into the ring  $Y$ ) given by:

$$(11) \quad R' = \{f' \mid f \in R\}$$

is an isomorphic image of the ring  $R$  (of functions  $f$  from  $X$  into the ring  $Y$ ).

Clearly, (10) and (11) establish (i) and (ii).

Next, let  $u' \in R'$  be the image (under the above isomorphism) of the unit element  $u$  of  $R$ . Obviously,  $u'$  is the unit of  $R'$ . Let  $u'$  be covered by the elements of a subset  $E'$  of  $R'$  (see the Definition above). To prove (iii) we must show that  $u'$  is already covered by some finitely many elements of  $E'$ . Let us assume to the contrary that  $u'$  is covered by no finite number of elements of  $E'$ . Thus, by the Lemma, the ideal  $I'$  (of  $R'$ ) generated by  $E'$  is such that

$$(12) \quad u' \notin I'$$

Obviously,

$$(13) \quad E' \subseteq I'$$

Since  $f \rightarrow f'$  is a ring isomorphism, we see that the subset  $I$  of  $R$  given by:

$$(14) \quad I = \{g \mid g' \in I'\}$$

is an ideal of  $R$  and by (12) we have  $u \notin I$ . Thus,  $I$  is a proper ideal of  $R$  and therefore there exists a proper ideal  $J$  of  $R$  satisfying (1). However, by (7) every such  $J$  ideal of  $R$  must be equal to an ideal  $J_z$  (of  $R$ ) for some  $z \in X'$ . Thus,

$$I \subseteq J_z \quad \text{and} \quad u \notin J_z \quad \text{for some } z \in X'$$

which by (9) and (14) implies

$$u'(z) \neq 0 \quad \text{and} \quad g'(z) = 0 \quad \text{for every } g' \in I'$$

which in turn, in view of (13) and (2) contradicts that  $u'$  is covered by the elements of  $E'$ . Hence our assumption is false and (iii) is also established.

**Remark 1.** Let the ring  $R$  instead of satisfying condition (1) satisfies the stronger condition (1\*) given by:

(1\*) *For every element  $f$  of  $R$  and every ideal  $I$  of  $R$  if  $f \notin I$  then there exists an ideal  $J$  of  $R$  such that  $I \subseteq J$  and  $f \notin J$  and such that  $R/J$  is isomorphic to a subring of  $Y$*

i.e., and ideal of  $R$  which misses an element  $f$  of  $R$  is contained in an ideal  $J$  of  $R$  which also misses  $f$  and for which  $R/J$  is isomorphic to a subring of  $Y$ .

Then the proof of our Theorem shows (via replacing in it  $u$  by an element  $f$  of  $R$ ) that (iii) of the conclusion of our Theorem (even with dropping the hypothesis that  $R$  must have a unit) will read:

(iii\*) *if an element  $f'$  (not necessarily the unit  $u'$ ) of  $R'$  is covered by the elements of a subset  $E'$  of  $R'$  then  $f'$  is already covered by finitely many elements of  $E'$ .*

**Remark 2.** In Topology, very often, the (common) domain  $X$  of a ring of functions (from  $X$ ) into a ring  $Y$  is topologized by an open basis whose members consist of those subsets  $B$  of  $X$  for which there exists an element  $f$  of  $R$  such that  $B$  is the set of all those points of  $X$  on each of which  $f$  does not vanish. Let us call such a topology the *Characteristic topology of  $R$  on  $X$* . But then our Theorem can be obviously rephrased as follows:

**Corollary.** *Let  $R$  be a ring of functions from a set  $X$  into a ring  $Y$  such that  $R$  has a unit and  $R$  satisfies (1). Then  $X$  can be extended to a set  $X'$  and every element  $f$  of  $R$  can be extended to a function  $f'$  from  $X'$  into  $Y$  such that the resulting set  $R' = \{f' \mid f \in R\}$  is a ring of functions  $f'$  from  $X'$  into  $Y$  and the correspondence  $f \rightarrow f'$  is a ring isomorphism from  $R$  onto  $R'$ . Moreover, the Characteristic topology of  $R'$  on  $X'$  (whose restriction to  $X$  is the Characteristic topology of  $R$  on  $X$ ) is compact.*

The Corollary shows how  $X$  endowed with a Characteristic topology can be compactified by extending it to  $X'$  which is endowed by a corresponding Characteristic topology.

*A. Abian  
Mathematical Institute  
University of Oxford  
Oxford, England*

*and  
Department of Mathematics  
Iowa State University  
Ames, Iowa 50011  
U.S.A.*