

Bedřich Pondělíček

Note on quasi-Hamiltonian semigroups

Časopis pro pěstování matematiky, Vol. 110 (1985), No. 4, 356–358

Persistent URL: <http://dml.cz/dmlcz/118250>

Terms of use:

© Institute of Mathematics AS CR, 1985

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

NOTE ON QUASI HAMILTONIAN SEMIGROUPS

BEDŘICH PONDĚLÍČEK, Praha

(Received February 15, 1984)

Following A. Cherubini and A. Varisco [1] a semigroup is said to be *quasi hamiltonian* if all its subsemigroups are permutable. In this note we shall show that every variety of quasi hamiltonian semigroups is commutative.

Let a be an element of a semigroup S . By $[a]$ we denote the subsemigroup of S generated by a . It is easy to show (see Lemma 3 of [2]) that a semigroup S is quasi hamiltonian if and only if we have

$$(1) \quad ab \in [b][a] \text{ for every } a, b \in S.$$

A semigroup S is called *quasicommutative* [3] if we have

$$(2) \quad ab \in [b]a \text{ for every } a, b \in S.$$

A semigroup S is σ -*reflexive* if for every $a, b \in S$ and every subsemigroup H of S , $ba \in H$ implies $ab \in H$. It is easy to prove that a semigroup S is σ -reflexive if and only if we have

$$(3) \quad ab \in [ba] \text{ for every } a, b \in S.$$

In [4] it has been shown that the class of all quasicommutative semigroups coincides with the class of all σ -reflexive semigroups. This together with (2) and (3) implies that a semigroup S is quasicommutative if and only if we have

$$(4) \quad ab \in b[a] \text{ for every } a, b \in S.$$

Theorem 1. *Let S be a noncommutative semigroup such that $S \times S$ is a quasi hamiltonian semigroup. Then S is a periodic semigroup.*

Proof. Let $S \times S$ be a quasi hamiltonian semigroup. Suppose that $ab \neq ba$ for some $a, b \in S$. By way of contradiction, assume that there exists a non periodic element c of S . According to (1), we have $(a, c)(b, c) = (b, c)^m(a, c)^n$ for some positive integers m, n . Then we obtain $m = 1 = n$. This implies that $ab = ba$, which is a contradiction. Hence S is a periodic semigroup.

Lemma. *Let S be a quasi hamiltonian semigroup, $a, b, e \in S$ and $a^k = e = e^2$ for a positive integer k . If in $S \times S$ we have $(a, a)(b, e) = (b, e)^m(a, a)^n$ for some positive integers m, n , then $(a, a)(b, e) = (b, e)^m(a, a)$.*

Proof. Suppose that $(a, a)(b, e) = (b, e)^m(a, a)^n$. We have

$$(5) \quad ab = b^m a^n \quad \text{and} \quad ae = ea^n.$$

We can suppose that $n > 1$. Since S is quasi hamiltonian, by (1) we have $ba = a^i b^j$ for some positive integers i, j and so $ba \in S^1 ab S^1$, where S^1 denotes the semigroup S with an identity adjoined. According to (1), we obtain

$$(6) \quad ba \in S^1 ab.$$

By induction we shall prove the following proposition:

$$(7) \quad ab \in S^1 ba(a^{n-1})^r \quad \text{for all positive integers } r.$$

Evidently (7) is true for $r = 1$ by (5). Suppose that (7) is true for some r . Then by (6) we have $ab \in S^1 ab(a^{n-1})^r$ and so by (5) we obtain $ab \in S^1 b^m a^n (a^{n-1})^r \subseteq S^1 ba(a^{n-1})^{r+1}$.

It is clear that $r(n-1) > k$ for a positive integer r . Then, by (7), we have $ab \in Se$ and so, by (6), we have $ba \in S^1 ab \subseteq Se$. Therefore $ba = bae$ and using (5) we get $ab = abe = b^m a^n e = b^m e a^n = b^m a e = b^m a$. Hence we have $(a, a)(b, e) = (b, e)^m(a, a)$.

Theorem 2. *Let S be a semigroup such that $S \times S$ is a quasi hamiltonian semigroup. Then S is a quasicommutative semigroup.*

Proof. Suppose that $S \times S$ is a quasi hamiltonian semigroup. It is easy to show that S is a quasi hamiltonian semigroup. We can assume that S is non commutative. It follows from Theorem 1 that S is periodic. Let $a, b \in S$. Then there exists a positive integer k such that $a^k = e = e^2$. According to (1), we have $(a, a)(b, e) = (b, e)^m \cdot (a, a)^n$ for some positive integers m, n . Using Lemma we get $(a, a)(b, e) = (b, e)^m(a, a)$ and so $ab = b^m a$. Then, by (2), S is quasicommutative.

Theorem 3. *Let S be a semigroup such that $S \times S$ is a quasicommutative semigroup. Then S is a commutative semigroup.*

Proof. Suppose that $S \times S$ is a quasicommutative semigroup. Evidently, S is quasicommutative. By way of contradiction, assume that S is not commutative. Then there exist elements a, b of S such that $ab \neq ba$. Theorem 1 implies that S is periodic. Thus we have $a^k = e = e^2$ for a positive integer k . It follows from (4) that $(a, a)(b, e) = (b, e)(a, e)^n$ for a positive integer n . Using Lemma we get $(a, a) \cdot (b, e) = (b, e)(a, a)$ and so $ab = ba$, which is a contradiction. Hence S is commutative.

Corollary 1. *Let S be a semigroup such that $S \times S \times S \times S$ is a quas hamiltonian semigroup. Then S is a commutative semigroup.*

Corollary 2. *The variety of all commutative semigroups is the largest variety of quasi hamiltonian semigroups.*

Corollary 3. *Let m, n be positive integers. Then every semigroup satisfying the identity $xy = y^n x^m$ is commutative.*

References

- [1] *A. Cherubini, A. Varisco:* Quasi hamiltonian semigroups. Czech. Math. J. 33 (1983), 131—140.
- [2] *B. Pondělíček:* Uniform semigroups whose proper quasi-ideals are power joined. Semigroup Forum 22 (1981), 331—337.
- [3] *N. P. Mukherjee:* Quasicommutative semigroup I. Czech. Math. J. 22 (1972), 449—453.
- [4] *A. Cherubini, A. Varisco:* Quasi commutative semigroups and σ -reflexive semigroups. Semigroup Forum 19 (1980), 313—321.

Author's address: 166 27 Praha 6, Suchbátarova 2 (Katedra matematiky FEL ČVUT).