

Jan Čerych

A note on pervasive algebras

Časopis pro pěstování matematiky, Vol. 110 (1985), No. 4, 375--377

Persistent URL: <http://dml.cz/dmlcz/118253>

Terms of use:

© Institute of Mathematics AS CR, 1985

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A NOTE ON PERVASIVE ALGEBRAS

JAN ČERYCH, Praha

(Received February 13, 1984)

By a function algebra (on a compact Hausdorff space X) we mean a closed subalgebra, separating the points of X , of the sup-norm algebra $C(X)$ of all continuous complex-valued functions on X .

A function algebra A is said to be *pervasive* provided it satisfies the following condition:

Whenever F is a nonvoid proper closed subset of X , then $A|_F$, the algebra of all restrictions of the functions in A to the set F , is dense in $C(F)$ (naturally with respect to $|\cdot|_F$, the sup-norm on F).

The notion "pervasiveness" is due to Hoffman and Singer [1] who also were the first to investigate the properties of such algebras.

$C(X)$ is of course a pervasive algebra. More interesting are its proper pervasive subalgebras; the simplest of them is the classical disc algebra, the set of all uniform limits of polynomials on the unit circle in the z -plane, and related algebras.

Pervasiveness is a rather strong property, and it is interesting to seek for a non-trivial additional property which guarantees the pervasive algebra to be equal to the whole $C(X)$. In this sense we have investigated pervasive algebras in [2]. Our aim here is to strengthen the following Theorem A proved therein:

Theorem A. *Let A be a function algebra on X . Suppose that for any closed nonvoid proper subset F of X and for any function f in $C(F)$ there exists a positive constant $k(F, f)$ with the following property:*

Whenever e is a positive number, then there exists a g in A satisfying

$$|f - g|_F \leq e, \quad |g| \leq k(F, f).$$

Then A is equal to $C(X)$.

Remark that the assumption of Theorem A comprises the pervasiveness of the algebra A .

In this note we shall require the pervasiveness of A , and the bounded approximation by functions in A solely of a single function on a certain set, and come to the same conclusion. More specifically, we shall prove the following.

Theorem B. *Let A be a pervasive algebra on X . Let F and H be a disjoint couple of closed subsets of X which both have nonvoid interiors. Suppose that there is a constant c with the following property:*

Whenever e is positive, then there is an f in A satisfying

$$(1) \quad |f|_F < e, \quad |f - 1|_H < e, \quad |f| \leq c.$$

Then A is equal to $C(X)$.

Proof. Fix an arbitrary g in $C(X)$ and e positive. To prove Theorem B, it suffices to find an h in A satisfying

$$(2) \quad |g - h| < e.$$

It is obvious that $\overline{X - F}$ and $\overline{X - H}$ (where the bar denotes the closure in X) are closed nonvoid proper subsets of X . A being pervasive contains a couple j, k of functions satisfying

$$(3) \quad |g - j|_{\overline{X - F}} < \frac{e}{4c}, \quad |g - k|_{\overline{X - H}} < \frac{e}{4c},$$

where c is the constant from (1). Remark that c is not less than 1.

Without loss of generality we may assume that j and k are not both identically zero (in the opposite case the function $h = 0$ satisfies (2)) and put

$$(4) \quad \check{e} = \frac{e}{2(|j| + |k|)}.$$

Take, with regard to (1), an f in A for which

$$(5) \quad |f|_F < \check{e}, \quad |f - 1|_H < \check{e}, \quad |f| \leq c.$$

The function

$$h = fj + (1 - f)k,$$

satisfies (2). In fact, it is undeniable that

$$F \subset \overline{X - H}, \quad H \subset \overline{X - F}, \quad X = F \cup H \cup (\overline{X - F} \cap \overline{X - H}),$$

and, by (3), (4) and (5)

$$\begin{aligned} |g - h|_F &= |g - fj - (1 - f)k|_F \leq \\ &\leq |g - k|_F + |f|_F(|j|_F + |k|_F) \leq \\ &\leq |g - k|_{\overline{X - H}} + |f|_F(|j| + |k|) < \frac{e}{4c} + \frac{e}{2} < e, \end{aligned}$$

$$\begin{aligned} |g - h|_H &= |g - j + j - fj - (1 - f)k|_H \leq \\ &\leq |g - j|_{\overline{X - F}} + |1 - f|_H(|j| + |k|) < \frac{e}{4c} + \frac{e}{2} < e, \end{aligned}$$

and finally

$$|g - h|_{\overline{X - F} \cap \overline{X - H}} = |g - fk - (1 - f)k + fk - fj|_{\overline{X - F} \cap \overline{X - H}} \leq$$

$$\begin{aligned} &\leq |g - k|_{\overline{X-H}} + |f| \cdot |j - k|_{\overline{X-F} \cap \overline{X-H}} < \\ &< \frac{e}{4c} + c(|g - j|_{\overline{X-F}} + |g - k|_{\overline{X-H}}) < \frac{e}{4c} + \frac{e}{2} < e. \end{aligned}$$

Theorem B is proved.

Remark. Evidently, the condition of pervasiveness for A may be omitted; it suffices to require an approximation of any continuous function on $X - F$ and $X - H$ by functions in A , and a norm-bounded approximation of the function which is equal to 0 on F and to 1 on H on the set $F \cup H$ by functions in A .

Problem. So far we have proved the following:

Whenever A is a proper pervasive algebra (i.e., a pervasive algebra which is a proper subalgebra of $C(X)$) and F, H are arbitrary disjoint closed proper fat (i.e., with interior points) subsets of X , then any approximation of the function 0 on F and 1 on H is unbounded in the norm of A .

Now we ask the following question: *Is, in general, the assumption of F and H being fat necessary?*

For the classical disc algebra mentioned above this is not the case:

It is well-known that the disc algebra A is pervasive; it follows, for instance, from the famous Wermer's Maximality Theorem [3]; also it is well-known that any nontrivial analytic measure on C (i.e., a measure m on the unit circle C which annihilates A in the sense that $\int f dm = 0$ for any f in A) and the Lebesgue measure on C are mutually absolutely continuous — this is the classical F. and M. Riesz Theorem.

Let F and H be closed disjoint subsets of C having positive Lebesgue measures. Let $\{f_n\}_n$ be a sequence of functions in A which approximates 0 on F and 1 on H . Then $\{f_n\}_n$ is unbounded.

Admit the boundedness of $\{f_n\}$ and fix an arbitrary nontrivial analytic measure m . Then the sequence $\{f_n m\}_n$ is a norm-bounded sequence of analytic measures and has, in the weak-star topology, a limit point, say p . It is evident that

$$p|_F = 0, \quad p|_H = m|_H,$$

where $y|_Y$ denotes the restriction of the measure y to the set Y . However, p is analytic and F has a positive measure, hence p has to be trivial, and at the same time $p|_H \neq 0$, which is a contradiction.

References

- [1] *K. Hoffman, I. M. Singer*: Maximal algebras of continuous functions. *Acta Math.* 103 (1960), 217–241.
- [2] *J. Čerych*: A characterization of $C(X)$, *Comment. Math. Univ. Carolinae* 21, 2 (1980), 379–383.
- [3] *J. Wermer*: On algebras of continuous functions, *Proc. Amer. Math. Soc.* 4 (1953), 866–869.

Author's address: 186 00 Praha 8, Sokolovská 83 (Matematicko-fyzikální fakulta UK).