

Dmitrii B. Shakhmatov

On zero-dimensionality of subgroups of locally compact groups

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 32 (1991), No. 3, 581--582

Persistent URL: <http://dml.cz/dmlcz/118435>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1991

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## On zero-dimensionality of subgroups of locally compact groups

DMITRII B. SHAKHMATOV

*Abstract.* Improving the recent result of the author we show that  $\text{ind } H = 0$  is equivalent to  $\dim H = 0$  for every subgroup  $H$  of a Hausdorff locally compact group  $G$ .

*Keywords:* zero-dimensionality, covering dimension, inductive dimension, subgroup, locally compact group

*Classification:* Primary 22A05, 22D05, 54F45; Secondary 54D45, 54H99

All topological groups considered in this note are assumed to be Hausdorff. A subset of a topological space is said to be *clopen* if it is both open and closed. A topological space  $X$  is *zero-dimensional* if it has a base consisting of clopen sets (i.e. if  $\text{ind } X = 0$ ), and  $X$  is *strongly zero-dimensional* if every (locally) finite open cover of  $X$  consisting of functionally open sets has a finite disjoint clopen refinement (i.e. if  $\dim X = 0$ ). Strongly zero-dimensional spaces are zero-dimensional [1, Theorem 6.2.6] but not vice versa [1, Example 6.2.20]. However, the implication can be reversed for totally bounded groups: Recently, the author proved that a zero-dimensional subgroup of a compact group is strongly zero-dimensional [3, Corollary 3.4]. The aim of our note is to extend this result over subgroups of locally compact groups.

**Theorem.** *A zero-dimensional subgroup of a locally compact group is strongly zero-dimensional.*

In the proof of this theorem, we need the notion of  $\mathbb{R}$ -factorizable group [4]: A topological group  $G$  is  *$\mathbb{R}$ -factorizable* if for every real-valued continuous function  $f : G \rightarrow \mathbb{R}$  defined on  $G$  there exist a topological group  $H$  with a countable base, a continuous homomorphism  $\pi : G \rightarrow H$  and a continuous mapping  $\varphi : H \rightarrow \mathbb{R}$  such that  $f = \varphi \circ \pi$ . The following proposition formally improves [3, Theorem 3.3]:

**Proposition.** *A zero-dimensional topological group having an open  $\mathbb{R}$ -factorizable subgroup is strongly zero-dimensional.*

**PROOF:** Let  $H$  be an open  $\mathbb{R}$ -factorizable subgroup of a zero-dimensional topological group  $G$ . Being a subspace of a zero-dimensional space  $G$ ,  $H$  is zero-dimensional, and since  $H$  is  $\mathbb{R}$ -factorizable, it is strongly zero-dimensional by [3, Theorem 3.3].

---

The author would like to thank cordially Dikran Dikranjan for his suggestion that [3, Corollary 3.4] might admit the improvement stated in the main theorem.

Observe that  $H$  is clopen in  $G$ , as is every open subgroup of any topological group [2, Chapter 2, Theorem 5.5]. Since  $G$  can be covered by disjoint clopen copies of  $H$  (namely, by some translations of  $H$ ) and  $H$  is strongly zero-dimensional,  $G$  is also strongly zero-dimensional [1, Theorem 6.2.13].  $\square$

PROOF OF THEOREM: Let  $H$  be a zero-dimensional subgroup of a locally compact group  $G$ , and let  $U$  be an open neighbourhood of the neutral element of  $G$  having compact closure  $\bar{U}$  in  $G$ . Then  $G^*$ , the smallest subgroup of  $G$  that contains  $\bar{U}$ , is  $\sigma$ -compact. As a subgroup of a  $\sigma$ -compact group,  $H^* = H \cap G^*$  is  $\mathbb{R}$ -factorizable [4, Corollary 1.13]. Since  $H^*$  contains the non-empty open set  $U \cap H$  and is a subgroup of  $H$ ,  $H^*$  is open in  $H$  [2, Chapter 2, Theorem 5.5]. Now, Proposition finishes the proof.  $\square$

In conclusion, let us mention that, quite surprisingly, the following question remains open:

**Question.** Is there a normal zero-dimensional group which is not strongly zero-dimensional?

Even if we drop “normal” here, the answer to this question seems to be unknown.

#### REFERENCES

- [1] Engelking R., *General Topology*, Warszawa, PWN, 1977.
- [2] Hewitt E., Ross K.A., *Abstract Harmonic Analysis, vol. 1. Structure of Topological Groups. Integration Theory. Group Representations*, Die Grundlehren der mathematischen Wissenschaften, Bd. 115, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963.
- [3] Shakhmatov D.B., *Imbeddings into topological groups preserving dimensions*, Topology Appl. **36** (1990), 181–204.
- [4] Tkačenko M.G., *Factorization theorems for topological groups and their applications*, Topology Appl. **38** (1991), 21–37.

DEPARTMENT OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY,  
119 889 Moscow, USSR

(Received March 26, 1991)