

Vladimir Vladimirovich Tkachuk

A note on splittable spaces

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 33 (1992), No. 3, 551--555

Persistent URL: <http://dml.cz/dmlcz/118522>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1992

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## A note on splittable spaces

VLADIMIR V. TKACHUK

*Abstract.* A space  $X$  is splittable over a space  $Y$  (or splits over  $Y$ ) if for every  $A \subset X$  there exists a continuous map  $f : X \rightarrow Y$  with  $f^{-1}fA = A$ . We prove that any  $n$ -dimensional polyhedron splits over  $\mathbf{R}^{2n}$  but not necessarily over  $\mathbf{R}^{2n-2}$ . It is established that if a metrizable compact  $X$  splits over  $\mathbf{R}^n$ , then  $\dim X \leq n$ . An example of  $n$ -dimensional compact space which does not split over  $\mathbf{R}^{2n}$  is given.

*Keywords:* splittable, polyhedron, dimension

*Classification:* 54A25

The notion of splittability was introduced by A.V. Arhangel'skii [1]. A space  $X$  is splittable (or splits) over a space  $Y$  if for any  $A \subset X$  there exists a continuous map  $f : X \rightarrow Y$  such that  $f^{-1}fA = A$ . Many results were obtained by A.V. Arhangel'skii and D.B. Shakhmatov ([1]–[3]) on spaces splittable over  $\mathbf{R}^\omega$ . The author had also written a paper [4] on this topic.

Recently, A.V. Arhangel'skii had shown that a compact space  $X$  splits over  $\mathbf{R}$  iff it embeds in  $\mathbf{R}$ . He also proved that any 1-dimensional polyhedron splits over  $\mathbf{R}^2$  so that not every compact space splittable over  $\mathbf{R}^2$  embeds in  $\mathbf{R}^2$ . We prove here that any  $n$ -dimensional polyhedron splits over  $\mathbf{R}^{2n}$  but not necessarily over  $\mathbf{R}^{2n-2}$ . We establish also that there exists a compact space  $X_n \subset \mathbf{R}^{2n+1}$  with  $\dim X_n = n$  and not splittable over  $\mathbf{R}^{2n}$ . Another result is Corollary 8 answering Question 2 and 3 in [1].

**Notations and terminology.** All spaces under consideration are Tychonoff ones. Given two spaces  $X$  and  $Y$ , denote by  $C(X, Y)$  the set of all continuous functions from  $X$  to  $Y$ . The space  $\mathbf{R}$  is the real line with its usual topology and  $2 = \{0, 1\}$ . If  $x, y \in \mathbf{R}^n$  then  $[x, y] = \{tx + (1-t)y : 0 \leq t \leq 1\}$  is the segment connecting  $x$  and  $y$ ,  $|x - y|$  is its length. It is always clear from the context whether  $|\cdot|$  denotes cardinality of some set or length of a segment. Of course,  $[x, y] = \{tx + (1-t)y : 0 < t \leq 1\}$  and  $(x, y) = \{tx + (1-t)y : 0 < t < 1\}$ . The simplex in  $\mathbf{R}^n$  with vertices  $a_0, \dots, a_k$  will be denoted by  $[a_0, \dots, a_k]$ , then  $\langle T \rangle = \{t_0a_0 + \dots + t_ka_k : t_i > 0 \text{ for all } i \in k + 1 \text{ and } \sum_{i=0}^k t_i = 1\}$ . Let  $x \in \mathbf{R}^n$  and  $A \subset \mathbf{R}^n$ . Then  $\text{con}(x, A) = \bigcup\{[x, a] : a \in A\}$ .

Other notations are standard and can be found in [9].

**1. Lemma.** *Given any spaces  $X$  and  $Y$  and an infinite cardinal  $k$  with  $d(X) \leq k$ , ( $d$  is the density character),  $|Y| \leq 2^k$ , suppose that  $\bigcup\{X_s : s \in 2^k\} \subset X$ ,  $X_s \cap X_t = \emptyset$  if  $s \neq t$  and no  $X_s$  can be continuously injected into  $Y$ . Then  $X$  does not split over  $Y$ .*

PROOF: Clearly,  $|C(X, Y)| \leq 2^k$ , so let  $C(X, Y) = \{f_s : s < 2^k\}$ . For any  $s < 2^k$  there is an  $x_s \in X$  such that  $|f_s^{-1}f_s(x_s) \cap X_s| > 1$  because  $f \upharpoonright X$  is not an injection. Then the set  $A = \{x_s : s < 2^k\}$  witnesses non-splittability of  $X$  over  $Y$ .  $\square$

**2. Example.** For any natural  $n$  there exists an  $n$ -dimensional metrizable compact space  $X_n$  which does not split over  $\mathbf{R}^{2n}$ .

PROOF: Take any metrizable compact space  $Y_n$  with  $Y_n \not\hookrightarrow \mathbf{R}^{2n}$  and  $\dim Y_n = n$ . Then let  $X_n = 2^\omega \times Y$ . It is obvious that the family  $\{\{s\} \times Y_n : s \in 2^\omega\}$  satisfies the conditions of Lemma 1 for  $X = X_n$ ,  $k = \omega$  and  $Y = \mathbf{R}^{2n}$ . Hence  $X_n$  does not split over  $\mathbf{R}^{2n}$ .  $\square$

**3. Example.** There exists an  $n$ -dimensional compact polyhedron  $P_n$  which is not splittable over  $\mathbf{R}^{2n-2}$ .

PROOF: There exists an  $(n - 1)$ -dimensional compact polyhedron  $Y_n$  which does not embed in  $\mathbf{R}^{2n-2}$ . Then  $P = Y_n \times [0, 1]$  is what was required, because the family  $\{Y_n \times \{t\} : t \in [0, 1]\}$  satisfies the conditions of Lemma 1 for  $X = P_n$ ,  $k = \omega$  and  $Y = \mathbf{R}^{2n-2}$ .  $\square$

**4. Theorem.** Let  $P$  be an  $n$ -dimensional compact polyhedron. Then  $P$  splits over  $\mathbf{R}^{2n}$ .

PROOF: Denote by  $a_1, \dots, a_k$  the vertices of  $P$ . Let  $\{S_1, \dots, S_r\}$  be the set of all  $(n - 1)$ -dimensional simplexes from  $P$  and let  $\mu = \{T_1, \dots, T_m\}$  be some set of its  $n$ -dimensional ones. Take any hyperplane  $H \subset \mathbf{R}^{2n}$  and let  $b_1, \dots, b_k$  be some points generally positioned in  $H$ . Define a polyhedron  $Q_{n-1}$  in the following way: the vertices of  $Q_{n-1}$  are  $b_1, \dots, b_k$  and a simplex  $[b_{i_1}, \dots, b_{i_l}]$ ,  $l \leq n$  belongs to  $Q_{n-1}$  iff the simplex  $[a_{i_1}, \dots, a_{i_l}]$  belongs to  $P$ . If  $P_{n-1}$  is the union of all  $\leq (n - 1)$ -dimensional simplexes, then the simplicial map  $f : P_{n-1} \rightarrow Q_{n-1}$  defined by  $f(a_i) = b_i$  is a homeomorphism because  $H$  is isomorphic to  $\mathbf{R}^{2n-1}$ . Pick any  $D \in \mathbf{R}^{2n} \setminus H$ .  $\square$

**5. Lemma.** There exist  $m$  sets  $L_1, \dots, L_m$  and a continuous map

$$g = g(f, D) : P_{n-1} \cup T_1 \cup \dots \cup T_m \rightarrow \mathbf{R}^{2n}$$

with the following properties:

- (1)  $L_i$  is a subset of  $\mathbf{R}^{2n}_D$ , where the last set is the component of  $\mathbf{R}^{2n} \setminus H$  containing  $D$ ,  $i = 1, \dots, m$ ;
- (2)  $L_i$  is homeomorphic to  $\langle T_i \rangle$ ,  $i = 1, \dots, m$ ;
- (3)  $L_i \cap L_j = \{D\}$  if  $i \neq j$ ;
- (4)  $g \upharpoonright P_{n-1} = f$ ;
- (5)  $g \upharpoonright (P_{n-1} \cup T_i)$  is a homeomorphism onto  $Q_{n-1} \cup L_i$ .

PROOF OF THE LEMMA: Let  $R = \text{con}(D, Q_{n-1})$ ,  $R_i = \text{con}(D, f(S_i))$ ,  $i = 1, \dots, r$ . The set  $R$  being compact, there is a sphere (in  $\mathbf{R}^{2n}$ ) containing it. Pick any  $E \in \mathbf{R}^{2n}_D$  outside this sphere and not belonging to any of  $(2n - 1)$ -dimensional planes, spanned in  $\mathbf{R}^{2n}$  by some  $2n$  points from the set  $\{b_1, \dots, b_k, D\}$ .

We are going to construct a continuous function  $q : R \rightarrow [0, 1)$  such that

$$(6) \quad q^{-1}(0) = Q_{n-1} \cup \{D\};$$

$$(7) \quad \text{if } x \in R \setminus (Q_{n-1} \cup \{D\}) \text{ and } y \in (x, E) \cap R, \text{ then } q(x) < \frac{|x - y|}{|y - E|}.$$

To do that let  $W_i = \{x \in R : (x, E) \cap R_i \neq \emptyset\}$ . For each  $x \in R$  and each  $i$ , there is at most one point in  $(x, E) \cap R_i$ , for otherwise  $E$  would belong to the  $n$ -dimensional plane spanned by  $R_i$ . Put for  $x \in R_i$ :

$$r_i(x) = \frac{|x - y|}{|y - E|} \text{ where } y \in (x, E) \cap R_i, \text{ and } r_i(D) = 0.$$

Let us prove that  $\text{dom } r_i = W_i \cup \{D\}$  is a closed subset of  $R$ . It suffices to show that  $\text{dom } r_i \cap R_j$  is closed for each  $j$ . If  $R_i \cap R_j \cap H \neq \emptyset$  then, owing to the choice of  $E$ ,  $\text{dom } r_i \cap R_j = \{D\}$ . Suppose  $R_i \cap R_j \cap H = \emptyset$ , hence  $R_i \cap R_j = \{D\}$ . Let  $x \in R_j \setminus \text{dom } r_i$ . Then  $(x, E) \cap R_i = \emptyset$  and  $[x, E] \cap R_i = \emptyset$  as well. Since  $[x, E]$  is compact and  $R_i$  is closed, the distance between these sets is positive, say  $\varepsilon$ , and whenever  $|z - x| < \varepsilon$  then clearly  $[z, E] \cap R_i = \emptyset$ . Since  $r_i$  is continuous, there exists a continuous  $q_i : R \rightarrow [0, 1)$  with  $q^{-1}(0) = Q_{n-1} \cup \{D\}$  and  $q_i(x) < r_i(x)$  for all  $x \in W_i \setminus (Q_{n-1} \cup \{D\})$ . Now it suffices to put

$$q(x) = \min\{q_i(x) : i = 1, \dots, r\}.$$

Let  $M_i = \text{con}(D, f(T_i \setminus \langle T_i \rangle))$ ,  $i = 1, \dots, m$ . It is clear that  $M_i$  is homeomorphic to  $T_i$ . Define an injective continuous map  $s_i : M_i \rightarrow \mathbf{R}_D^{2n}$  in the following way: if  $x \in M_i$  then find the point  $y \in [x, E]$  with

$$\frac{|x - y|}{|y - E|} = \frac{q(x)}{i + 1}$$

and put  $s_i(x) = y$ .

Evidently,  $s_i$  is a homeomorphism. Let  $L_i = s_i(M_i \setminus f(T_i \setminus \langle T_i \rangle))$ . We are going to define the map  $g = g(f, D)$  and verify (1)–(5). Take any homeomorphism  $u_i : T_i \rightarrow M_i$  with  $u_i \upharpoonright (T_i \setminus \langle T_i \rangle) = f$ . Then let  $g(x)$  be equal to  $f(x)$  if  $x \in P_{n-1}$  and to  $s_i(u_i(x))$  for  $x \in T_i \setminus P_{n-1}$ ,  $i = 1, \dots, m$ .

Only (3) needs to be verified.

Let  $x \in M_i \setminus (\{D\} \cup Q_{n-1})$ ,  $y \in M_j \setminus (\{D\} \cup Q_{n-1})$ . If  $g(x) = g(y)$  then  $x, y$  and  $E$  are linearly dependent. We may assume without loss of generality that  $y \in [x, E]$ . There are two possibilities:  $x = y$ , and  $x \neq y$ .

If  $x = y$  then

$$\frac{|x - g(x)|}{|g(x) - E|} = \frac{q(x)}{i + 1} \quad \text{and} \quad \frac{|x - g(y)|}{|g(y) - E|} = \frac{q(y)}{j + 1} = \frac{q(x)}{j + 1} \neq \frac{q(x)}{i + 1},$$

so that  $g(x) \neq g(y)$ , which is a contradiction.

If  $x \neq y$  and  $x \in R_{t_1}, y \in R_{t_2}, R_{t_1} \cap R_{t_2} \cap Q_{n-1} \neq \emptyset$  then it is impossible that  $y \in [x, E]$  — a contradiction.

If  $R_{t_1} \cap R_{t_2} \cap Q_{n-1} = \emptyset$  then

$$\frac{|x - g(x)|}{|g(x) - E|} < \frac{|x - y|}{|y - E|}$$

and therefore  $g(x) \in [x, y]$  while  $g(y) \in [y, E]$  and  $g(x) \neq g(y)$  — a contradiction again and we established (3) together with our lemma.  $\square$

We have all we need to split  $P$  over  $\mathbf{R}^{2n}$ . Let  $A \subset P$ . Pick a point  $D_1 \in \mathbf{R}^{2n} \setminus (H \cup \mathbf{R}_D^{2n})$ . Let  $T_1, \dots, T_m, T_{m+1}, \dots, T_{m_1}$  be all  $n$ -dimensional simplexes of  $P$  numerated in such a way that  $A \cap \langle T_i \rangle \neq \emptyset, i = 1, \dots, m, (P \setminus A) \cap \langle T_i \rangle \neq \emptyset, i = m + 1, \dots, m_1$ . Using Lemma 5 find the sets  $L_1, \dots, L_{m_1}$  and maps  $g = g(f, D)$  and  $g_1 = g(f, D_1)$  such that

(8)  $L_i \subset \mathbf{R}_D^{2n}, i = 1, \dots, m, L_i \subset \mathbf{R}_{D_1}^{2n}, i = m + 1, \dots, m_1;$

(9)  $L_i$  is homeomorphic to  $\langle T_i \rangle, i = 1, \dots, m_1;$

(10)  $L_i \cap L_j = \{D\}, i \neq j, j \in 1, \dots, m;$

(11)  $L_i \cap L_j = \{D\}, i = j, i, j \in m + 1, \dots, m_1;$

(12)  $g \upharpoonright P_{n-1} = g_1 \upharpoonright P_{n-1} = f;$

(13)  $g \upharpoonright (P_{n-1} \cup T_i)$  is a homeomorphism onto  $Q_{n-1} \cup L_i, i = 1, \dots, m;$

(14)  $g \upharpoonright (P_{n-1} \cup T_i)$  is a homeomorphism onto  $Q_{n-1} \cup L_i, i = m + 1, \dots, m_1;$

Pick some points  $c_1, \dots, c_{m_1}$  with  $c_i \in A \cap \langle T_i \rangle, i = 1, \dots, m, c_i \in (P \setminus A) \cap \langle T_i \rangle, i = m + 1, \dots, m_1$  and the points  $d_1, \dots, d_{m_1}$  with  $g(d_i) = D, i = 1, \dots, m, g_1(d_i) = D_1, i = m + 1, \dots, m_1$  (observe that automatically  $d_i \in \langle T_i \rangle$  for each  $i$ ). Let  $G = g \cup g_1$ . Then  $G$  is a continuous map,  $G : P \rightarrow \mathbf{R}^{2n}$ . There exists a homeomorphism  $h : P \rightarrow P$  with  $h \upharpoonright P_{n-1} = \text{id}_{P_{n-1}}$  and  $h(c_i) = d_i, i = 1, \dots, m_1$ . The map  $F = G \circ h$  separates  $A$  from  $P \setminus A$ , because  $|F^{-1}(x)| = 1$  if  $x \notin \{D, D_1\}$  and  $F^{-1}(D) = \{c_1, \dots, c_m\} \subset A, F^{-1}(D_1) = \{c_{m+1}, \dots, c_{m_1}\} \subset P \setminus A$ , and our theorem is proved.  $\square$

**6. Proposition.** *Let  $X$  be a compact space and  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and any compact  $K \subset X_i$  is scattered ( $i = 1, 2$ ). Assume that  $X$  splits over a space  $Y$  with  $\dim Y \leq n$ . Then  $\dim X \leq n$ .*

PROOF: Take a continuous  $f : X \rightarrow Y$  with  $f^{-1}f(X_1) = X_1$ . If  $y \in Y$  then  $f^{-1}(y)$  is a compact subset of some  $X_i$  ( $i = 1, 2$ ) and is thus scattered. Hence  $\dim f^{-1}(y) = 0$  for every  $y \in f(X)$ . But  $\dim X \leq \dim f(X) + \dim f \leq n$  [10] and the proof is over.  $\square$

**7. Corollary.** *If a compact space  $X$  is splittable over  $\mathbf{R}^n$ , then  $\dim X \leq n$ .*

PROOF: The space  $X$  must be metrizable [3]. It is widely known (see e.g. [5]) that metrizable compact spaces satisfy the assumptions of Proposition 6, so our proof is over.  $\square$

This corollary answers Questions 2 and 3 in [1].

**8. Corollary ( $ACP^\#$ ).** *If  $X$  is a compact space splittable over a space  $Y$  then  $\dim X \leq \dim Y$ . (The definition of  $ACP^\#$  can be found in [5]).*

**9. Corollary.** *If  $X$  is a metrizable compact space splittable over a space  $Y$  then  $\dim X \leq \dim Y$ .*

**10. Example.** *Compactness is essential in 7–9, for there exist infinite-dimensional second countable spaces which can be injectively mapped in  $\mathbf{R}$  [6].*

**11. Proposition.** *If  $X$  is an infinite extremally disconnected compact space splittable over a space  $Y$  then  $\beta\omega \hookrightarrow Y$ .*

PROOF: It is true in ZFC (see [7]) that  $X = X_1 \cup X_2$ ,  $X_1 \cap X_2 = \emptyset$  and every compact  $K \subset X$  is finite ( $i = 1, 2$ ). Pick a continuous map  $f : X \rightarrow Y$  with  $f^{-1}f(X_i) = X_i$ . The space  $\beta\omega$  embeds in  $X$  and  $f \upharpoonright \beta\omega$  has finite point-inverses, so that  $\beta\omega \hookrightarrow f(\beta\omega)$  [8] and our proposition is proved.

**12. Corollary.** *If  $\beta\omega$  splits over a space  $Y$  then  $\beta\omega$  embeds in  $Y$ .*

□

## REFERENCES

- [1] Arhangel'skii A.V., *A general concept of splittability of a topological space* (in Russian), in: Proceedings of the Fifth Tyraspol Symposium on General Topology and its Applications, Kishinev, Shtiintsa, 1985, 8–10.
- [2] Arhangel'skii A.V., Shakhmatov D.B., *Splittable spaces and questions of functions approximations* (in Russian), in: Proceedings of the Fifth Tyraspol Symposium on General Topology and its Applications, Kishinev, Shtiintsa, 1985, 10–11.
- [3] ———, *On pointwise approximation of arbitrary functions by countable families of continuous functions* (in Russian), Trudy Seminara I.G. Petrovskogo **13** (1988), 206–227.
- [4] Tkachuk V.V., *Approximation of  $\mathbf{R}^X$  with countable subsets of  $C_p(X)$  and calibers of the space  $C_p(X)$* , Comment. Math. Univ. Carolinae **27** (1986), 267–276.
- [5] Bregman Yu.H., Šapirovskii B.E., Šostak A.P., *On partition of topological spaces*, Časopis pro pěstování mat. **109** (1984), 27–53.
- [6] Mazurkiewicz S., *Sur les problèmes  $\kappa$  et  $\lambda$  de Urysohn*, Fund. Math. **10** (1927), 311–319.
- [7] Tkachuk V., *Remainders over discrete spaces — some applications* (in Russian), Vestnik MGU, Mat., Mech., no. 4, 1990, 18–21.
- [8] Malyhin V.I.,  *$\beta N$  is prime*, Bull. Acad. Polon. Sci., Ser. Mat. **27** (1979), 295–297.
- [9] Engelking R., *General Topology*, PWN, Warszawa, 1977.
- [10] Skljarenko E.G., *A theorem on maps, lowering dimension (in Russian)*, Bull. Acad. Polon. Sci., Ser. Mat. **10** (1962), 429–432.

DEPARTMENT OF MECHANICS AND MATHEMATICS, SECTION OF GENERAL TOPOLOGY AND GEOMETRY, MOSCOW STATE UNIVERSITY, 119899 MOSCOW, RUSSIA

(Received May 6, 1991, revised April 21, 1992)