

Michal Fečkan

Problems with nonlinear boundary value conditions

Commentationes Mathematicae Universitatis Carolinae, Vol. 33 (1992), No. 4, 597--604

Persistent URL: <http://dml.cz/dmlcz/118530>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1992

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Problems with nonlinear boundary value conditions

MICHAL FEČKAN

Abstract. The existence and multiplicity results are shown for certain types of problems with nonlinear boundary value conditions.

Keywords: nonlinear boundary value problems, multiple solutions, Melnikov functions

Classification: 34B15, 34L30

Introduction.

The purpose of this paper is to study several problems with nonlinear boundary value conditions. Mostly we study problems which are small perturbations of linear boundary value problems. The author was stimulated by the paper [1]; but in this paper we shall use several approaches to solve our problems: the implicit function theorem, the Mawhin coincidence degree theory, the Nielsen fixed point theory and when an unperturbed linear boundary value condition is a periodic one, we derive a Melnikov function for this problem [2].

Results.

We study

$$(1 - \varepsilon) \quad \begin{aligned} x' &= f_1(x) + \varepsilon \cdot f_2(t, x) \\ Ax(0) + Bx(T) &= \varepsilon \cdot \phi(x(0), x(T)), \end{aligned}$$

where f_1, f_2, ϕ are continuous on $R^m, R \times R^m, R^m \times R^m$, respectively, $A, B \in \mathcal{L}(R^m), T > 0, \varepsilon \in R$ is small.

Theorem 1. *Let us consider $(1 - \varepsilon)$ for the case when $f_1(c_0) = 0$ for some $c_0 \in R^m, Ac_0 + Bc_0 = 0, f_1, f_2, \phi$ are C^1 -smooth. If $\det(A + B \cdot e^{Df_1(c_0) \cdot T}) \neq 0$, then $(1 - \varepsilon)$ has a solution x_ε defined on $[0, T]$ for each ε small satisfying $x_\varepsilon(\cdot) \rightarrow c_0$ as $\varepsilon \rightarrow 0$.*

PROOF: We consider

$$\begin{aligned} F_\varepsilon: C^1 &\rightarrow C^0 \times R^m \\ F_\varepsilon(x) &= (x' - f_1(x) - \varepsilon f_2(t, x), Ax(0) + Bx(T) - \varepsilon \phi(x(0), x(T))). \end{aligned}$$

We see that

$$\begin{aligned} F_0(c_0) &= 0 \\ D_x F_0(c_0)v &= (v' - Df_1(c_0)v, Av(0) + Bv(T)). \end{aligned}$$

Thus $\ker D_x F_0(c_0) = \{v \mid v' = Df_1(c_0)v, Av(0) + Bv(T) = 0\}$ and by using our assumptions we have $\ker D_x F_0(c_0) = \{0\}$.

Let us solve

$$\begin{aligned} v' - Df_1(c_0)v &= r \\ Av(0) + Bv(T) &= w. \end{aligned}$$

Then $v(t) = \int_0^t e^{Df_1(c_0)(t-s)} r(s) ds + e^{Df_1(c_0)t} c$. Hence

$$Ac + B.e^{Df_1(c_0)T} c = -B \int_0^T e^{Df_1(c_0)(T-s)} r(s) ds$$

and the last equation we can solve in c . This completes the proof, since we can use the implicit function theorem. □

Corollary 2. *Let $f_1 \equiv 0$. Then the conditions of Theorem 1 are $c_0 = 0$, $A + B$ is invertible.*

Now we consider $(1 - \varepsilon)$ for the case $A = -B = Id$ and $x' = f_1(x)$ has an isolated T -periodic nonconstant solution $x_0(\cdot)$. Hence $(1-\varepsilon)$ has the form

$$(2 - \varepsilon) \quad \begin{aligned} x' &= f_1(x) + \varepsilon.f_2(t, x) \\ x(0) - x(T) &= \varepsilon.\phi(x(0), x(T)). \end{aligned}$$

Let ϕ, f_1, f_2 be C^2 -smooth mappings. We note that $(2 - 0)$ has the family of solutions $\Gamma = \{x_0(\cdot + c), c \in [0, T]\}$. We are interested in bifurcations of solutions of $(2-\varepsilon)$ from Γ for ε small. We apply the following theorem from [5, pp. 397]:

Theorem 3. *Let $F_\varepsilon: X \rightarrow Y$ be a C^2 -smooth mapping, X, Y are Hilbert spaces and F_0 possesses a compact C^2 -manifold \mathcal{M} such that $F_0(\mathcal{M}) = 0$, $\ker D_x F_0(m) = T_m \mathcal{M}$, $\text{index } D_x F_0(m) = 0$, $D_x F_0(m)$ is a Fredholm operator for each $m \in \mathcal{M}$. Here $T_m \mathcal{M}$ is the tangent space of \mathcal{M} at m . Let $P(m) \in \mathcal{L}(Y)$ be the orthogonal projection onto $(\text{im } D_x F_0(m))^\perp$ for each $m \in \mathcal{M}$. Consider the map $M(m) = P(m).D_\varepsilon F_0(m)$, $M: \mathcal{M} \rightarrow Y$. If there is a $m_0 \in \mathcal{M}$ such that $M(m_0) = 0$, $DM(m_0)$ is injective. Then for any ε small the equation $F_\varepsilon(m) = 0$ has a solution near \mathcal{M} . We note that M can be considered as a map from $R^{\dim \mathcal{M}}$ into $R^{\dim \mathcal{M}}$ in the local coordinates.*

We shall derive M for the special case $(2 - \varepsilon)$. We put $X = H^1([0, T], R^m)$, $Y = H^0([0, T], R^m) \times R^m$, $F_\varepsilon(x) = (x' - f_1(x) - \varepsilon.f_2(t, x), x(0) - x(T) - \varepsilon.\phi(x(0), x(T)))$ and $\mathcal{M} = \{x_0(\cdot + c), c \in [0, T]\}$. Hence \mathcal{M} is homeomorphic to a circle and

$$D_x F_0(m)v = (v' - Df_1(x_0(\cdot + c)).v, v(0) - v(T)).$$

Since x_0 is an isolated T -periodic nonconstant solution of (2 - 0) we have $\ker D_x F_0(m) = T_m \mathcal{M}$ for each $m \in \mathcal{M}$. Now we derive $\text{im } D_x F_0(m)$ and thus let us solve

$$\begin{aligned} v' - Df_1(x_0(\cdot + c))v &= r \\ v(0) - v(T) &= v_1 \in R^m. \end{aligned}$$

We put $w(t) = v(t) + \frac{t v_1}{T}$, hence

$$\begin{aligned} w' - Df_1(x_0(t + c))w &= r + \frac{v_1}{T} - Df_1(x_0(t + c)) \cdot \frac{v_1}{T} \\ w(0) &= w(T). \end{aligned}$$

It is well-known that this equation has a solution if and only if

$$\int_0^T \tilde{x}_0(s + c) \cdot (r(s) + \frac{v_1}{T} - Df_1(x_0(s + c)) \cdot \frac{v_1}{T}) ds = 0,$$

where \tilde{x}_0 is a nonzero T -periodic solution of $x' + (Df_1(x_0))^\top x = 0$.

Hence $(r, v_1) \in \text{im } D_x F_0(x_0(\cdot + c))$ if and only if

$$\begin{aligned} \langle w(c), (r, v_1) \rangle_Y &= \\ &= \int_0^T \tilde{x}_0(s + c) \cdot r(s) ds + \frac{1}{T} \int_0^T \tilde{x}_0(s + c) (v_1 - Df_1(x_0(s + c)) \cdot v_1) ds = 0, \end{aligned}$$

where $\langle \cdot, \cdot \rangle_Y$ is the scalar product on Y . Then

$$P(x_0(\cdot + c))w_1 = \langle w(c), w_1 \rangle_Y \cdot \frac{1}{\|w(c)\|_Y} \cdot w(c)$$

and

$$M(c) = \langle w(c), (-f_2(\cdot, x_0(\cdot + c)), -\phi(x_0(\cdot + c), x_0(\cdot + c))) \rangle_Y / \|w(c)\|_Y \cdot w(c).$$

Now we shall use the fact: let $\frac{a(c)}{b(c)} = d(c)$, where a, b, d are real smooth functions, $b(c_0) \neq 0$. Then for $a(c_0) = 0$ it follows $d'(c_0) \neq 0$ if and only if $a'(c_0) \neq 0$. Thus instead of $M(c)$ we can consider the map

$$\begin{aligned} \bar{M}(c) &= \langle w(c), (f_2(\cdot, x_0(\cdot + c)), \phi(x_0(\cdot + c), x_0(\cdot + c))) \rangle_Y = \\ &= \int_0^T \tilde{x}_0(s + c) \cdot f_2(s, x_0(s + c)) ds + \\ &+ \frac{1}{T} \int_0^T \tilde{x}_0(s + c) \cdot (\phi(x_0(s + c), x_0(s + c)) - \\ &- Df_1(x_0(s + c)) \cdot \phi(x_0(s + c), x_0(s + c))) ds. \end{aligned}$$

Summing up we obtain

Theorem 4. *If there is $c_0 \in [0, T]$ such that $\bar{M}(c_0) = 0$, $\bar{M}'(c_0) \neq 0$, then for each ε small, $(2 - \varepsilon)$ has a solution on $[0, T]$.*

Remark 5. We see that for $\phi \equiv 0$ \bar{M} is the subharmonic Melnikov function [2] and thus \bar{M} we can consider as a Melnikov function for $(2 - \varepsilon)$.

Now we consider

$$(3) \quad \begin{aligned} x' &= f(t, x) \\ Ax(0) + Bx(T) &= \phi(x(0), x(T)), \end{aligned}$$

where f, ϕ are continuous. Let $G \subset R^m$ be an open bounded subset, $0 \in G$.

Theorem 6. *Assume that*

- (i) $x' = \lambda f(t, x)$, $Ax(0) + Bx(T) = \lambda \phi(x(0), x(T))$ has no solution for each $\lambda \in (0, 1)$ satisfying $x(\cdot) \subset \bar{G}$, $x(\cdot) \cap \partial G \neq \emptyset$.

Moreover

- (ii) $D = \{z \in R^m \mid Az + Bz = 0, z \in G\} \neq \{0\}$, $g(z) \neq 0$ for each $z \in \partial D$, where

$$g(z) = JP(\phi(z), z) - B \cdot \int_0^T f(s, z) ds$$

here $P: R^m \rightarrow (\text{im}(A + B))^\perp$ is a projection and

$J: (\text{im}(A + B))^\perp \rightarrow \{z, Az + Bz = 0\}$ is an isomorphism.

- (iii) $\text{deg}(g, D, 0) \neq 0$.

Then (3) has a solution x , $x(\cdot) \subset G$.

PROOF: We shall apply a theorem of Mawhin [3, p. 41]. We put

$$\begin{aligned} X &= C^0([0, T], R^m), Y = X \times R^m \\ Lx &= (x', Ax(0) + Bx(T)) \\ N(x) &= (f(\cdot, x), \phi(x(0), x(T))) \\ \Omega &= \{x \in X, x(\cdot) \in G\}. \end{aligned}$$

By our assumptions $Lx = \lambda N(x)$, $\lambda \in (0, 1)$ has no solution on $\partial\Omega$. We compute $\ker L = \{x' = 0, Ax(0) + Bx(T) = 0\} = \{x \mid x = \text{constant} = c_1, Ac_1 + Bc_1 = 0\}$. Now $\text{im } L = \{(v, w) \mid x' = v, Ax(0) + Bx(T) = w\}$. But

$$\begin{aligned} x(t) &= \int_0^t v(s) ds + c_1, \quad Ac_1 + B \int_0^T v(s) ds + Bc_1 = w, \\ Ac_1 + Bc_1 &= w - B \cdot \int_0^T v(s) ds. \end{aligned}$$

This equation has a solution if and only if

$$P(w - B. \int_0^T v(s) ds) = 0.$$

Hence

$$\text{im } L = \{(v, w), P(w - B. \int_0^T v(s) ds) = 0\}.$$

Thus $\dim \text{coker } \text{im } L = \dim \ker L \neq 0$. We take

$$\bar{P}(v, w) = (0, P(w - B. \int_0^T v(s) ds)).$$

Then $\text{im}(I - \bar{P}) = \text{im } L$.

Finally consider the map

$$J.\bar{P}.N / \ker L \cap \Omega \rightarrow 0 \times R^m \circlearrowright R^m$$

defined in the following way

$$g(z) = J.P(\phi(z, z) - B. \int_0^T f(s, z) ds), \quad z \in D.$$

Since $g(z) \neq 0$ for $z \in \partial D$ and $\text{deg}(g, D, 0) \neq 0$ we see that also the last assumption of the theorem of Mawhin is satisfied. The proof is finished. □

Theorem 7. *Let us consider*

$$(4) \quad \begin{aligned} x' &= \varepsilon.f(t, x) \\ Ax(0) + Bx(T) &= \varepsilon.\phi(x(0), x(T)) \end{aligned}$$

and assume the existence of G as in Theorem 6 possessing the properties (ii), (iii). Then (4) has a solution for each ε small.

PROOF: The proof is similar as for Theorem 6. □

Theorem 7 expresses only the existence result. Now we shall apply a theorem of [4] to show a multiplicity result.

Theorem 8 ([4]). *Let $X \subset Y$ be Banach spaces, X is compactly embedded into Y . Consider $Lx = \varepsilon N(x)$, where $L: X \rightarrow Y$ is continuous, linear, Fredholm with index $L = 0$, $\ker L \neq \{0\}$ and $N: Y \rightarrow Y$ maps bounded sets into bounded sets, continuous. Moreover we assume that the map $\Pi(z) = z + JPN(z)$ is μ -retractible onto S with a retraction π , where J is an isomorphism from $\text{im } P$ onto $\ker L$, $P: Y \rightarrow Y$ is a projection, $\text{im}(I - P) = \text{im } L$, S is a compact, nonempty, locally contractible subset of $\ker L$, $\mu > 0$. Then the equation $Lx = \varepsilon.N(x)$ has for each ε small at least $N(\pi.\Pi)$ solutions. Here $N(\pi.\Pi)$ is the Nielsen number of the map $\pi.\Pi/S: S \rightarrow S$.*

Theorem 9. Consider (4) and assume that there is S a compact, nonempty, locally contractible subset of $\{c \in R^m, Ac + Bc = 0\} = W$ and the map

$$\psi(z) = z + JP(\phi(z, z) - B \int_0^T f(s, z) ds), \psi: W \rightarrow W$$

is μ -retractible onto S with respect to π . Then (4) has at least $N(\pi.\psi)$ solutions for each ε small. (The operators J, P are from Theorem 6.)

PROOF: We put

$$X = C^1([0, T], R^m), Y = C^0([0, T], R^m) \times R^m,$$

L, N as in the proof of Theorem 6. It is clear that $\Pi = \psi$ and thus the assertion follows by Theorem 8. □

Example 1. Consider

$$\begin{aligned} x'_1 &= \varepsilon.f_1(t, x_1, x_2), 0 \leq t \leq T \\ x'_2 &= \varepsilon.f_2(t, x_1, x_2) \\ (5 - \varepsilon) \quad a_1x_1(0) + a_2x_2(0) &= \varepsilon.\phi_1(x_1(0), x_2(0)) \\ b_1x_1(T) + b_2x_2(T) &= \varepsilon.\phi_2(x_1(T), x_2(T)). \end{aligned}$$

In this case

$$A = \begin{pmatrix} a_1 & a_2 \\ 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 \\ b_1 & b_2 \end{pmatrix}.$$

Hence

$$A + B = \begin{pmatrix} a_1 & a_2 \\ b_1 & b_2 \end{pmatrix}.$$

Applying Corollary 2 we obtain

Proposition 10. If $a_1.b_2 - a_2.b_1 \neq 0$ and f_1, f_2, ϕ_1, ϕ_2 are C^1 -smooth then (5- ε) has a solution for each ε small tending to 0 as $\varepsilon \rightarrow 0$.

Consider the case $a_1.b_2 - a_2.b_1 = 0, a_1^2 + a_2^2 \neq 0 \neq b_1^2 + b_2^2$. Then (see Theorem 9)

$$\begin{aligned} W &= \{(c_1, c_2) \mid a_1c_1 + a_2c_2 = 0, b_1c_1 + b_2c_2 = 0\} \\ &= \{c.(a_2, -a_1), c \in R\} \\ (\text{im}(A + B))^\perp &= \{c.(b_1, -a_1), c \in R\}. \end{aligned}$$

Hence

$$\begin{aligned} P(v_1, v_2) &= \frac{(v_1b_1 - v_2a_1)}{b_1^2 + a_1^2}(b_1, -a_1) \\ J(c.(b_1, -a_1)) &= c.(b_1^2 + a_1^2).(a_2, -a_1). \end{aligned}$$

Thus

$$g(c) = b_1 \phi_1(c, a_2, -c, a_1) + b_1 \cdot a_1 \int_0^T f_1(s, c, a_2, -c, a_1) ds - a_1 \cdot \phi_2(c, a_2, -c, a_1) + b_2 \cdot a_1 \int_0^T f_2(s, c, a_2, -c, a_1) ds,$$

since $\dim W = 1$.

Proposition 11. *Let f_1, f_2, ϕ_1, ϕ_2 be continuous and*

$$\limsup_{|c| \rightarrow \infty} g(c)/c > 0 \text{ or } \liminf_{|c| \rightarrow \infty} g(c)/c < 0.$$

Then $(5 - \varepsilon)$ has a solution for each ε small.

PROOF: The assertion follows by Theorem 7. □

Example 2. Consider

$$(6 - \varepsilon) \quad \begin{aligned} x'_1 &= \varepsilon \cdot f_1(t, x_1, x_2), \quad 0 \leq t \leq T \\ x'_2 &= \varepsilon \cdot f_2(t, x_1, x_2) \\ a_1 x_1(0) + a_2 x_1(T) &= \varepsilon \cdot \phi_1(x_1(0), x_2(0)) \\ b_1 x_2(0) + b_2 x_2(T) &= \varepsilon \cdot \phi_2(x_1(T), x_2(T)). \end{aligned}$$

In this case

$$A = \begin{pmatrix} a_1 & 0 \\ 0 & b_1 \end{pmatrix}, \quad B = \begin{pmatrix} a_2 & 0 \\ 0 & b_2 \end{pmatrix}.$$

Hence

$$A + B = \begin{pmatrix} a_1 + a_2 & 0 \\ 0 & b_1 + b_2 \end{pmatrix}.$$

According to Corollary 2 we obtain

Proposition 12. *If $(a_1 + a_2)(b_1 + b_2) \neq 0$ and f_1, f_2, ϕ_1, ϕ_2 are C^1 -smooth then $(6 - \varepsilon)$ has a solution for each ε small tending to 0 as $\varepsilon \rightarrow 0$.*

Let $a_1 = -a_2 \neq 0, b_1 + b_2 \neq 0$. Then (see Theorem 9)

$$\begin{aligned} W &= \{(c, 0), c \in R\} \\ P(v_1, v_2) &= (v_1, 0) \\ (\text{im}(A + B))^\perp &= \{(c, 0), c \in R\} \\ J((c, 0)) &= c. \end{aligned}$$

Thus

$$(7) \quad g(c) = \phi_1(c, 0) - a_2 \cdot \int_0^T f_1(s, c, 0) ds.$$

Proposition 13. *Let f_1, f_2, ϕ_1, ϕ_2 be continuous and*

$$\limsup_{|c| \rightarrow \infty} g(c)/c > 0 \quad \text{or} \quad \liminf_{|c| \rightarrow \infty} g(c)/c < 0.$$

Then $(6 - \varepsilon)$ has a solution for each ε small. Here g is defined by (7).

Lastly, consider $a_1 = -a_2 \neq 0, b_1 = -b_2 \neq 0$. Then (see Theorem 9) $W = R^2, P = J = Id$ and

$$\psi(c_1, c_2) = (8) \quad (c_1 + \phi_1(c_1, c_2) - a_2 \int_0^T f_1(s, c_1, c_2) ds, c_2 + \phi_2(c_1, c_2) - b_2 \int_0^T f_2(s, c_1, c_2) ds)$$

$$\psi: R^2 \rightarrow R^2.$$

Applying Theorem 9 we obtain

Proposition 14. *Let f_1, f_2, ϕ_1, ϕ_2 be continuous and S be a compact, locally contractible subset of R^2 . If the map ψ defined by (8) is μ -retractible onto S with respect to a retraction π then $(6 - \varepsilon)$ has at least $N(\pi, \psi)$ solutions for any ε small.*

To be more concrete we take $S = A_{r,p} = \{z \in R^2, r \leq |z| \leq p\}$ for fixed $0 < r < p$. We have constructed in [4] a family of mappings q for each $m \in \mathcal{N} \setminus \{1\}$ satisfying $N(\rho_{r,p}, q) = m - 1$, where $\rho_{r,p}$ is the usual retraction on $A_{r,p}$ (see [4]) and q is μ -retractible onto $A_{r,p}$ with respect to $\rho_{r,p}$ for some $\mu > 0$.

If $T = 1 = a_2 = b_2$ and

$$(9) \quad \begin{aligned} f_i(s, c_1, c_2) &= 2 \cdot q_i(c_1, c_2) \cdot s \\ \phi_i(c_1, c_2) &= 2 \cdot q_i(c_1, c_2) - c_i, \quad i = 1, 2, \end{aligned}$$

where $q = (q_1, q_2)$. Then easy computations show that the map ψ from (8) has the form $\psi = q$ and $\pi = \rho_{r,p}$. Summing up we have

Proposition 15. *Consider the special case (9) of the problem discussed in Proposition 14. Then in this case $(6 - \varepsilon)$ has at least $m - 1$ solutions for each ε small.*

REFERENCES

- [1] Samoilenko A.M., Le Loyong Tai, *On a method of study of boundary value problems with nonlinear boundary value conditions* (in Russian), Ukrainian Math. Journal **42** (1990), 951–957.
- [2] Guckenheimer J., Holmes P., *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer-Verlag, New York, 1983.
- [3] Mawhin J., *Nonlinear functional analysis and periodic solutions of ordinary differential equations*, in Summer School on Ordinary Differential Equations-Difford 74, Brno (1974), 37–60.
- [4] Fečkan M., *Nielsen fixed point theory and nonlinear equations*, to appear in Journal of Differential Equations.
- [5] Chow S.-N., Hale J.K., *Methods of Bifurcations Theory*, Springer-Verlag, New York, 1982.

MATHEMATICAL INSTITUTE, SLOVAK ACADEMY OF SCIENCES, ŠTEFÁNKOVA 49, BRATISLAVA, CZECHOSLOVAKIA

(Received January 17, 1992)