

Jiří Adámek; Jan Reiterman

The category of uniform spaces as a completion of the category of metric spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 33 (1992), No. 4, 689–693

Persistent URL: <http://dml.cz/dmlcz/118540>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1992

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

The category of uniform spaces as a completion of the category of metric spaces

JIŘÍ ADÁMEK, JAN REITERMAN

Abstract. A criterion for the existence of an initial completion of a concrete category \mathbf{K} universal w.r.t. finite products and subobjects is presented. For \mathbf{K} = metric spaces and uniformly continuous maps this completion is the category of uniform spaces.

Keywords: universal completion, metric space, uniform space

Classification: 18A35, 54B30, 54E15

Introduction.

We investigate initial completions of concrete categories, i.e. initially complete categories \mathbf{K}^* such that \mathbf{K} is a full, concrete subcategory of \mathbf{K}^* . Recall that \mathbf{K}^* is said to be a universal initial completion of \mathbf{K} , see [H], provided that (a) \mathbf{K} is closed under initial sources in \mathbf{K}^* and (b) for each initially complete \mathbf{L} and each concrete functor $F : \mathbf{K} \rightarrow \mathbf{L}$ preserving initial sources there exists an extension to an initial-sources preserving functor $F^* : \mathbf{K}^* \rightarrow \mathbf{L}$, unique up-to natural isomorphism. More in general, let Δ be a collection of sources in the base-category, then a Δ -universal initial completion of \mathbf{K} , see [E], is an initial completion \mathbf{K}^* such that

- (a) \mathbf{K} is closed in \mathbf{K}^* under initial sources carried by Δ -sources (shortly: initial Δ -sources)

and

- (b) for each initially complete category \mathbf{L} and each concrete functor $F : \mathbf{K} \rightarrow \mathbf{L}$ preserving initial Δ -sources there exists an extension to an initial-sources preserving functor $F^* : \mathbf{K}^* \rightarrow \mathbf{L}$, unique up-to a natural isomorphism.

A description of the Δ -universal initial completion (as a category of “ Δ -complete sources”) has been presented in [E]. Unfortunately, the conclusion made in that paper that \mathbf{K}^* is always legitimate (i.e. lives in the universe U of classes), formulated in Theorem 4, is false: For example, if \mathbf{K} is a discrete, large category, then the Δ -universal completion is illegitimate, being codable by the collection of all subclasses of \mathbf{K} . In the case $\Delta =$ all sources or $\Delta = \emptyset$, the legitimacy of the Δ -universal completion is characterized in [AHS].

In the present paper we concentrate on the case

$$\Delta_{fm} = \text{all finite products and monomorphisms.}$$

(Or, equivalently, all finite monosources.) We prove that each fibre-small, hereditary concrete category has a fibre-small (thus, legitimate) Δ_{fm} -universal initial completion. For

$\mathbf{K} = \mathbf{Met}$, metric spaces and uniformly continuous maps

this completion is

$\mathbf{K}^* = \mathbf{Unif}$, uniform spaces and uniformly continuous maps.

In this special case the same result also holds for $\Delta =$ all finite sources, or $\Delta =$ all countable sources, but it does not hold for $\Delta =$ all sources. In fact, a concrete description of the universal initial completion of \mathbf{Met} is not known.

The paper has been inspired by Zdeněk Frolík who asked us about a categorical motivation of uniform spaces from the point of view of metric spaces. This paper is, most unfortunately, the end of a long, fruitful, and happy collaboration of the authors: Jan Reiterman died precisely when it was completed.

I. Δ -universal completion.

Recall from [A] that a construct (i.e. a concrete category over \mathbf{Set}) is called hereditary provided that every subset of the underlying set of any object K gives rise to an initial subobject of K . This can be generalized to concrete categories over a base category \mathbf{X} (i.e. pairs consisting of a category \mathbf{K} and a faithful, amnesic functor $|\cdot| : \mathbf{K} \rightarrow \mathbf{X}$) provided that a fixed factorization structure (\mathbf{E}, \mathbf{M}) for \mathbf{X} -morphisms is given:

Definition. A concrete category \mathbf{K} over an (\mathbf{E}, \mathbf{M}) -base category is called hereditary provided that given an object K of \mathbf{K} , every \mathbf{M} -morphism $m : X \rightarrow |K|$ has an initial lift.

Recall that a concrete category is fibre-small provided that for each object X of the underlying category the fibre $\{K \in \mathbf{K} \mid |K| = X\}$ is a set.

Theorem. *Let \mathbf{X} be an \mathbf{E} -co-wellpowered (\mathbf{E}, \mathbf{M}) -category, and let Δ be a collection of sources containing all \mathbf{M} -maps (considered as singleton sources). Then each hereditary, fibre-small concrete category over \mathbf{X} has a fibre-small Δ -universal initial completion.*

PROOF: The Δ -universal initial completion has been described in the proof of Theorem 4 of [E] (for the case of $\Gamma =$ all sources and $P =$ forgetful functor of \mathbf{K}) as the category of all Δ -complete sources in \mathbf{K} . It is our task to show that this category is fibre-small (thus, legitimate).

For each Δ -complete source $\sigma = (X \xrightarrow{f_i} |S_i|)_{i \in I}$ and each i we factorize $f_i = m_i \cdot e_i$ ($m_i \in \mathbf{M}$, $e_i \in \mathbf{E}$) and denote by $m_i : S'_i \rightarrow S_i$ the initial morphism induced by m_i (which exists since \mathbf{K} is hereditary). From the fact that Δ contains $\{m_i\}$ it follows that $X \xrightarrow{e_i} |S'_i|$ is a member of the (Δ -complete) source σ . Thus,

$$\sigma' = (X \xrightarrow{e_i} |S'_i|)_{i \in I}$$


is a subsource of σ . This subsource fully determines σ (in other words, given two Δ -complete sources $\sigma_1 \neq \sigma_2$, it follows that $\sigma'_1 \neq \sigma'_2$). In fact, from the Δ -completeness

it follows that for each $i \in I$ and each morphism $h : S'_i \rightarrow S$ in \mathbf{K} we have $X \xrightarrow{he_i} |S|$ in σ ; consequently, σ is precisely the source of all composites of the members of σ' with morphisms of \mathbf{K} . Thus, to prove the fibre-smallness, it is sufficient to observe that for each object X of \mathbf{X} , all the possible sources σ' form a set. In fact, since \mathbf{X} is \mathbf{E} -co-wellpowered, we have a set of representatives for all e_i 's, and for each such a representative $e_i : X \rightarrow Y$ we have (since \mathbf{K} is fibre-small) only a set of possible objects S'_i with $|S'_i| = Y$. Thus, there exists a set A of representative structured maps $e_i : X \rightarrow |S'_i|$, and the collection of all Δ -complete sources with the domain X can, obviously, be coded by the set of all subsets of A . \square

Remark. For constructs \mathbf{K} , i.e. concrete categories over \mathbf{Set} , the Δ -universal initial completion \mathbf{K}^* can be described as follows.

Objects on the underlying set X are all collections σ of \mathbf{K} -objects K with the underlying sets $|K| = X / \sim$ (where \sim is an equivalence relation on X) such that:

- (1) If $K \in \sigma$ then $K' \in \sigma$ whenever $|K'| = X / \sim$ and the canonical map $c : X \rightarrow |K'|$ of $|K'|$ (assigning to each $x \in X$ the equivalence class of x) factorizes as the canonical map of $|K|$ followed by a \mathbf{K} -morphism $K \rightarrow K'$;
- (2) For each initial Δ -source $(S \xrightarrow{f_i} S_i)_{i \in I}$ in \mathbf{K}



and each map $h : X \rightarrow |S|$ such that every $f_i h$ factorizes as the canonical map of $|K_i|$ for some $K_i \in \sigma$ followed by a \mathbf{K} -morphism $h_i : K_i \rightarrow S_i$ it follows that σ contains the initial lift of the inclusion $X / \ker h \hookrightarrow |S|$.

Morphisms from σ to σ' are maps $f : |\sigma| \rightarrow |\sigma'|$ such that for each $K' \in \sigma'$ (with the canonical map $c : |\sigma'| \rightarrow |K'|$) σ contains the initial lift of the inclusion $|\sigma| / \sim \hookrightarrow |K|$, where \sim is the kernel equivalence of c.f.

II. A completion of the category of metric spaces.

Proposition. *The category \mathbf{Unif} is a Δ_{fm} -universal initial completion of the category \mathbf{Met} .*

PROOF: We apply the general result above to the special case of $\Delta = \Delta_{fm}$ and $\mathbf{K} = \mathbf{Met}$. The objects of Δ_{fm} -universal initial completion \mathbf{K}^* with the underlying set X are collections σ of metric spaces $K = (X/\sim, d_K)$, where \sim is an equivalence on X , such that (1) and (2) are satisfied. Each K induces a pseudometric d_K^* on the set X by $d_K^*(x, y) = d_K([x], [y])$ (where $x \mapsto [x]$ denotes the canonical map). The conditions (1) and (2) guarantee that the resulting set of pseudometrics on X forms a uniformity (defined as a collection of pseudometrics). Conversely, for each uniformity on X and each pseudometric d of that uniformity we have a corresponding metric space on the set X/\sim where $x \sim y$ means that $d(x, y) = 0$. The axioms of a uniformity guarantee that the resulting set of metric spaces is an object of \mathbf{K}^* . The morphisms of \mathbf{K}^* correspond precisely to the uniformly continuous maps. Thus, \mathbf{Unif} is a Δ_{fm} -universal initial completion of \mathbf{Met} . \square

Remark. Since \mathbf{Met} is, obviously, closed under countable initial sources in \mathbf{Met} , we can also say that for $\Delta =$ all countable sources, \mathbf{Unif} is a Δ -universal initial completion of \mathbf{Met} . However, the universal completion ($\Delta =$ all sources) is different:

Example of an initial source in \mathbf{Met} which is not initial in \mathbf{Unif} .

Let δ be the discrete metric (with value 1) on the set N of natural numbers. Let \mathbf{F} be a free ultrafilter on N containing the set E of even numbers. For each $F \in \mathbf{F}$ let δ_F be the following metric on N :

$$\delta_F(x, y) = \begin{cases} \frac{1}{n} & \text{if } \{x, y\} = \{2n - 1, 2n\} \text{ with } 2n \in F \\ 0 & \text{if } x = y \\ 1 & \text{else.} \end{cases}$$

As proved in [PRRS], the set of all $\delta_F, F \in \mathbf{F}$, is a base of a uniformity σ on N which is an atom in the fibre of N (i.e. the only strictly finer uniformity is that induced by δ). Consequently, the following source

$$((N, \sigma) \xrightarrow{id} (N, \delta_F))_{F \in \mathbf{F}}$$

is initial in \mathbf{Unif} . We will show that, nevertheless,

$$((N, \delta) \xrightarrow{id} (N, \delta_F))_{F \in \mathbf{F}}$$

is initial in \mathbf{Met} . In fact, let (X, δ_0) be a metric space, and let $f : X \rightarrow N$ be a map such that $f : (X, \delta_0) \rightarrow (N, \delta_F)$ is uniformly continuous for each $F \in \mathbf{F}$. We will prove that the image of f , considered as a uniform subspace of (N, σ) , is discrete — thus, $f : (X, \delta_0) \rightarrow (N, \delta)$ is uniformly continuous.

Suppose that image of f is not discrete. Then it is isomorphic to (N, σ) . [In fact, every uniform subspace A of (N, σ) is isomorphic to (N, σ) or is discrete, according to whether the set $\{2n \mid n \in N, 2n \in A, 2n + 1 \in A\}$ is a member of \mathbf{F} or not.] Thus, we can assume that f is surjective. Since σ is an atom, it follows that f is a final morphism. However, (N, σ) is not a quotient of a metric space, since it is not generated by a single pseudometric — this is a contradiction.

REFERENCES

- [A] Adámek J., *Theory of Mathematical Structures*, Reidel Publ. Comp., Dordrecht, 1983.
- [AHS] Adámek J., Herrlich H., Strecker G.E., *Least and largest initial completion*, Comment. Math. Univ. Carolinae **20** (1979), 43–75.
- [C] Čech E., *Topological Spaces*, Academia Prague, 1966.
- [E] Ehlersmann A.E., *Partial completions of concrete functors*, Cahiers Topo. Géom. Diff. **22** (1981), 315–328.
- [H] Herrlich H., *Initial completions*, Math. Z. **150** (1976), 101–110.
- [PRRS] Pelant J., Reiterman J., Rödl V., Simon P., *Ultrafilters on ω and atoms in the lattice of uniformities I*, Topology and Appl. **30** (1988), 1–17.

FACULTY OF ELECTRICAL ENGINEERING, TECHNICAL UNIVERSITY PRAGUE, TECHNICKÁ 2,
PRAHA 6, CZECHOSLOVAKIA

(Received September 25, 1992)