

Angelo Bella; Camillo Costantini
On the Novak number of a hyperspace

Commentationes Mathematicae Universitatis Carolinae, Vol. 33 (1992), No. 4, 695--698

Persistent URL: <http://dml.cz/dmlcz/118541>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1992

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On the Novak number of a hyperspace

ANGELO BELLA, CAMILLO COSTANTINI

Abstract. An estimate for the Novak number of a hyperspace with the Vietoris topology is given. As a consequence it is shown that this cardinal function can decrease passing from a space to its hyperspace.

Keywords: hyperspace, Vietoris topology, Novak number, netweight

Classification: 54A25, 54B20

The motivation for this paper comes from a question posed in [3]. There it was proved (relatively to the locally finite topology, but the same reasoning applies to the Vietoris topology) that for any dense in itself T_1 space X the Novak number of $\exp(X)$ is not greater than that of X . The point left open was whether the two cardinal numbers can actually be different.

Here we present an estimate for the Novak number of a hyperspace in terms of the netweight of the base space. Using this theorem we give a couple of examples showing that the Novak number can actually decrease passing to the hyperspace. This fact is rather surprising considering the behaviour of practically all other cardinal functions.

Given a topological space X , the hyperspace $\exp(X)$ is the set of all non empty closed subsets of X .

If \mathcal{S} is a family of subsets of X then the symbol $\langle \mathcal{S} \rangle$ denotes the set of all $A \in \exp(X)$ satisfying $A \subset \cup \mathcal{S}$ and $A \cap S \neq \emptyset$ for every $S \in \mathcal{S}$.

The Vietoris topology on $\exp(X)$ is defined by taking as a base all the sets of the form $\langle U_1, \dots, U_n \rangle$, where U_1, \dots, U_n are open subsets of X . For more information on the Vietoris topology the reader is referred to [5].

Notice that if C_1, \dots, C_n are closed subsets of X then the set $\langle C_1, \dots, C_n \rangle = \exp(X) \setminus (\langle X \setminus C_1 \rangle \cup \dots \cup \langle X \setminus C_n \rangle \cup \langle X, X \setminus (C_1 \cup \dots \cup C_n) \rangle)$ is closed in $\exp(X)$.

A network \mathcal{B} for the topological space X is a family of subsets having the property that for any open set $U \subset X$ and any $x \in U$ there exists a member $B \in \mathcal{B}$ such that $x \in B \subset U$.

The netweight of the space X , denoted by $nw(X)$, is the smallest cardinality of a network for X .

Given a dense in itself T_1 space X , the Novak number of X , denoted by $n(X)$, is the smallest cardinality of a cover of X consisting of nowhere dense sets.

More details on the Novak number can be found in [1] and [2] and the bibliography listed there.

Theorem 1. *If X is a dense in itself regular T_1 space then $n(\text{exp}(X)) \leq nw(X)^{\aleph_0}$.*

PROOF: Let \mathcal{B} be a network of X satisfying $|\mathcal{B}| = nw(X)$. As the space X is regular, we can assume that \mathcal{B} consists of closed sets. Denote by \mathcal{B}_1 the collection of all countable infinite subsets of \mathcal{B} consisting of pairwise disjoint elements. For any $\mathcal{C} = \{C_1, \dots, C_n, \dots\} \in \mathcal{B}_1$ let

$$F_{\mathcal{C}} = \bigcap_{n \in \mathbb{N}^+} \langle X, C_n \rangle.$$

Since every C_n is closed, it follows that also $F_{\mathcal{C}}$ is closed. Moreover, it is clear that every point in $F_{\mathcal{C}}$ is an infinite subset of X . On the other hand, every basic open set in $\text{exp}(X)$ contains finite subsets of X and therefore it follows that each $F_{\mathcal{C}}$ is nowhere dense. We claim that every infinite closed subset A of X is contained in some $F_{\mathcal{C}}$. To see this, let us begin by taking two disjoint elements $C', C'' \in \mathcal{B}$ such that $C' \cap A \neq \emptyset \neq C'' \cap A$. At least one of these two sets, say C' , satisfies $|A \setminus C'| \geq \aleph_0$. Let $C_1 = C'$ and suppose we have already chosen pairwise disjoint sets $C_1, \dots, C_n \in \mathcal{B}$ in such a way that $(*) C_i \cap A \neq \emptyset$ for $i \leq n$ and $|A \setminus (C_1 \cup \dots \cup C_n)| \geq \aleph_0$. Then we proceed by induction selecting $C_{n+1} \in \mathcal{B}$ disjoint from C_1, \dots, C_n and satisfying $(*)$. Letting $\mathcal{C} = \{C_1, \dots, C_n, \dots\}$ it is clear that $A \in F_{\mathcal{C}}$. Now let F_n be the subset of $\text{exp}(X)$ consisting of all subsets of X having cardinality not bigger than n . Since X is dense in itself and T_2 , it is not difficult to see that F_n is closed and nowhere dense in $\text{exp}(X)$. To finish, observe that $\{F_n : n \in \mathbb{N}^+\} \cup \{F_{\mathcal{C}} : \mathcal{C} \in \mathcal{B}_1\}$ is a nowhere dense cover of $\text{exp}(X)$ of cardinality not exceeding $nw(X)^{\aleph_0}$. \square

In order to obtain our first example, we recall the construction of a certain linearly ordered topological group.

For any ordinal ν denote by \mathfrak{R}^ν the set of all functions $f : \nu \rightarrow \mathfrak{R}$ ordered lexicographically, that is $f < g$ if and only if $f \neq g$ and $f(\alpha) < g(\alpha)$, where $\alpha = \min\{\beta : f(\beta) \neq g(\beta)\}$. The order so defined is actually a linear order and \mathfrak{R}^ν can be equipped with the standard interval topology. If, moreover, we define $f + g$ by the rule $(f + g)(\alpha) = f(\alpha) + g(\alpha)$ then \mathfrak{R}^ν becomes a linearly ordered topological abelian group.

For any $\alpha \in \nu$ denote by ε_α the element of \mathfrak{R}^ν defined by $\varepsilon_\alpha(\alpha) = 1$ and $\varepsilon_\alpha(\beta) = 0$ if $\beta \neq \alpha$.

Let $\mathfrak{R}^{<\nu} = \bigcup_{\alpha \in \nu} \mathfrak{R}^\alpha$ and for any $\varphi \in \mathfrak{R}^{<\nu}$ denote by $\|\varphi\|$ the ordinal α such that $\varphi \in \mathfrak{R}^\alpha$. Let $[\varphi] = \{f \in \mathfrak{R}^\nu : f \upharpoonright \alpha = \varphi\}$. Observe that $[\psi] \subset [\varphi]$ if and only if $\varphi \subset \psi$.

Each $[\varphi]$ is open in \mathfrak{R}^ν , in fact if $\varphi \in \mathfrak{R}^\alpha$ and $f \in [\varphi]$ then the interval $(f - \varepsilon_{\alpha+1}, f + \varepsilon_{\alpha+1})$ is contained in $[\varphi]$. Furthermore, the collection of all sets of the form $[\varphi]$ is a base for the topology of \mathfrak{R}^ν . To see this, fix an interval (f, g) and an element $h \in (f, g)$ and let $\alpha_1 = \min\{\beta : f(\beta) \neq h(\beta)\}$ and $\alpha_2 = \min\{\beta : h(\beta) \neq g(\beta)\}$. If $\alpha = \max\{\alpha_1, \alpha_2\} + 1$ then it is easily seen that $h \in [h \upharpoonright \alpha] \subset (f, g)$.

The next proposition is somewhat related to a result of Sikorski ([6, 4.15]).

Proposition 1. *If ν is a regular cardinal then $n(\mathfrak{R}^\nu) > \nu$.*

PROOF: It is enough to show that any family $\{A_\alpha : \alpha \in \nu\}$ of dense open subsets of \mathfrak{R}^ν has a non empty intersection. We construct by induction the sequence

$\{\varphi_\alpha : \alpha \in \nu\} \subset \mathfrak{R}^{<\nu}$ satisfying the condition

$$[\varphi_\alpha] \subset A_\alpha \cap (\bigcap_{\beta \in \alpha} [\varphi_\beta]).$$

Assume that the family $\{\varphi_\beta : \beta \in \alpha\}$ has already been constructed. Since ν is regular, the set $\{|\varphi_\beta| : \beta \in \alpha\}$ is bounded in ν . Thus the function $\psi = \bigcup\{\varphi_\beta : \beta \in \alpha\}$ is a member of $\mathfrak{R}^{<\nu}$. To finish the induction, select $\varphi_\alpha \in \mathfrak{R}^{<\nu}$ in such a way that $[\varphi_\alpha] \subset A_\alpha \cap [\psi]$. Now let $\varphi = \bigcup\{\varphi_\alpha : \alpha \in \nu\}$. If φ is defined on all ν then $\varphi \in \mathfrak{R}^\nu$ and $\varphi \in \bigcap_{\alpha \in \nu} A_\alpha$. If, on the other hand, $\|\varphi\| < \nu$ then any $f \in [\varphi]$ belongs to $\bigcap_{\alpha \in \nu} A_\alpha$. \square

Recall that assuming Martin's axiom (MA) the cardinality of the continuum \mathfrak{c} is regular and, for every $\kappa < \mathfrak{c}$, $2^\kappa \leq \mathfrak{c}$ (see [4, Section 2.2]). Taking this into account, we see that the space $\mathfrak{R}^\mathfrak{c}$ has a base (and a fortiori a network) of cardinality not exceeding $|\mathfrak{R}^{<\mathfrak{c}}| = \sum_{\alpha \in \mathfrak{c}} 2^{|\alpha|} = \mathfrak{c}$. Thus from Theorem 1 and Proposition 1 we get:

Corollary 1 (MA). *There exists a linearly ordered topological group X , namely $\mathfrak{R}^\mathfrak{c}$, for which $n(\text{exp}(X)) < n(X)$.*

Under more complicate set-theoretic assumptions, it is possible to find a compact space for which the Novak number decreases passing to the hyperspace. Indeed Theorem 5.2 in [1] describes two models of ZFC in which the Novak number of N^* , the Čech-Stone remainder of N , is greater than \mathfrak{c} . Taking into account that N^* is a compact space of netweight \mathfrak{c} , another application of Theorem 1 gives:

Corollary 2. *It is consistent with ZFC the existence of a compact space X , namely N^* , for which $n(\text{exp}(X)) < n(X)$.*

Observe that, since for any compact space the locally finite topology coincides with the Vietoris topology, Corollary 2 also furnishes a direct answer to the question posed in [3].

We do not know whether the inequality $n(X) \leq 2^{n(\text{exp}(X))}$ holds for every dense in itself T_1 space X . Certainly, however, it cannot be improved. In fact, in one of the models described in [1], it is true that $n(N^*) = 2^\mathfrak{c} = 2^{n(\text{exp}(N^*))}$.

REFERENCES

- [1] Balcar B., Pelant J., Simon P., *The space of ultrafilters on N covered by nowhere dense sets*, Fund. Math. **110** n.1 (1980), 11–24.
- [2] Bella A., *Some remarks on the Novak number*, Proceedings of the 6th Prague Topological Symposium – General Topology and its Relations to Modern Analysis and Algebra VI – Z. Frolík ed., Heldermann Verlag, Berlin (1988), pp. 43–48.
- [3] ———, *Some cardinality properties of a hyperspace with the locally finite topology*, Proc. Amer. Math. Soc. **104** n. 4 (1988), 1274–1278.
- [4] Kunen K., *Set Theory. An introduction to independence proofs*, North Holland Pub. Co., 1980.

- [5] Michael E., *Topologies on spaces of subsets*, Trans. Amer. Math. Soc. **71** (1951), 152–182.
- [6] Sikorski R., *Spaces of high power*, Fund. Math. **37** (1950), 125–136.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MESSINA, CONTRADA PAPARDO, SALITA SPERONE 31, 98166 MESSINA, ITALY

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MILANO, VIA C. SALDINI 50, 20133 MILANO, ITALY

(Received May 13, 1992)